

Conclusion Here we have shown for the first time that vasopressin and oxytocin have direct contractile effects on human isolated stomach muscle. The effective concentrations of vasopressin are within the range induced by nausea in humans. This indicates a potential direct role of vasopressin in signalling the induction of nausea in humans.

Competing interests None declared.

REFERENCES

1. **Stern RM**, Koch KL, Andrews PL. *Nausea: Mechanisms and Management*. Oxford Univ Press, 2011.
2. **Sanger GJ**, Holbrook JD, Andrews PL. The translational value of rodent gastrointestinal functions: a cautionary tale. *Trends Pharmacological Sci* 2011;**32**:402–9.

PWE-004 A STUDY OF FAECAL VOLATILE ORGANIC COMPOUNDS METABOLOME IN HEALTHY POPULATION ACROSS THE COUNTRIES

doi:10.1136/gutjnl-2012-302514d.4

¹I Ahmed,* ²V De Preter, ³K Rioux, ⁴N Ratcliffe, ⁵C S Probert. ¹Department of Gastroenterology, University of Bristol, Bristol, UK; ²Department of Gastroenterology, Translational Research Centre for Gastrointestinal Disorders, Leuven, Belgium; ³Department of Gastroenterology, University of Calgary, Calgary, Canada; ⁴Department of Department of sensors and Technology, University of West of England, Bristol, UK; ⁵Department of Gastroenterology, University of Liverpool, Liverpool, UK

Introduction Faecal biomarkers are emerging non-invasive tools for diagnosing gastrointestinal disorders. Faecal volatile organic compounds (VOCs) have been studied more recently in clinical diagnosis. Pattern of faecal VOCs in healthy population may provide basis for understanding changes in disease conditions. The VOCs within the metabolome may be different across the countries due to differences in dietary habits and environmental conditions and may have implications in developing their clinical utility.

Methods We aim to study the faecal VOCs of the healthy population from three different countries that is, England, Belgium and Canada. A total of 159 health volunteers (English=109, F=69), (Belgium=20, F=14), (Canada=30, F=17) donated faecal samples. Fresh samples were aliquoted in 18 mls sealed vials. VOCs were extracted using solid phase micro extraction and were analysed using gas chromatography–mass spectrometry. VOCs were identified using NIST library search comparing their fragment pattern.

Results A total of 232 VOCs were identified. Using binary data (presence or absence of VOCs), univariate analysis was used to identify those VOCs which were statistically significant ($p < 0.05$) in discerning differences between the three population groups. Alcohols, ketones and esters were predominantly associated with English volunteers compared to both Canadian and Belgium volunteers while aldehydes and alkenes were predominantly detected VOCs in the Canadian and Belgium groups respectively. A multivariate discriminant function analysis utilising these VOCs was able to differentiate three groups with a sensitivity of 96% and specificity of 90%.

Conclusion The observed differences in the faecal VOCs metabolites of the healthy population in different countries may provide important basis in the clinical utility of faecal biomarkers. It may also provide information in studying the differences in disease prevalence and behaviour in different countries. Further studies are warranted to explore this area.

Competing interests None declared.

REFERENCES

1. **Probert CS**, Ahmed I, Khalid T, et al. Volatile organic compounds as diagnostic biomarkers in gastrointestinal and liver diseases. *J Gastrointest Liver Dis* 2009;**18**:337–43.
2. **Angriman I**, Scarpa M, D'Inca R, et al. Enzymes in feces: useful markers of chronic inflammatory bowel disease. *Clin Chim Acta* 2007;**381**:63–8.

PWE-005 HIGH RESOLUTION ANORECTAL MANOMETRY: FIRST STUDY ESTABLISHING NORMAL VALUES IN HEALTHY VOLUNTEERS

doi:10.1136/gutjnl-2012-302514d.5

¹J M Burke,* ¹W Jackson, ¹M M Withers, ¹H O'Grady, ²G S Duthie. ¹GI Physiology, Hull and East Yorkshire Hospitals NHS Trust, Cottingham, UK; ²Hull York Medical School, Hull, UK

Introduction High Resolution Anorectal Manometry (HRAM) combined with interpretive software allows for the interpolation of manometric recordings into highly detailed topographical plots of intraluminal pressure events. HRAM has previously been shown to correlate highly with conventional water perfused manometry measurements.¹ This preliminary study is the first report establishing HRAM pressures in healthy volunteers. The advantages of the detection of pressure changes over a longer length of the anal canal have already been shown to improve accuracy and the detection of abnormalities in the anorectum.²

Methods HRAM was performed using the Medical Measurement System (Enschede, Netherlands) consisting of an 8-channel HRAM catheter with sensors spaced at 0.8 cm intervals. Pressure data are displayed in topographic form using Medical Measurement System analysis software that is integrated into the system. Measurements of anal sphincter pressure at rest, cough, during voluntary squeeze, endurance squeeze and pushdown were evaluated. Volunteers also completed a questionnaire which provided a Wexner score.

Results A total of 20 healthy volunteers (11 Female, 9 Male) with a mean age of 40 (range 19–60) constituted the study population. The Wexner scores ranged from 0 to 1 (median 0).

Conclusion These preliminary measurements of HRAM pressures in healthy volunteers could serve as a valuable resource of normative data when performing HRAM studies in disease specific groups such as incontinence and constipation.

Abstract PWE-005 Table 1

Anal sphincter	Range	Median
Resting pressure	30–163 cm H ₂ O	109 cm H ₂ O
Cough pressure increase	39–305 cm H ₂ O	143 cm H ₂ O
Voluntary squeeze pressure	50–922 cm H ₂ O	275 cm H ₂ O
Endurance squeeze time	18–125 s	52 s
% of relaxation during pushdown	0–42% (17/20 relaxed)	14%

Competing interests None declared.

REFERENCES

1. **Jones MP**, Post J, Crowell MD. High-resolution manometry in the evaluation of anorectal disorders: a simultaneous comparison with water-perfused manometry. *Am J Gastroenterol* 2007;**102**:850–5.
2. **Rao SS**. Advances in diagnostic assessment of fecal incontinence and dyssynergic defecation *Clin Gastroenterol Hepatol* 2010;**8**:910–19.

PWE-006 DIAGNOSTIC YIELD AND CLINICAL OUTCOME FOR DEFAECATING PROCTOGRAPHY AND ANORECTAL MANOMETRY IN PATIENTS WITH CHRONIC CONSTIPATION

doi:10.1136/gutjnl-2012-302514d.6

¹M Bhalme,* ²S Murugesan, ²V Jayasekeran, ²E Wriughtham, ²B Unsworth, ²A Mondino, ²J McLaughlin, ²P Paine. ¹North Manchester General Hospital, Manchester, UK; ²Salford Royal Foundation Trust, Salford, UK

Introduction Defaecating proctography (DFP) and anorectal manometry (ARM) are both used to investigate chronic

constipation but their relative clinical performance is unclear. Our aim was to investigate the diagnostic yield and clinical outcomes of DFP and ARM in chronic constipation.

Methods Patients who had undergone both DFP and ARM over a 3-year period were studied retrospectively. Demographics, treatment and clinical outcomes were recorded. The diagnosis was recorded as "mixed" if investigation showed evidence of both anismus and anatomical problems such as rectocele, intussusception or prolapse. The clinical outcome was defined as positive if the test resulted in treatment with symptomatic improvement, or resolution at follow-up. To determine whether there was a selection bias in those undergoing both DFP and ARM we additionally looked at the two groups having solely DFP or ARM from the same period.

Results DFP and ARM group: 43 patients (40 female, 58% surgical referrals; age range 17–85 years; median 46) underwent both DFP and ARM. The diagnostic yield for DFP was higher at 98% (anismus 44%, anatomical 40%, mixed 14%; normal 2%) vs 47% for ARM (anismus 26%, mixed 21%; normal 53%). There was diagnostic concordance in only 11 (26%), partial concordance in 9 (21%) and discordance in 23 (53%) patients. Although the diagnostic yield of DFP was much greater than ARM in this combined group, both tests led to similar positive outcomes regardless (47% in DFP vs 45% in ARM) when tests revealed a pathology. Single investigation groups: 10 patients had DFP alone (8 female, 60% surgical referrals; age range 22–73 years, median 55) with a diagnostic yield of 90% (anismus 30%, anatomical 50%, mixed 10%; normal 10%). The positive outcome in those with a detectable pathology was 33%. 15 patients had ARM alone (14 female, 27% surgical referrals; age range 19–75 years, median 50) with a diagnostic yield of 67% (anismus; 33% normal). The positive outcome in those with a detectable pathology was 70%.

Conclusion DFP had a higher diagnostic yield than ARM, but concordance was poor. Greater diagnostic yield did not translate into more positive clinical outcomes either. The clinical impact of additional DFP-based diagnoses is therefore questionable. The single test cohort data suggest that patients having DFP alone are a different clinical population from those who accessed both tests, since diagnostic yields and clinical outcomes were higher for ARM alone. The latter group were predominantly medical gastroenterology referrals. Further study is required to design optimal investigation strategies for chronic constipation.

Competing interests None declared.

PWE-007

PARACRINE PROSTAGLANDIN-E SIGNALLING MODULATES CANINE GASTRIC EPITHELIAL CELL MIGRATION

doi:10.1136/gutjnl-2012-302514d.7

¹R Hollins,* ²A I Freeman, ³P J Cripps, ³A D Radford, ¹P J M Noble. ¹*Institute of Translational Medicine, University of Liverpool, Liverpool, UK;* ²*School of Veterinary Science, University of Liverpool, Liverpool, UK;* ³*Institute of Infection and Global Health, University of Liverpool, Liverpool, UK*

Introduction Gastric ulceration is a limiting complication of therapy with cyclooxygenase (COX) antagonists, widely used anti-inflammatory/analgesic drugs in both humans and dogs. COX-derived prostaglandin E (PGE) has an important role in gastric defense and cytoprotection via promotion of blood flow and mucus secretion and inhibition of gastric acid secretion. Given the importance of gastric epithelial cell migration in reestablishing epithelial integrity following gastric damage, we have investigated whether paracrine PGE signalling has a role in the modulation of gastric epithelial cell migration.

Methods In order to retain paracrine signalling between different cell types, we isolated intact gastric glands via collagenase digestion of

canine gastric mucosal tissue. Isolated glands spread *in vitro* to form islands of cells. The rate of gland spreading over 48h was measured as a surrogate for cell migration speed. Lamellipodia protrusion was analysed as an index of spreading activity. A value for lamellopodia area was calculated by measuring spread area minus area bounded by nuclei of cells at the edge of spread glands. Spread glands, when serum-starved, exhibit a reduction in area. We added both a selective and a non-selective COX antagonist and PGE₂ to serum-starved glands to assess their effects on migration. All treatments were added blindly to eliminate bias. Statistical significance was assessed using univariate analysis of variance. Expression of COX-2 and PGE receptors (EP-3 and EP-4) was assessed by RT-PCR and immunohistochemistry.

Results RT-PCR confirmed COX-2, EP-3 and EP-4 expression in our samples. COX-2 immunoreactivity was present in the majority of gland cells. The COX 1/2 antagonist indomethacin (50 µM) decreased spreading (0.85-fold, *p*<0.05, *n*=5–9 for all experiments) and lamellipodia area (0.3-fold, *p*<0.05). The COX-2 selective antagonist NS-398 (10 µM) caused similar decreases to indomethacin (0.8-fold and 0.65-fold respectively, *p*<0.05). PGE (1 µM) prevented a 0.7-fold reduction in island area elicited by incubation in serum free medium (*p*<0.05).

Conclusion This data shows a role for COX-2 derived PGE in the promotion of gastric cell migration and cellular lamellipodia formation. A reduction of mucosal PGE via COX-2 antagonism may therefore inhibit gastric epithelial cell migration contributing to COX-antagonist elicited gastric ulceration in both humans and dogs.

Competing interests None declared.

Oesophageal II

PWE-008

DO STATINS PREVENT THE HISTOLOGICAL SUBTYPES OF OESOPHAGEAL CANCER? PROSPECTIVE DATA FROM THE UK GENERAL PRACTICE RESEARCH DATABASE (GPRD)

doi:10.1136/gutjnl-2012-302514d.8

¹H Y Bhutta, ^{2,3}L Alexandre,* ³A Clark, ⁴S Holt, ^{1,3}M Lewis, ^{2,3}A Hart. ¹*Department of General Surgery, Norfolk and Norwich Hospital, Norwich, UK;* ²*Department of Gastroenterology, Norfolk and Norwich Hospital, Norwich, UK;* ³*Norwich Medical School, University of East Anglia, Norwich, UK;* ⁴*Roundwell Medical Centre, Norwich, UK*

Introduction The incidence of oesophageal adenocarcinoma (OAC) has risen dramatically in the Western world and is associated with a poor prognosis. Statins show anti-cancer properties in experimental work with OAC cell lines for example reduced cell proliferation, increased apoptosis. This study aimed to investigate if statins are negatively associated with the development of two different histological subtypes of oesophageal cancer, OAC and oesophageal squamous cell cancer (OSCC), in a prospective cohort study.

Methods The cohort was over 4 million people in the General Practice Research Database (GPRD), a UK database of 488 nationwide general practices. Information is recorded on medication use prior to development of other illnesses, including cancers. Statin use was defined as a prescription for a minimum of 10 months preceding diagnosis of oesophageal cancer. Approximately half the GP practices in the GPRD are linked to the NHS cancer registry, allowing identification and sub-classification of histologically confirmed cases of OAC and OSCC. Each case was matched with four controls and conditional logistic regression estimated the OR plus 95% CIs for the development of each type of cancer, adjusted for diabetes, BMI, smoking, aspirin, PPIs and drugs that relax the lower oesophageal sphincter.

Results 581 histologically confirmed cases of OAC (77.8% men, mean age 70.7 years, SD=11.4) and 332 cases of OSCC (38.6% men,