



ORIGINAL ARTICLE

Neurotensin—regulated miR-133 α is involved in proinflammatory signalling in human colonic epithelial cells and in experimental colitis

Ivy Ka Man Law,¹ Kyriaki Bakirtzi,¹ Christos Polytarchou,² Angelos Oikonomopoulos,¹ Daniel Hommes,¹ Dimitrios Iliopoulos,² Charalabos Pothoulakis¹

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/gutjnl-2014-307329>).

¹Inflammatory Bowel Disease Center, Division of Digestive Diseases, David Geffen School of Medicine, UCLA, Los Angeles, California, USA

²Center for Systems Biomedicine, Division of Digestive Diseases, David Geffen School of Medicine, UCLA, Los Angeles, California, USA

Correspondence to

Professor Charalabos Pothoulakis, Inflammatory Bowel Disease Center, Division of Digestive Diseases, David Geffen School of Medicine, University of California at Los Angeles, 675 Charles E Young Dr, South MRL building 1240, Los Angeles, CA 90095, USA; cpothoulakis@mednet.ucla.edu

Received 26 March 2014

Revised 2 July 2014

Accepted 22 July 2014

ABSTRACT

Objective Neurotensin (NT) mediates colonic inflammation through its receptor neurotensin receptor 1 (NTR1). NT stimulates miR-133 α expression in colonic epithelial cells. We investigated the role of miR-133 α in NT-associated colonic inflammation in vitro and in vivo.

Design miR-133 α and aftiphilin (AFTP) levels were measured by quantitative PCR. Antisense (as)-miR-133 α was administrated intracolonically prior to induction of 2, 4, 6-trinitrobenzene sulfonic acid (TNBS)-induced colitis and dextran sodium sulfate (DSS)-induced colitis. The effect of AFTP was examined by gene silencing in vitro.

Results NT increased miR-133 α levels in NCM-460 overexpressing NTR1 (NCM460-NTR1) and HCT-116 cells. NT-induced p38, ERK1/2, c-Jun, and NF- κ B activation, as well as IL-6, IL-8 and IL-1 β messenger RNA (mRNA) expression in NCM-460-NTR1 cells were reduced in miR-133 α -silenced cells, while overexpression of miR-133 α reversed these effects. MiR-133 α levels were increased in TNBS (2 day) and DSS (5 day) colitis, while NTR1 deficient DSS-exposed mice had reduced miR-133 α levels, compared to wild-type colitic mice. Intracolonic as-miR-133 α attenuated several parameters of colitis as well expression of proinflammatory mediators in the colonic mucosa. In silico search coupled with qPCR identified AFTP as a downstream target of miR-133 α , while NT decreased AFTP expression in NCM-460-NTR1 colonocytes. Gene silencing of AFTP enhanced NT-induced proinflammatory responses and AFTP levels were downregulated in experimental colitis. Levels of miR-133 α were significantly upregulated, while AFTP levels were downregulated in colonic biopsies of patients with ulcerative colitis compared to controls.

Conclusions NT-associated colitis and inflammatory signalling are regulated by miR-133 α -AFTP interactions. Targeting of miR-133 α or AFTP may represent a novel therapeutic approach in inflammatory bowel disease.

INTRODUCTION

Neurotensin (NT) is a neuropeptide expressed in the central nervous system and the intestine,^{1 2} while its high-affinity receptor, NT receptor 1 (NTR1)³ is expressed in neurones,⁴ colonic epithelial,^{2 5–7} immune cells,^{8 9} as well as different colon cancer cell lines.¹⁰ In the colon, NT/NTR1 signalling promotes inflammation in acute colitis animal models.^{2 5} Colonic NT and NTR1 expression increases during enterotoxin-mediated intestinal inflammation,² dextran sodium sulfate (DSS) colitis,^{6 11 12} and in the colon of patients with

Significance of this study

What is already known on this subject?

- Neurotensin (NT) mediates colonic inflammation through its high-affinity receptor (neurotensin receptor 1 (NTR1)).
- NT and NTR1 expressions are upregulated in inflamed colon tissues in experimental colitis and ulcerative colitis (UC).
- NT modulates differential expression of 38 microRNAs in colonocytes, among them miR-133 α .
- MicroRNA-133 α modulates signalling pathways related to proliferation, differentiation and hypertrophy in cancer cell lines and myocytes.

What are the new findings?

- miR-133 α modulates NT-induced MAP kinase, and NF- κ B activation, as well as transcription of proinflammatory cytokines in vitro.
- Reducing the levels of miR-133 α in human colonocytes and mouse colon attenuates proinflammatory signalling and development of experimental colitis in vitro and in vivo.
- Aftiphilin (AFTP) is a novel downstream target of NT-induced miR-133 α expression in human colonic epithelial cells involved in proinflammatory signalling pathways.
- Colon tissues from patients with UC and mice with colitis have higher miR-133 α but lower AFTP levels when compared with controls.

How might it impact on clinical practice in the foreseeable future?

- Inhibiting miR-133 α and restoring AFTP expression in inflammatory bowel disease (IBD) colon may represent promising approaches for the treatment of IBD.
- The reverse association of the levels of miR-133 α and AFTP together with increased expression of NTR1 in UC may represent novel biomarkers for UC.

ulcerative colitis (UC).⁶ Moreover, NTR1 deficiency in mice is associated with reduced weight loss and mortality in experimental colitis.^{5 13}

MicroRNAs (miRs) are short (19–25 nucleotides), single-stranded RNA molecules, acting as

To cite: Law IKM, Bakirtzi K, Polytarchou C, et al. Gut Published Online First: [please include Day Month Year] doi:10.1136/gutjnl-2014-307329

negative transcriptional regulators. They bind to the 3' untranslated regions (UTR) of transcripts¹⁴ and lead to messenger RNA (mRNA) degradation, or inhibition of translation into protein.¹⁵ Recently, several studies identified the differential miR expression in colonic mucosa in patients with UC,^{16–19} Crohn's Disease (CD)^{18 20} and experimental colitis.¹⁷ However, the role of individual miRs in inflammatory bowel disease (IBD) pathophysiology is still under investigation.

We have recently shown that NT/NTR1 coupling in human colonic epithelial NCM460 overexpressing NTR1 (NCM460-NTR1) cells induces differential expression of miRs,¹⁰ including miR-133α. Dysregulation of miR-133α and its target genes has been correlated with colorectal cancer development^{21–28} through regulating ERK^{29 30} and Akt^{21 31} activation, pathways also targeted by NT/NTR1 signalling in human colonic epithelial cells.^{7 10 32} Based on these considerations, we examined the role of NT-miR-133α interactions in colonic inflammation in vitro and in two experimental colitis models induced by trinitrobenzene sulfonic acid (TNBS) and DSS in vivo. We also performed studies to identify the target of NT-NTR1-modulated miR-133α that mediates these effects. Our results point to aftiphilin (AFTPH), as a novel downstream target for miR-133α, and present evidence for the association of miR-133α and its downstream target, AFTPH, in the pathophysiology of colitis.

MATERIALS AND METHODS

Cell culture and reagents

NCM460-NTR1 cells were generated from human colonic epithelial NCM460 cells (INCELL, San Antonio, Texas, USA) transduced with lentivirus particles as previously described¹⁰ and maintained in M3:D culture media (INCELL) supplemented with 10% fetal bovine serum (FBS, Life Technologies, Grand Island, New York, USA). Human embryonic kidney fibroblasts, HEK293, and colonic cancer HCT-116 cells were maintained in Eagle's Minimum Essential Medium (MEM, Life Technologies) and McCoy5a (ATCC, Manassas, Virginia, USA), respectively, and supplemented with 10% FBS. Lipofectamine 2000, lipofectamine RNAiMAX and OptiMEM were from Life Technologies. NT was from Bachem Americas (Torrance, California, USA). TNBS and SR48962 were from Sigma Aldrich (St Louis, Missouri, USA), and DSS was from MP Biomedicals (Santa Ana, California, USA). Goat anti-AFTPH (sc-167055), mouse antivillin and rabbit anti-β tubulin were from Santa Cruz Biotechnology (Santa Cruz, California, USA).

Transfection experiments

Small interfering RNA (siRNA) was purchased from Santa Cruz Biotechnology (Santa Cruz). NCM460-NTR1 cells were transfected with siRNA against AFTPH (si-AFTPH) using Lipofectamine RNAiMAX. All miRs were purchased from Life Technologies. For miR-133α silencing or overexpression, cells were transfected with antisense-miR-133α (as-miR-133α) or miR-133α precursor (miR-133α), respectively, using Lipofectamine RNAiMAX. Cells transfected with siRNA-A (si-control), antisense-control miR (as-miR-control) or miR-negative control precursor (miR-control) served as controls.

NT/NTR1-mediated phosphoprotein activation and cytokine production

NCM460-NTR1 cells were transfected with as-miR-133α, miR-133α and si-AFTPH and their respective controls. Cells were incubated in serum-free media overnight and stimulated with NT (100 nM). Cell lysates were collected for

phosphoprotein detection, and media were collected 6 h after stimulation for cytokine measurements. Custom designed Bio-Plex Pro Cell Signalling assay panel (Bio-Rad, Hercules, California, USA) and Bio-Plex Pro Human Cytokine 27-plex Assay (Bio-Rad) were used for phosphoprotein activation and cytokine production (except IL-8), respectively, according to manufacturer's instructions.

Animal models and establishment of experimental colitis models

Neurotensin receptor 1 knockout mouse model

Ntsr1^{tm1Dgen} (hereafter called NTR1KO) were purchased from Jackson Laboratories and bred in our facility. Animals were obtained as fifth generation backcross of 129 onto C57BL6/J. We performed one additional backcross before intercrossing animals as heterozygous NTR1KO crosses to generate littermate controls.

TNBS-induced colitis

C57BL6/J wild-type (WT) mice were purchased from Jackson Laboratories. TNBS was dissolved in 30% ethanol (5 mg/kg) was intracolonically administered (50 µL) to mice as previously described.^{5 33} Briefly, TNBS was slowly infused via a 1 mL syringe (Becton Dickinson, Laguna Hills, California, USA) fitted with a polyethylene cannula (Intramedic PE-20 tubing; Becton Dickinson). Colon tissues were collected 48 h after TNBS administration.

DSS-induced colitis

NTR1KO and its WT counterparts, or C57BL6/J WT animals, were treated with 5% DSS in drinking water as previously described,^{13 34} and colon tissues were collected 5 days after the start of the treatment.

In vivo NT infusion. NT was dissolved in PBS supplemented with 1% bovine serum albumin (BSA)³⁵ and was administered intracolonically (300 µg/kg) to C57BL6/J WT mice twice per day for 4 days. Control mice were infused with 1% BSA in PBS. Colon tissues were collected 5 days after NT administration.

In vivo knockdown of miR-133α

Endogenous expression of miR-133α in colon tissues of C57BL6/J WT mice was silenced by intracolonic administration of miRCURY LNA Inhibitor probe, *in vivo* against *mmu-miR-133a* (20 µg/mouse, Exiqon). Briefly, the appropriate amount of oligonucleotides against *mmu-miR-133a* and its respective control were resuspended in 100 µL Opti-MEM with 2 µL lipofectamine 2000 and administered intracolonically 24 h and 72 h prior to TNBS or DSS treatment.

MicroRNA *in situ* hybridisation

Colon tissues (3 cm from distal end) were obtained from WT mice 2 day after TNBS treatment. The tissues were immediately fixed with 4% paraformaldehyde, paraffin-embedded and sectioned (5 µm) for histological study. 5'-DIG and 3'-DIG labelled detection probes specific for mouse miR-133α (Cat. No. 39270-15) and miRCURY LNA microRNA ISH optimisation Kit (FFPE) were purchased from Exiqon, and the experiments were performed according to the manufacturer's instructions.

RNA expression studies in patient samples

Total RNAs from colon tissues obtained from patients with UC (n=12), CD (n=8) and normal subjects (n=9) were purchased from Origene (Rockville, Maryland, USA). Conversion of cDNA

of RNA samples was performed as described above, and levels of miR-133 α and AFTPH were determined by qPCR analysis.

Immunohistochemistry

Frozen tissue sections (5 μ m) obtained from chronic active patients with UC (n=3) and their normal control (n=3) were purchased from Origene. The tissues were fixed with 4% paraformaldehyde and blocked with 0.3% Triton X-100 in PBS (PBS-Triton) supplemented with 5% normal bovine serum. Appropriate antibodies were incubated with the tissue sections overnight at 4°C, washed with PBS-Triton and incubated with appropriate secondary antibodies (Santa Cruz Biotechnology). The tissues were imaged using Axio Imager.Z1 microscope (Carl Zeiss).

Please refer to online supplementary methods for site-directed mutagenesis, luciferase assay, NF- κ B p65 translocation, measurement of interleukin-8 (IL-8) production, immunoblot analysis, mRNA and miR expression analysis and microRNA profiling in blood samples.

Statistical analysis

All in vitro results derived from at least three sets of experiments, expressed as means \pm SD and analysed with Student t tests. Results from animal experiments and studies in human tissues were analysed with Student t tests and expressed as means \pm SEM. In all statistical comparisons, $p<0.05$ was used to indicate significant differences.

RESULTS

Increased miR-133 α levels in human colonic epithelial cells after NT exposure

NT exerts its effect on colonic epithelial cells^{2 5–7} through its high-affinity receptor NTR1.³ NTR1 overexpression is observed in colon cancer cell lines,^{10 36} colon tissues from mice with experimental colitis,^{6 11 12} intestinal tumors³⁷ and in tissue biopsies from patients with UC.⁶ In a microarray analysis, we previously showed that NT/NTR1 coupling in human colonic epithelial NCM460-NTR1 increased miR-133 α expression.¹⁰ In our current study, we first verified this result by examining the levels of miR-133 α in human colonic epithelial NCM460-NTR1 cells and in colon cancer adenocarcinoma HCT-116 cells by qPCR. Consistent with our previous findings,¹⁰ levels of miR-133 α were upregulated after NT exposure by 2.6 ± 0.43 -fold ($p<0.01$, figure 1A) in NCM460-NTR1 cells, while in HCT-116 cells, levels of miR-133 α were increased by 1.5 ± 0.07 -fold ($p<0.01$, figure 1B) after addition of NT. Therefore, NT increases miR-133 α expression in two different human colonic epithelial cell lines.

NT induces proinflammatory signalling in human colonic epithelial cells via miR-133 α expression

NT coupling of NTR1 stimulates inflammatory responses through MAPK^{38 39}-dependent and NF- κ B^{7 40 41}-dependent pathways. Here, we examined whether miR-133 α is involved in NT-related proinflammatory signalling and cytokine expression in NCM460-NTR1 cells with an antisense approach using as-miR-133 α . First, we showed that transfection of as-miR-133 α blocked NT-induced miR-133 α overexpression in NCM460-NTR1 cells (figure 2A), compared to those transfected with control as-miR, validating this approach. Moreover, NT stimulation activated p38 and ERK1/2 signalling only in cells transfected with control as-miR, but not as-miR-133 α , while NT-induced c-Jun activation was significantly attenuated in as-miR-133 α -transfected cells (figure 2B, $p<0.05$). Moreover,




Figure 1 Levels of miR-133 α were upregulated after neurotensin (NT) exposure in human colonic epithelial cells. (A) Expression of miR-133 α in human colonic epithelial NCM460 cells after NT exposure was analysed by qPCR analysis. ** $p<0.01$, compared to vehicle control. (B) miR-133 α levels in human colon cancer HCT-116 cells after NT exposure were analysed by qPCR analysis. ** $p<0.01$, compared to vehicle control.


NT-induced NF- κ B p65 phosphorylation was blocked in as-miR-133 α -transfected cells (figure 2C), while NT-induced IL-6 and IL-8 mRNA (figure 2D, E, $p<0.05$) and IL-8 secretion were also significantly attenuated upon miR-133 α knock-down (figure 2F, $p<0.05$). Analysis of the cell culture media showed that NT stimulation increased the production of the proinflammatory IL-1 β , but reduced the production of the anti-inflammatory IL-1ra (figure 2G, $p<0.05$). Transfection of as-miR-133 α significantly reduced NT-induced IL-1 β production, while prevented the decrease in IL-1ra production (figure 2G). Together, these results suggest an important role for miR-133 α in the induction of proinflammatory signalling pathways and cytokine production in response to NTR1 activation in human colonocytes.

We further verified the proinflammatory role of NT and miR-133 α in NCM460 cells transfected with NTR1 and miR-133 α . Overexpression of miR-133 α further increased NT-induced ERK1/2, c-Jun (figure 3A, $p<0.05$) and NF- κ B p65 (figure 3B, $p<0.05$) activation. Additionally, overexpression of miR-133 α alone, in the absence of NT stimulation, promoted nuclear translocation of NF- κ B p65 (figure 3C, $p<0.05$) when compared to the cells expressing control miR. Furthermore, IL-8, IL-1 β and TNF- α mRNA was increased in cells overexpressing miR-133 α when compared to their controls (figure 3D, $p<0.05$). For cytokine production, IL-8 secretion was increased upon miR-133 α overexpression (figure 3E, $p<0.05$). Cells overexpressing miR-133 α had also significantly reduced IL-1ra production ($p<0.05$), while IL-1 β production was increased slightly, but not significantly (figure 3F). Together, these results strongly suggest an important role for miR-133 α in proinflammatory signalling in human colonocytes.

MiR-133 α overexpression in vivo is NT/NTR1-dependent

Since, as shown above, NT regulates miR-133 α in colonocytes in vitro, we next examined whether this effect can be recapitulated in vivo. To do this, C57/BL6 WT (WT) mice received intracolonically NT (300 μ g/kg).³⁵ After 4 days, mice were euthanased and colonic tissues were processed for RNA purification. qPCR analysis showed increased colonic miR-133 α levels (by 1.3 ± 0.04 -fold, figure 4A, $p<0.05$) in mice administered with NT, compared to intracolonic administration of vehicle. Because NT is an important mediator of acute colonic inflammation,^{2 13}

Figure 2 Downregulation of miR-133 α attenuated neurotensin (NT)-associated proinflammatory signalling in NCM460-neurotensin receptor 1 (NTR1) cells. (A) Levels of miR-133 α in NCM460-NTR1 cells upon NT stimulation after transfection of as-miR-133 α and its control were analysed by qPCR analysis. * $p<0.05$, compared to cells transfected with as-miR-control. (B) Levels of MAP kinase phosphorylation, including p38, ERK1/2 and c-Jun and (C) NF- κ B phosphorylation in NCM460-NTR1 cells transfected with as-miR-133 α and its control were analysed as stated in Methods. * $p<0.05$, compared to cells transfected with as-miR-control; ** $p<0.01$, compared to cells treated with NT. (D) IL-6 and (E) IL-8 levels in NCM460-NTR1 cells transfected with as-miR-133 α and its control was analysed by qPCR analysis. ** $p<0.01$, compared to cells transfected with as-miR-control and # $p<0.05$, compared with NT-treated cells. (F) IL-8 production in NCM460-NTR1 cells transfected with as-miR-133 α and its control was measured using ELISA. ** $p<0.01$, compared to cells treated with vehicle control; # $p<0.05$, compared with NT-treated cells. (G) IL-1 β and IL-1ra production in NCM460-NTR1 cells transfected with as-miR-133 α and its control was measured as stated in Methods. * $p<0.05$, compared to cells treated with vehicle control.



we also examined colonic levels of expression of miR-133 α in two colitis models. Acute colitis was induced in C57BL6/J WT mice by intracolonic administration of 5 mg/kg TNBS, or by addition of 5% DSS in their drinking water. qPCR analysis showed increased miR-133 α levels (by 3.2 ± 1.01 -fold, figure 4B, $p<0.05$) 2 days after TNBS administration, but decreased by 63% at day 7 (figure 4B, $p<0.05$). Mice receiving DSS showed a 3.6 ± 0.61 -fold increase in miR-133 α levels (figure 4B, $p<0.001$) at the end of the DSS treatment, compared to controls.

To study the mucosal distribution of miR-133 α in colon tissue during colitis, *in situ* hybridisation against miR-133 α in colon tissue sections obtained from TNBS-treated WT mice and their control counterparts was performed as described in Methods. Our result show that while miR-133 α expression was higher in inflamed colons, the majority of miR-133 α expression was localised in epithelial cells (figure 4C, arrows) and less in other cells of the lamina propria (figure 4C, arrowheads).


The direct relationship of endogenous NT/NTR1 signalling on miR-133 α expression in colon tissues during colitis development was further investigated in NTR1-deficient NTR1KO mice. NTR1KO mice and their WT counterparts were allowed access to 5% DSS in their drinking water, and after 5 days the mice were euthanased and miR-133 α levels were measured. As shown in figure 4D, colonic miR-133 α levels were increased in WT mice after DSS treatment (2.4 ± 0.32 -fold, $p<0.01$), but decreased in NTR1KO mice ($p<0.05$). Thus, increased

miR-133 α expression during acute colitis involves NT/NTR1 interactions.

Intracolonic silencing of miR-133 α attenuates the development of acute colitis

We next determined the role of miR-133 α in the development of experimental colitis. C57BL6/J WT mice were administered intracolonically two doses of as-miR-133 α or its control 24 h and 72 h prior to intracolonic administration of TNBS (5 mg/kg) or vehicle. Colonic tissues were harvested 48 h post-TNBS-treatment for gene expression and histological analysis. Histological examination on colon tissues collected from mice without TNBS-treatment did not show any differences (figure 5A). As expected, TNBS administration induced acute colonic inflammation in mice administered with as-miR-control, while administration of as-miR-133 α prior to TNBS treatment attenuated colitis development (figure 5A). Additionally, quantitative PCR analysis showed that intracolonic administration of as-miR-133 α significantly reduced endogenous colonic expression of miR-133 α ($p<0.05$, figure 5B). Mice treated with as-miR-133 α and TNBS showed a significant reduction in colon weight after normalisation against colon length (figure 5C, $p<0.05$), neutrophil infiltration, and an improvement in mucosal integrity which resulted in a significantly improved histological score (figure 5D, $p<0.05$). Proinflammatory cytokine production was generally increased in colon tissues treated with TNBS. Pretreatment with as-miR-133 α prior to TNBS administration

Figure 3 Overexpression of miR-133 α promoted proinflammatory signalling in NCM460-neurotensin receptor 1 (NTR1) cells. (A) Levels of MAP kinase phosphorylation, including ERK1/2 and c-Jun and (B) NF- κ B p65 phosphorylation in NCM460-NTR1 cells transfected with miR-133 α and its control was analysed as stated in Methods. *p<0.05, compared to cells transfected with control miRNA precursor. (C) Translocation of NF- κ B p65 in NCM460-NTR1 cells transfected with miR-133 α and its control was analysed by ELISA. *p<0.05, compared to cells transfected with control miRNA precursor. (D) Expression of IL-8, IL-1 β and TNF- α in NCM460-NTR1 cells transfected with miR-133 α and its control was analysed by qPCR analysis. *p<0.05, compared to cells transfected with control miRNA precursor. (E) IL-8 production in NCM460-NTR1 cells transfected with miR-133 α and its control was measured using IL-8 ELISA. *p<0.05, compared to cells transfected with control miRNA precursor. (F) IL-1 β and IL-1 α production in NCM460-NTR1 cells transfected with miR-133 α and its control was measured as stated in Methods. *p<0.05, compared to cells treated with vehicle control; #p<0.05, compared to cells transfected with miR control precursor.



blocked increased expression of the neutrophil product lipocalin 2 (lcn2), and TNF- α , a major proinflammatory cytokine in colitis. The production of cxcl1, an important neutrophil chemoattractant (figure 5E, p<0.05) was significantly reduced in mice administered with as-miR-133 α , when compared to control as-miR-administered counterparts (figure 5E). We also investigated the effect of as-miR-133 α treatment in the DSS-induced colitis model. Administration of as-miR-133 α and its control was performed at 24 h and 72 h before DSS treatment. Mice administered with as-miR-133 α and DSS showed reduced weight loss 5 days after DSS treatment (see online supplementary figure S1A, p<0.05) while levels of TNF- α and IL-1 β were increased 5 days after DSS-induced colitis and significantly reduced in mice administered with as-miR-133 α (see online supplementary figure S1B, C, p<0.05). In conclusion, these results suggested that miR-133 α is involved in the pathophysiology of colitis.

MiR-133 α directly regulates AFTPH expression through binding its 3' UTR in colonic epithelial cells

MiRs act as gene-silencers by inhibiting the expression of mRNA transcripts through binding to the 3' UTRs of target genes.¹⁴ To identify candidate genes involved in the NTR1-miR-133 α response, we searched for genes with possible miR-133 α binding sites in their 3' UTRs. In silico search using three online databases (TargetScanHuman (<http://www.targetscan.org>); miRBase (<http://www.mirbase.org>) and PicTar (<http://pictar.mdc-berlin.de>)), identified aftiphilin (AFTPH) with a miR-133 α binding site at its 3' UTR highly conserved across species (figure 6A).

The NTR1-miR-133 α -AFTPH interaction was first validated in NCM460-NTR1 cells transfected with as-miR-133 α and exposed to NT. NT exposure downregulated AFTPH mRNA levels, as shown in quantitative PCR analysis (figure 6B, p<0.05) and AFTPH 3'UTR-regulated luciferase activity assay (figure 6C, p<0.05) when compared with control cells. To confirm the direct interaction between miR-133 α and AFTPH, we deleted the miR-133 α binding sequence on the AFTPH 3' UTR and showed that NT-induced downregulation of AFTPH 3' UTR luciferase activity was not observed in the absence of the miR-133 α binding site (figure 6D). Overexpression of miR-133 α in HEK293 cells significantly reduced AFTPH-3' UTR-associated luciferase activity in the absence of NT stimulation (figure 6E, p<0.05). Moreover, treatment with the NTR1-specific antagonist of SR48962 prevented the reduction in AFTPH protein expression induced by NT (figure 6F). The above results strongly indicate that AFTPH is a downstream target of miR-133 α in colonocytes after NT exposure, suggesting that NT/NTR1 signalling suppresses AFTPH expression through miR-133 α overexpression.

To confirm our in vitro findings in the in vivo setting, we compared expression of AFTPH mRNA in colon tissues from control, NT-exposed and TNBS-exposed (2d) mouse colon. Our results show significantly decreased colonic AFTPH mRNA expression in NT-administered (figure 6G, p<0.05) and TNBS-injected mice (figure 6H, p<0.05) compared to vehicle-exposed control colon, consistent with the increased miR-133 α levels observed in both models (figures 4A and 5F). In NTR1KO mice exposed to DSS, only a small (by 1.28 ± 0.17-fold, n=5 per group) but not significant increase in

Colon

Figure 4 Levels of miR-133 α were upregulated during experimental colitis in vivo. (A) miR-133 α levels in colon tissues from C57BL6/J mice intracolonically administered with NT (300 μ g/kg) was analysed by qPCR analysis. * $p<0.05$, compared to mice receiving vehicle control. (B) Expression of miR-133 α in colon tissues from C57BL6/J mice collected 2 and 7 days after 5 mg/kg trinitrobenzene sulfonic acid (TNBS) administration and 5 days after 5% dextran sodium sulfate (DSS) feeding was analysed by qPCR analysis. * $p<0.05$, compared to treatment control. (C) Representative images of in situ hybridisation of miR-133 α of colon tissues from TNBS-treated C57BL6/J mice and their control counterparts (arrows, epithelial cells; arrow heads, lamina propria cells infiltration). Scale: 50 μ m. (D) Expression of miR-133 α in colon tissues from neurotensin receptor 1 (NTR1) KO and its control counterparts collected 5 days after 5% DSS feeding was analysed by qPCR analysis. * $p<0.05$, compared to wild-type mice with same treatment.


AFTPH levels were observed. Along these lines, silencing of miR-133 α by intracolonic administration of as-miR-133 α increased AFTPH levels in normal and inflamed colon tissues, when compared to their control counterparts (figure 6H, $p<0.05$).

AFTPH gene silencing promotes colonic proinflammatory responses in human colonic epithelial cells

To examine the functionality of AFTPH in NT-mediated proinflammatory signalling, we silenced expression of AFTPH in NCM460-NTR1 cells with si-RNA transfection prior to NT stimulation. Our results show that AFTPH gene silencing enhanced NT-induced c-Jun and NF- κ B activation (figure 7A, B, $p<0.05$), consistent with our findings for a role of miR-133 α in NT-mediated signalling activation. Additionally, gene silencing of AFTPH increased basal IL-1 β production (figure 7C, $p<0.05$), while ameliorated NT-mediated IL-1ra reduction in human colonic epithelial cells (figure 7C), in line with our results from miR-133 α overexpression (figure 3F).

Levels of miR-133 α and AFTPH are dysregulated in colon of patients with UC

To confirm our mouse findings, we compared levels of miR-133 α and AFTPH in colonic tissue samples from UC and normal colon tissues. Levels of miR-133 α were significantly increased by 2.6 ± 1.82 -fold ($p=0.048$, figure 8A) in patients with UC with active disease, while AFTPH mRNA levels were decreased by 0.72 ± 0.17 -fold ($p=0.0073$, figure 8A) in the same patients. By contrast, levels of miR-133 α obtained from patients with CD did not show significant difference compared with normal controls ($n=8$ per group, data not shown). Additionally, immunohistochemistry of colon tissues from patients with UC showed reduced AFTPH expression in the




colonic mucosa when compared with normal colon tissues (figure 8B), with AFTPH expressed primarily in colonic epithelial cells, and colocalised with villin, an epithelial cell marker (figure 8B). The above results suggested that as in mouse colitis, miR-133 α and AFTPH expression are inversely dysregulated in patients with UC.

DISCUSSION

Coupling of NT to its high-affinity receptor NTR1 triggers MAPK^{38 39} and NF- κ B^{7 40 41} signalling and promotes intestinal inflammation.^{2 5 13} We have previously shown that NT alters expression of several microRNAs, including miR-133 α in human colonic epithelial cells, and presented evidence that microRNAs represent an important functional component of the effects of NT in the intestine.¹⁰ In this current study, we examined the role of miR-133 α in the NT-associated intestinal inflammatory circuit in vitro and in vivo. Our results demonstrate that expression of miR-133 α is increased following NT stimulation of colonocytes in vitro and normal mouse colon in vivo. We also present strong evidence that miR-133 α is involved in the inflammatory cascade activated by NT/NTR1 interactions in human colonocytes, and that silencing of endogenous miR-133 α in the colon diminishes acute experimental colitis in two different mouse models. Last, our results point to AFTPH as a novel downstream target of miR-133 α that mediates intestinal proinflammatory signalling and cytokine transcription following NT-miR-133 α interactions. Thus, the NT-regulated miR-133 α and its target AFTPH may represent new targets for colonic inflammatory responses in vivo.

Our present study focused on understanding the role of miR-133 α in NT-related colonic inflammation. NT/NTR1 signalling promotes acute colitis,^{2 5} while NTR1 deficiency

Figure 5 Intracolonic administration of as-miR-133 α attenuated trinitrobenzene sulfonic acid (TNBS)-induced colitis development in wild-type mice. (A) Representative images from H&E staining of colon tissues from C57BL6/J mice intracolonically administered with as-miR-133 α and its control with or without TNBS treatment. Scale: 50 μ m. (B) Levels of miR-133 α in colon tissues from C57BL6/J mice intracolonically administered with as-miR-133 α and its control were analysed using qPCR analysis. * $p<0.05$, compared to mice administered with control as-miRNA. (C) Colon weight from C57BL6/J mice intracolonically administered with as-miR-133 α and its control with or without TNBS treatment were measured and normalised by colon length. * $p<0.05$, compared to mice administered with control as-miRNA. (D) Total histological score, neutrophil infiltration and mucosal integrity of colon tissues from C57BL6/J mice intracolonically administered with as-miR-133 α and its control with TNBS treatment were scored. * $p<0.05$, ** $p<0.01$, *** $p<0.005$ compared to mice administered with control as-miRNA. (E) Levels of lcn2, TNF- α and cxcl1 in colon tissues from C57BL6/J mice intracolonically administered with as-miR-133 α and its control with or without TNBS treatment were analysed using quantitative PCR analysis. * $p<0.05$, compared to mice administered with control as-miRNA.



reduces the severity of experimental colitis.^{5–13} Additionally, NT and NTR1 are overexpressed in colon tissues from patients with UC,⁶ and as shown in our current study, NTR1KO mice have reduced miR-133 α colonic levels compared to wild-type mice. On the other hand, miR-133 α and its downstream target expression have been associated with different disease states, such as colorectal cancer,^{21–28} myocyte development, and hypertrophy,^{42–44} and fibrosis in the cardiac muscle and liver.^{45–46} However, its role in colonic inflammation has never been studied. Several studies have shown that miR-133 α modulates ERK^{29–30} and PI3 K/Akt^{21–31} signalling pathways, or directly targets molecules involved in these pathways.²² Consistent with the above findings, we have shown that miR-133 α modulates ERK1/2 and p38 activation upon NT stimulation. We also demonstrate for the first time that miR-133 α directly regulates NF- κ B activation and NT exposure, together with miR-133 α overexpression, in cells increased the production of proinflammatory IL-1 β and suppressed that of the anti-inflammatory IL-1ra. IL-1ra antagonises IL-1 β in colitis in vivo.^{47–48} Interestingly, a reduced IL-1ra/IL-1 β ratio is considered as an important factor for the severity of inflammation in patients with IBD.⁴⁹ Taken together, our results suggest that NT/NTR1 coupling triggers

proinflammatory signalling pathways by modulating miR-133 α levels in human colonic epithelial cells.

Our results showed that miR-133 α levels were significantly increased in the colonic mucosa of mice with acute TNBS-induced and DSS-induced colitis, as well as colon tissues of patients with UC with active disease, compared to controls. MiR-133 α was primarily expressed in colonic epithelial cells as shown by *in situ* hybridisation in mouse colon, with increased expression of this miR during TNBS-induced colitis. However, miR-133 α was not detectable in sera from patients with UC and their normal counterparts (data not shown). To study the role of endogenous miR-133 α in colitis, we introduced as-miR-133 α intracolonically in the two experimental colitis models to avoid perturbation of normal cellular functions of non-target tissues. Our results showed that mice receiving intracolonial as-miR-133 α had reduced endogenous miR-133 α levels, improved mucosa histology and reduced proinflammatory cytokine expression in colon tissues of mice with colitis. Intracolonial delivery of antisense oligonucleotides is largely confined to colonic epithelial cells in vivo,⁵⁰ our results suggest that miR-133 α levels in colonic epithelial cells can influence the pathophysiology of colitis. However, since expression of miR-133 α has also been identified in circulating monocytes,⁵¹

Colon




Figure 6 Aftiphilin (AFTPH) was the downstream target of neurotensin (NT)-induced miR-133 α upregulation. (A) Conservative miR-133 α binding sites were identified in 3' untranslated region (UTR) region of aftiphilin across different species. (B) Levels of miR-133 α in NCM460-neurotensin receptor 1 (NTR1) cells transfected with as-miR-133 α and its control was analysed by qPCR analysis. *p<0.05, compared to cells stimulated by vehicle. (C) AFTPH 3'UTR-driven luciferase activities from NCM460-NTR1 cells transfected with as-miR-133 α and its control were measured after NT stimulation. *p<0.05, compared to cells stimulated by vehicle. (D) Luciferase activity driven by AFTPH 3'UTR with or without miR-133 α binding sequence was measured after NT stimulation. *p<0.05, compared to cells stimulated by vehicle. (E) AFTPH 3'UTR-driven luciferase activities from HEK293 cells transfected with miR-133 α precursor (miR-133 α) and its control were measured. *p<0.05, compared to cells transfected with control miRNA precursor. (F) Immunoblot analysis of AFTPH expression in NCM460-NTR1 cells 30 min and 6 h after NT exposure in the presence or absence of SR48962 (10 nM). (G) AFTPH levels in colon tissues from C57BL/6J mice intracolonically administered with NT and its control and (H) from C57BL/6J mice intracolonically administered with as-miR-133 α and its control with or without trinitrobenzene sulfonic acid treatment were analysed using qPCR analysis. *p<0.05, compared to mice administered with control as-miRNA.

and in subepithelial cells of the colonic mucosa, participation of miR-133 α in inflammatory signalling in colonic lamina propria cell types cannot be excluded.

We identified a novel miR-133 α target, AFTPH, in which its expression level is regulated by NT/NTR1 interactions. Several




Figure 7 Gene silencing of aftiphilin (AFTPH)-promoted neurotensin (NT)-associated proinflammatory response in vitro. Levels of (A) c-Jun and (B) NF-κB phosphorylation in NCM460-neurotensin receptor 1 (NTR1) cells transfected with si-AFTPH and its control was measured as stated in Methods. *p<0.05, compared to cells transfected with control siRNA; #p<0.05, compared to NT-treated cells. (C) IL-1 β and IL-1 α production of NCM460-NTR1 cells transfected with si-AFTPH and its control in the presence or absence of NT stimulation was analysed as stated in Methods. *p<0.05, compared to cells transfected with control siRNA.

pieces of evidence support our identification of AFTPH as a miR-133 α downstream target. Incubation of human colonic epithelial cells with NT decreases AFTPH mRNA levels, as well as AFTPH 3'UTR-regulated luciferase activity. Most importantly, expression of AFTPH with a deleted miR-133 α binding sequence on its 3' UTR is no longer downregulated in response to NT exposure. The presence of AFTPH has not been previously recognised in the intestine, and its cellular function is not well characterised. AFTPH is essential to intracellular trafficking with binding sites for clathrin and adaptor proteins (AP-1, AP-2).⁵² AFTPH is localised in trans-golgi network (TGN), and its gene silencing leads to unregulated exocytosis of Weibel-Palade bodies in endothelial cells⁵⁴ without altering TGN morphology.⁵⁵ However, the role of AFTPH or TGN in proinflammatory signalling cascades has not been investigated. Here, we demonstrate that AFTPH levels are significantly downregulated in colon tissues from patients with UC, as well as in the colon of mice with TNBS-induced and DSS-induced colitis. At the cellular level, AFTPH gene silencing activates the NF-κB pathway and promotes proinflammatory cytokine IL-1 β production in vitro. NF-κB activation is involved in IL-1 β expression⁵⁶ and administration of NF-κB inhibitor reduce IL-1 β production and ameliorates colonic inflammation in mice.⁵⁷ Thus, dysregulated AFTPH expression may modulate proinflammatory responses, representing a novel regulator in proinflammatory signalling pathways and colitis development. Moreover, measurements of its expression levels, together with measurements of the levels of miR-133 α , may serve as biomarkers in UC.




Figure 8 Levels of miR-133 α and aftiphilin (AFTP) were significantly dysregulated in patients with ulcerative colitis (UC). (A) Levels of miR-133 α and AFTP in colon tissues from normal and patients with UC were analysed using qPCR analysis. (B) Representative images from immunohistochemistry against AFTP and villin of colon tissues from normal and patients with UC. Scale: 50 μ m.

Overall, our results represent a previously unrecognised paradigm whereby activation of a neuropeptide receptor (NTR1) in the colonic mucosa stimulates the expression of a miR (miR-133 α) that modulates the development of colitis. The therapeutic potential of miR-133 α silencing and stabilisation of colonic AFTP levels against colitis deserves further investigation.

Contributors Overall project design and hypotheses were developed by IKML and CP. IKML conducted experiments, analysed the data and wrote the manuscript. KB generated the human colon epithelial cells overexpressing NTR1, NCM460-NTR1. DH and AO collected and provided sera from UC and control patients and for analysing these new results now included in the revised manuscript. CP, DI and CP contributed to experimental design and provided animals and reagents. All authors participated in revising the manuscript and agreed to the final version. CP and DI supervised the overall project.

Funding This work was supported by NIH grant DK60729 (CP), the Neuroendocrine Assay Core, NIDDK P50 DK 64539 (CP), the Blinder Research Foundation for Crohn's Disease (IKML), and the Crohn's and Colitis Foundation of America (KB).

Competing interests None.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

- Polak JM, Sullivan SN, Bloom SR, et al. Specific localisation of neuropeptid Y to the N cell in human intestine by radioimmunoassay and immunocytochemistry. *Nature* 1977;270:183–4.
- Castagliuolo I, Wang CC, Valenick L, et al. Neuropeptid Y is a proinflammatory neuropeptide in colonic inflammation. *J Clin Invest* 1999;103:843–9.
- Tanaka K, Masu M, Nakanishi S. Structure and functional expression of the cloned rat neuropeptid Y receptor. *Neuron* 1990;4:847–54.
- Fassio A, Evans G, Grissommer R, et al. Distribution of the neuropeptid Y receptor NTS1 in the rat CNS studied using an amino-terminal directed antibody. *Neuropharmacology* 2000;39:1430–42.
- Koon HW, Kim YS, Xu H, et al. Neuropeptid Y induces IL-6 secretion in mouse preadipocytes and adipose tissues during 2,4,6-trinitrobenzenesulphonate acid-induced colitis. *Proc Natl Acad Sci USA* 2009;106:8766–71.
- Brun P, Mastrotto C, Beggio E, et al. Neuropeptid Y stimulates intestinal wound healing following chronic intestinal inflammation. *Am J Physiol Gastrointest Liver Physiol* 2005;288:G621–9.
- Zhao D, Keates AC, Kuhnt-Moore S, et al. Signal transduction pathways mediating neuropeptid Y-stimulated interleukin-8 expression in human colonocytes. *J Biol Chem* 2001;276:44464–71.
- Sundman L, Saarialho-Kere U, Vendelin J, et al. Neuropeptid Y receptor 1 expression in the intestine and skin—putative role in peptide hormone secretion. *Neurogastroenterol Motil* 2010;22:79–87, e30.
- da Silva L, Neves BM, Moura L, et al. Neuropeptid Y downregulates the pro-inflammatory properties of skin dendritic cells and increases epidermal growth factor expression. *Biochim Biophys Acta* 2011;1813:1863–71.
- Bakirtzi K, Hatzipostolou M, Karagiannides I, et al. Neuropeptid Y signalling activates microRNAs-21 and -155 and Akt, promotes tumor growth in mice, and is increased in human colon tumors. *Gastroenterology* 2011;141:1749–61.e1.
- Margolis KG, Gershon MD. Neuropeptides and inflammatory bowel disease. *Curr Opin Gastroenterol* 2009;25:503–11.
- Zhao D, Pothoulakis C. Effects of NT on gastrointestinal motility and secretion, and role in intestinal inflammation. *Peptides* 2006;27:2434–44.
- Bugni JM, Rabadi LA, Jubbal K, et al. The neuropeptid Y receptor-1 promotes tumor development in a sporadic but not an inflammation-associated mouse model of colon cancer. *Int J Cancer* 2012;130:1798–805.
- McKenna LB, Schug J, Vourekas A, et al. MicroRNAs control intestinal epithelial differentiation, architecture, and barrier function. *Gastroenterology* 2010;139:1654–64, 1664 e1.
- Bartel DP. MicroRNAs: target recognition and regulatory functions. *Cell* 2009;136:215–33.
- Wu F, Zikusoka M, Trindade A, et al. MicroRNAs are differentially expressed in ulcerative colitis and alter expression of macrophage inflammatory protein-2 alpha. *Gastroenterology* 2008;135:1624–35.e24.
- Bian Z, Li L, Cui J, et al. Role of miR-150-targeting c-Myb in colonic epithelial disruption during dextran sulphate sodium-induced murine experimental colitis and human ulcerative colitis. *J Pathol* 2011;225:544–53.
- Fasseu M, Treton X, Guichard C, et al. Identification of restricted subsets of mature microRNA abnormally expressed in inactive colonic mucosa of patients with inflammatory bowel disease. *PLoS ONE* 2010;5:e13160.
- Takagi T, Naito Y, Mizushima K, et al. Increased expression of microRNA in the inflamed colonic mucosa of patients with active ulcerative colitis. *J Gastroenterol and Hepatol* 2010;25:S129–33.
- Wu F, Zhang S, Dassopoulos T, et al. Identification of microRNAs associated with ileal and colonic Crohn's disease. *Inflamm Bowel Dis* 2010;16:1729–38.
- Josse C, Bouznad N, Geurts P, et al. Identification of a microRNA landscape targeting the PI3K/Akt signalling pathway in inflammation-induced colorectal carcinogenesis. *Am J Physiol Gastrointest Liver Physiol* 2014;306:G229–43.
- Wang H, An H, Wang B, et al. miR-133a represses tumour growth and metastasis in colorectal cancer by targeting LIM and SH3 protein 1 and inhibiting the MAPK pathway. *Eur J Cancer* 2013;49:3924–35.
- Arndt GM, Dossey L, Cullen LM, et al. Characterization of global microRNA expression reveals oncogenic potential of miR-145 in metastatic colorectal cancer. *BMC Cancer* 2009;9:374.
- Varner AL, French AJ, Borralho PM, et al. Human colon cancer profiles show differential microRNA expression depending on mismatch repair status and are characteristic of undifferentiated proliferative states. *BMC Cancer* 2009;9:401.
- Dong Y, Zhao J, Wu CW, et al. Tumor suppressor functions of miR-133a in colorectal cancer. *Mol Cancer Res* 2013;11:1051–60.
- Ma Y, Zhang P, Yang J, et al. Candidate microRNA biomarkers in human colorectal cancer: systematic review profiling studies and experimental validation. *Int J Cancer* 2012;130:2077–87.
- Necela BM, Carr JM, Asmann YW, et al. Differential Expression of MicroRNAs in Tumors from Chronically Inflamed or Genetic (APC^{Min/+}) Models of Colon Cancer. *PLoS ONE* 2011;6:e18501.
- Li Z, Gu X, Fang Y, et al. microRNA expression profiles in human colorectal cancers with brain metastases. *Oncol Lett* 2012;3:346–50.
- Li Q, Lin X, Yang X, et al. NFATc4 is negatively regulated in miR-133a-mediated cardiomyocyte hypertrophic repression. *Am J Physiol Heart Circ Physiol* 2010;298:H1340–7.
- Feng Y, Niu LL, Wei W, et al. A feedback circuit between miR-133 and the ERK1/2 pathway involving an exquisite mechanism for regulating myoblast proliferation and differentiation. *Cell Death Dis* 2013;4:e934.
- Cui W, Zhang S, Shan C, et al. microRNA-133a regulates the cell cycle and proliferation of breast cancer cells by targeting epidermal growth factor receptor through the EGFR/Akt signalling pathway. *FEBS J* 2013;280:3962–74.
- Zhao D, Bakirtzi K, Zhan Y, et al. Insulin-like growth factor-1 receptor transactivation modulates the inflammatory and proliferative responses of neuropeptid Y in human colonic epithelial cells. *J Biol Chem* 2011;286:6092–9.
- Koon HW, Shih D, Karagiannides I, et al. Substance P modulates colitis-associated fibrosis. *Am J Pathol* 2010;177:2300–9.

- 34 Koon HW, Zhao D, Xu H, et al. Substance P-mediated expression of the pro-angiogenic factor CCN1 modulates the course of colitis. *Am J Pathol* 2008;173:400–10.
- 35 Evers BM, Izukura M, Townsend CM Jr, et al. Neurotensin prevents intestinal mucosal hypoplasia in rats fed an elemental diet. *Dig Dis Sci* 1992;37:426–31.
- 36 Maoret JJ, Pospai D, Rouyerfessard C, et al. Neurotensin receptor and its mRNA are expressed in many human colon cancer cell lines but not in normal colonic epithelium: binding studies and RT-PCR experiments. *Biochem Biophys Res Commun* 1994;203:465–71.
- 37 Gui X, Guzman G, Dobner PR, et al. Increased neurotensin receptor-1 expression during progression of colonic adenocarcinoma. *Peptides* 2008;29:1609–15.
- 38 Zhao D, Zhan Y, Koon HW, et al. Metalloproteinase-dependent transforming growth factor-alpha release mediates neurotensin-stimulated MAP kinase activation in human colonic epithelial cells. *J Biol Chem* 2004;279:43547–54.
- 39 Zhao D, Zhan Y, Zeng H, et al. Neurotensin stimulates expression of early growth response gene-1 and EGFR receptor through MAP kinase activation in human colonic epithelial cells. *Int J Cancer* 2007;120:1652–6.
- 40 Zhao D, Kuhnt-Moore S, Zeng H, et al. Neurotensin stimulates IL-8 expression in human colonic epithelial cells through Rho GTPase-mediated NF-kappa B pathways. *Am J Physiol Cell Physiol* 2003;284:C1397–404.
- 41 Zhao D, Zhan Y, Zeng H, et al. Neurotensin stimulates interleukin-8 expression through modulation of I kappa B alpha phosphorylation and p65 transcriptional activity: involvement of protein kinase C alpha. *Mol Pharmacol* 2005;67:2025–31.
- 42 Rao PK, Kumar RM, Farkhondeh M, et al. Myogenic factors that regulate expression of muscle-specific microRNAs. *Proc Natl Acad Sci USA* 2006;103:8721–6.
- 43 Chen X, Wang K, Chen J, et al. In vitro evidence suggests that miR-133a-mediated regulation of uncoupling protein 2 (UCP2) is an indispensable step in myogenic differentiation. *J Biol Chem* 2009;284:5362–9.
- 44 Hua Y, Zhang Y, Ren J. IGF-1 deficiency resists cardiac hypertrophy and myocardial contractile dysfunction: role of microRNA-1 and microRNA-133a. *J Cell Mol Med* 2012;16:83–95.
- 45 Roderburg C, Luedde M, Vargas Cardenas D, et al. miR-133a mediates TGF- β -dependent derepression of collagen synthesis in hepatic stellate cells during liver fibrosis. *J Hepatol* 2013;58:736–42.
- 46 Castoldi G, di Gioia CRT, Bombardi C, et al. MiR-133a regulates collagen 1A1: potential role of miR-133a in myocardial fibrosis in angiotensin II-dependent hypertension. *J Cell Physiol* 2012;227:850–6.
- 47 Cominelli F, Nast CC, Clark BD, et al. Interleukin 1 (IL-1) gene expression, synthesis, and effect of specific IL-1 receptor blockade in rabbit immune complex colitis. *J Clin Invest* 1990;86:972–80.
- 48 Ferretti M, Casini-Raggi V, Pizarro TT, et al. Neutralization of endogenous IL-1 receptor antagonist exacerbates and prolongs inflammation in rabbit immune colitis. *J Clin Invest* 1994;94:449–53.
- 49 Casini-Raggi V, Kam L, Chong YJ, et al. Mucosal imbalance of IL-1 and IL-1 receptor antagonist in inflammatory bowel disease. A novel mechanism of chronic intestinal inflammation. *J Immunol* 1995;154:2434–40.
- 50 Huang Z, Shi T, Zhou Q, et al. miR-141 Regulates colonic leukocytic trafficking by targeting CXCL12 β during murine colitis and human Crohn's disease. *Gut* 2014;63:1247–57.
- 51 Wang Y, Li L, Moore BT, et al. MiR-133a in human circulating monocytes: a potential biomarker associated with postmenopausal osteoporosis. *PLoS ONE* 2012;7:e34641.
- 52 Mattera R, Ritter B, Sidhu SS, et al. Definition of the consensus motif recognized by gamma-adaptin ear domains. *J Biol Chem* 2004;279:8018–28.
- 53 Burman JL, Wasiak S, Ritter B, et al. Aftiphilin is a component of the clathrin machinery in neurons. *FEBS Lett* 2005;579:2177–84.
- 54 Lui-Roberts WW, Ferraro F, Nightingale TD, et al. Aftiphilin and gamma-synergins are required for secretagogue sensitivity of Weibel-Palade bodies in endothelial cells. *Mol Biol Cell* 2008;19:5072–81.
- 55 Hirst J, Borner GH, Harbour M, et al. The aftiphilin/p200/gamma-synergins complex. *Mol Biol Cell* 2005;16:2554–65.
- 56 Carvalho FA, Aitken JD, Gewirtz AT, et al. TLR5 activation induces secretory interleukin-1 receptor antagonist (sIL-1Ra) and reduces inflammasome-associated tissue damage. *Mucosal Immunol* 2011;4:102–11.
- 57 Funakoshi T, Yamashita K, Ichikawa N, et al. A novel NF- κ B inhibitor, dehydroxymethylepoxyquinomicin, ameliorates inflammatory colonic injury in mice. *J Crohns Colitis* 2012;6:215–25.