DUOGASTRONE® direct healing of duodenal ulcer

Trials suggest*

- radiological disappearance of ulcer crater
- relief of symptoms within a few days
- superiority over antacid or anticholinergic therapy

- patients can lead a normal life
- special diets are unnecessary
- even chronic cases, with a long history of symptoms, respond

Unique 'positioned release' capsules deliver the active ingredient (50 mg carbenoxolone sodium) into the duodenum.


* Further trials are in progress to study the effects of DUOGASTRONE in long-term therapy



Full information available on request from

Berk Pharmaceuticals Limited Godalming Surrey England
Made under licence from Biorex Laboratories Ltd., London. Brit. Pat. Nos. 843,133 & 1,093, 286

R Regd. Trade Mark



GRYSTALLINE AMINO-ACIDS



VAMIN provides, for the first time, a physiological mixture of pure crystalline amino acids, all in the L-form, for intravenous use.

For the few occasions on which a crystalline solution may be preferred to the casein hydrolysate (Aminosol), VAMIN offers a balanced combination of essential and non-essential amino acids together with fructose and minerals to meet basic nutritional requirements.

1 litre VAMIN provides 9.4 G amino nitrogen and 650 calories. Additional calories are available from Intralipid.


Packing: 500 ml. Descriptive literature on request.

ALL IN THE LFORM



PAINES & BYRNE LIMITED, PABYRN LABORATORIES, GREENFORD, MIDDX.

This little tablet takes such a lot of trouble



out of amylase testing

First of the Pharmacia diagnostic reagents for the determination of biologically active substances is the Phadebas amylase *tablet* test.


Phadebas® Amylase Test

- No tedious preparation of reagents.
- Standardization—from tablet to tablet and bottle to bottle, reproducibility is certain. (Standard curve supplied).
- Accuracy ensured by entirely new type of starch substrate.
- Stability-no refrigeration required.
- Simplicity—less apparatus required.
- Rapidity of result.
- · Highly sensitive.
- · Highly precise.
- Wide range.
- Suitable for all body fluids.

Pharmacia (Great Britain) Limited Paramount House, 75 Uxbridge Road, London, W.5. Tel: 01-579 0102/7




An inadequate barrier is no protection



Prolonged clinical evaluation has shown NOXYFLEX to be singularly effective against the intrusion of bacteria.

NOXYFLEX literature outlines its proven uses in detail. Have you a copy?

Protection: NOXYFLEX 2.5G. Noxytiolin and 10 mgm. Amethocaine HC1 Geistlich Chester



The pre-and post-operative patient may require calories in the form of carbohydrates. These are best provided with minimal fluid volume. Just one litre of Sorbitol 30% 'Egic' provides your patient with 1200 calories, whereas three litres of the conventional dextrose 10% solution would be needed. Also with Sorbitol 30%—the risk of venous thrombosis is dramatically reduced and there is no danger of hyperglycaemia, or reactional hypoglycaemia, it has superior antiketogenic properties over fructose and is not insulin dependent. Sorbitol 30% 'Egic'—today's intravenous carbohydrate provider.

"In clinical practice Sorbitol has proved to be almost fully retained in the body and can replace intravenous dextrose as a carbohydrate source. Although like dextrose it yields only 4 cal. per gramme of carbohydrate, it can be safely administered as a 30 per cent solution containing 1200 cal. per litre. In a grossly infected surgical patient requiring intravenous feeding for 3 months, Sorbitol was used as the main source of calories without alteration in the patient's nitrogen balance."

1. Reference: "The Utilisation and Metabolism of Intravenous Sorbitol" Brit. J. Surgery (1969), 56, 653.

Sorbitol 30% 'Egic' Bottles of 500ml. Complete with giving set.

Trophysan. Wherever oral protein intake is insufficient Trophysan provides more readily assimilable free-amino nitrogen as pure crystalline amino acids in balanced proportions plus Sorbitol, Vitamins and Minerals. Trophysan often produces an earlier return to normal oral feeding and speeds recovery.

two better solutions for parenteral nutrition from Servier Laboratories . .



Full information from

Servier Laboratories Limited, formerly Selpharm Laboratories Ltd. Percival House, Pinner Road, Harrow, Middlesex, HA1 4HQ.

products of L'Equilibre Biologique, France



Have a closer look...

Examine the facts. Scrutinise the bibliography. Test the rationale. It's all there in 'A Scientific Assessment of Alka-Seltzer' new presentation of the science behind a familiar home remedy which has antacid and analgesic properties. This is an unusual

book about an unusual product. Worth reading. Send for a copy. Miles Laboratories Limited, Medical Dept.11A Stoke Court. Stoke Poges, Slough, Bucks.



Alka-Seltzer is a trade mark

Maxolon a powerful anti-emetic... which also restores gastric motility

Maxolon is of proven value in nausea and vomiting...

Maxolon acts centrally on the chemoreceptor trigger zone, and also acts locally on the afferent nerves of the pylorus and duodenum leading to the vomiting centre. Maxolon is remarkably effective in the treatment of nausea and vomiting from a wide range of gastric disorders. In a clinical trial (1) Maxolon achieved an 80% reduction score in

achieved an 80% reduction score nausea and 87% in vomiting.

... Maxolon also restores gastric motility

This makes it unique among products used in the alimentary tract. Spasm where present is relieved and stasis gives way

to normal peristalsis. This action is of great importance in the treatment of nausea and vomiting but also leads to a consideration of further therapeutic applications. Among these are its known value in the treatment of post vagotomy symptoms, post operative gastric hypotonia, duodenal intubation and diagnostic radiology.

Maxolon has a good safety record. It does not sedate the patient, nor affect gastric or salivary secretion.

Maxolon is a unique compound—with a well proven clinical record.

Are you finding in practice how much more there is to Maxolon?

(1) Brit. J. clin. Pract. 1967, 21, 457

Further information is available on request

Maxolon

a powerful anti-emetic and so much more









Digestion

Index Vol. 3, No. 2 1970 Smoldas, J. (Olomouc):

Some Problems of the Ultrasonic Diagnosis of Bile Stones

Pirk, F. and Skála, I. (Prague):

Motility of the Digestive Tract After Administration of Medium Chain Triglycerides (MCT) as Compared with Long Chain Triglycerides (LCT)

Pedersen, S. A. (Odense):

Pentagastrin as a Stimulant in Routine Clinical Testing of Gastric Acid Secretion

Elder, J. B. (Glasgow):

Achalasia of the Cardia in Childhood

Beno, I.: Bučko, A.: Chorváthová, V. and Babala, J. (Bratislava):

Absorption of d-xylose and Fat and Histological Changes of the Small Intestine in Patients after Partial Gastrectomy

Mauer, I. and Levitan, R. (Chicago, IL):

Regional Enteritis and Celiac Artery Stenosis with Negative Small Bowel Studies. A Case Report

Bianchi Porro, G. and Maiolo, A. T. (Milano):

Cerebral Hemodynamics and Metabolism in Hepatic Cirrhosis. Effects of Portacaval Shunt Operation

EDITORIAL

Berndt, H. (Berlin):

Malabsorption in Skin Diseases

BOOK REVIEWS

Index Vol. 3, No. 3 1970 Colombi, A. (Basel):

Early Diagnosis of Fatal Hepatitis

Steinitz, H.; Talis, B. and Stein, B. (Tel-Aviv):

Entamoeba histolytica and Dientamoeba fragilis and the Syndrome of Chronic Recurrent Intestinal Amoebiasis in Israel

Takada, A.; Kobayashi, K. and Takeuchi, J. (Kazanawa):

Gastroenteric Clearance of Albumin in Liver Cirrhosis; Relative Protein-losing Gastroenteropathy

Hoedemaeker, Ph.J. (Groningen):

Heterotopic Gastric Mucosa in the Duodenum

Penzes, L. (Budapest):

Data of the Chemical Composition of the Ageing Intestine

CASE REPORT

Chowers, I.; Eyal, Z. and Saltz, N. J. (Jerusalem):

Cushing's Syndrome Associated with Polyps of the Colon

EDITORIAL

BOOK REVIEWS

Berndt, H. (Berlin-Buch):


Is Appendectomy Followed by Increased Cancer Risk?

Ask for specimen copies

. .----

All orders from the U.K. to be placed with J. Wiley and Sons Ltd., Baffins Lane, Chichester, Sussex

S. Karger Basel (Switzerland) New York



Pharmacia (Great Britain) Ltd, Paramount House, 75 Uxbridge Road, London W.5. Telephone: 01-579 0102/7.



Pharmacia

ARCHIVES FRANÇAISES DES MALADIES DE L'APPAREIL DIGESTIF



The most important French periodical on the subject

Organe Officiel de la Société Nationale Française de Gastro-Entérologie et de ses filiales

Publishing all articles concerning the work of physicians, surgeons, electro-radiologists and biologists which are relevant to Gastro-enterology.

Starting in 1968, the "Archives Françaises des maladies de l'appareil digestif" — publish two numbers a year entitled "Biologie et Gastroentérologie" devoted to reports on experimental work in gastro-enterology carried out in the new research units of the various French University Hospital Centres.

These articles are original works, clinical, radiological, biological and anatomy-pathological papers (with illustrations and bibliography).

The articles are written in French and followed by a summary in French and English.

The Archives also publishes summaries of the Proceedings of the Societies of which they are the

official organ, the texts of conferences or communications which are presented to them, Proceedings of Congresses or Colloquia, book and journal reviews.

The Archives publishes announcements of national and international meetings on all subjects concerned with Gastro-enterology.

SOME ARTICLES RECENTLY PUBLISHED

Thrombo-embolic complications in transcutaneous retrograde arteriography using Seldinger's method by L. P. DOUTRE, G. DELORME, J. TAVERNIER, J. PÉRISSAT and J. PACCALIN (Bordeaux)

The value of guanase assessment in hepatology by Ch. Tamarelle, A. Quinton, J.-J. Dubarry (Bordeaux)

Recurrent rectorrhagias due to rectal varices by J. Faivre, C. Balabaud and C. Béraud (Bordeaux)

A statistical study of the effect antienzymes on acute pancreatites. In connection with 160 surgically verified cases of acute pancreatites by H. SARLES and R. CAMATTE.

Cancers of the gastric stump after resection of the stomach for ulcers. Treatment. Results by M. F. SAEGESSER

Carbonic anhydrase activity from gastric mucosa in the experimental gastric ulcer by D. Dobrescu and A. Cicotti

Subscription (11 numbers) 1971. France F.135. Other countries F.186.

Specimen copy available on application to:

MASSON & CIE-Editeurs. 120 Boulevard Saint Germain, 75 - PARIS 6°

ACTA GASTRO-ENTEROLOGICA BELGICA

SOMMAIRE du FASCICULE 7-1970

Bibliographie.

- J.-P. BARROY, G. WILLEMS, S. VERBEUSTEL et A. GERARD (Bruxelles). Altérations histologiques de la paroi gastrique chez le chien soumis à l'action de la phénylbutazone.
- J.-P. GOVAERTS (Bruxelles). Les répercussions digestives des vagotomies sélectives et totales. (première partie).
- M. De La Pierre, C. Sategna-Guidetti et C. Saracco (Turin). La colite ulcéreuse. Données cliniques ressortant de l'étude de 95 cas.

SOMMAIRE du FASCICULE 8.9-1970

Bibliographie.

- J.-P. GOVAERTS (Bruxelles). Les répercussions digestives des vagotomies sélectives et totales. (deuxième partie).
- G. Ligny, et J. Manouiloff (Charleroi). Débits d'acide et d'électrolytes gastriques chez des individus normaux, des patients dyspeptiques et gastritiques au cours de l'épreuve histaminique 'forte dose'.
- J. CAHEN et Y. BARBIER (Bruxelles). Exérèse locale dans les néoplasmes bas situés de l'ampoule rectale et du canal anal. Indications et résultats.
- A. A. ARIANOFF, PH. VAN REEPINGEN et G. VIELLE (Bruxelles). A propos de deux cas de perforation spontanée du cholédoque chez l'adulte.
 - E. Furnemont (Bruxelles). Lipome sous-muqueux pédiculé du caecum.
- TH. FIRICA, V. GRADINARU et M. RADULESCO (Bucarest). La dilatation kystique congénitale de la voie biliaire principale.
- Th. Gheorghiu (Cologne). Une méthode simple pour l'enlèvement d'un fragment de sonde gastrique retenu accidentellement dans l'estomac.

SOMMAIRE du FASCICULE 10-1970

Bibliographie.

- A. Brassinne (Liège). Importance des hémorragie occultes dans la pathogénie de l'anémie après résection intestinale étendue et dans le syndrome de l'anse aveugle.
 - M. LEVRAT et E. PHELIP (Lyon). Les ictères postopératoires précoces, en apparance primitifs.
- J.-P. GOVAERTS (Bruxelles). Les répercussions digestives des vagotomies sélectives et totales. (Troisième partie et fin).
 - J. STALPORT, Y. STAINIER et P. LETAWE (Huy). Lymphoréticulosarcome du grêle.
- H. DANCOT et W. FEREMANS (Bruxelles). Le traitement des lympho- et réticulosarcomes digestifs primitifs au stade localisé ou régional. Chirurgie ou radiothérapie.
 - J. DE MEERSMAN, J. BOULANGER et M. HORNICK (Arlon). Le syndrome de Peutz-Jeghers.

Acta Gastro-enterologica Belgica. Secrétariat général: rue des Champs Elysées. 43. B 1050. Bruxelles. Belgique.

GASTRIC ULCER PATIENTS STILL FILLING YOUR BEDS?

ambulant patients may result in slightly less rapid ulcer healing than can be achieved by the addition of inpatient care, but the difference is not statistically significant. For practical purposes conventional inpatient care of a gastric ulcer carries no ultimate advantage over ambulant therapy with

that it a gastric older has failed to heal after three months of ambulant therapy was earbenoxolone it is very unlikely to heal carbenoxolone.


subsequently with inpatient care.

The Practitioner, 1969, 202, 402.

BIOGASTRONE®

frees beds and liberates the patient

Full information available on request.




Made under licence from Biorex Laboratories Ltd., London Brit. Pat. No. 843, 133 R Regd. Trade Mark. J. 2995

RIFOCIN-M

in BILIARY TRACT INFECTIONS

"...it is unique in its action in acute cholecystitis and other bacterial diseases of the biliary tract, for which it is the only specific antibiotic."

"A much higher proportion of the organisms found in cholangitis would be expected to be inhibited by this drug than by other antibiotics that achieve effective bile levels ..."²



* Active against penicillinase-producing organisms.

* No cross-resistance or cross-allergenicity with other antibiotics has been demonstrated.

* Well tolerated. No toxic reactions have been reported.

* Dose: 150 mg intramuscularly every 8-12 hours.

RIFOCIN-M (rifamide) is a member of the RIFAMYCINS – a new group of antibiotics discovered and developed in the Research Laboratories of LEPETIT.

References: 1. Med. J. Aust., 1966, 1, 1-7. 2. Brit. J. Pharmacol., 1967, 31, 506-512.

Full information is available on request.



LEPETIT Pharmaceuticals Ltd.

Nicholson House, High Street, Maidenhead, Berks. Tel: Maidenhead 32277

GASTROENTEROLOGY:

The monthly seminar

You may object to our calling a medical journal a seminar. But consider it this way: each month, you share in the results of research and clinical studies of your confreres. When the after-hours free you at the end of the day's appointments, consult GASTROENTEROLOGY. You'll find:

- 1.) Selected summaries. Abstracts of relevant new work, with a brief expert critique.
- 2.) Original papers. Studies of the liver, alimentary tract, pancreas and biliary tract.
- 3.) Case Studies. Dealing with clinical research.

Too, there'll be editorials, symposium reports, and a "special comments" section for readers to present their views of papers that have appeared in recent issues—your chance to participate in the "monthly seminar" when you so desire.

GASTROENTEROLOGY. The official organ of the American Gastroenterological Association. From the Williams and Wilkins Company.

Published monthly/2 volumes per year/Subscription: \$25 (Resident rate: \$15.00) Overseas postage: \$3.00

Sample Selections

- Presentation of the Julius Friedenwald Medal to Andrew Conway Ivy.

 M. I. Grossman
- Effect of Long Term Anticholinergic Therapy on Gastric Acid Secretion, with Observations on the Serial Measurement of Peak Histalog Response. R. P. Norgaard, D. E. Polter, J. W. Wheeler, Jr., and J. S. Fordtran.
- A Controlled Study on the Use of Propantheline and Amylopectin Sulfate (SN-263) for Recurrences in Duodenal Ulcer. D. C. H. Sun and M. L. Ryan.
- Basal and Histamine-stimulated Human Gastric Acid Secretion. Lack of Effect of Indomethacin in Therapeutic Doses. D. H. Winship and G. C. Bernhard.
- Gastric Juice Lactic Acidosis in the Presence of Gastric Carcinoma. D. W. Piper, M. L. Kemp, B. H. Fenton, M. J. Croydon, and A. D. Clarke.

The Williams & Wilkins Co./428 E. Preston St./ Baltimore, Md./21202/U.S.A.

TODAY'S DRUGS

New 1970 Edition

The problem facing the prescribing doctor today is choice. Catalogues of drugs—no matter how detailed—are of little help unless accompanied by advice on which drug to give for what condition. So this new edition of "Today's Drugs" not only gives a full account of the drugs but also includes clear accounts of the management of common conditions such as urinary infections and epilepsy.

The new edition differs from the earlier one in other ways. It deals with only three subjects—infections, vitamin deficiencies and anaemias, and disorders of the nervous system. Further volumes will be published at intervals of about a year, and by the time the whole field of therapeutics has been covered this first volume will have become obsolete and due for replacement. In order to further emphasize the ephemeral nature of books on therapeutics—and to cut costs—the book is a paperback.

Demy Octavo

Paperback

224 pages

PRICE 20s. (U.S.A. \$3.00) including postage

ORDER YOUR COPY NOW FROM

The Publisher, BRITISH MEDICAL JOURNAL, B.M.A. House, Tavistock Square, London WC1H 9JR, or through any leading bookseller