

**Do you want
to stop his pain...
..or heal his ulcer?**

Why not do both at once?

DUOGASTRONE[®]




BIOGASTRONE[®]



Full details available on request

J.3572

A completely new concept in the




treatment of hiatus hernia

'Gaviscon' is an advance on previous therapies because it has physical and chemical properties that are particularly suited to the problems of the conservative management of hiatus hernia.

'Gaviscon' stays longer where it's needed most

In the treatment of hiatus hernia, 'Gaviscon' has a unique physical mode of action. Taken after each meal it forms a demulcent gel which floats on top of the stomach contents and gently occludes the cardiac sphincter for up to 4 hours. Gastric reflux is thus physically suppressed and the irritated sphincter allowed to recover and resume its normal competence. Being completely physical in its mode of action, Gaviscon is more efficient than chemical agents and does not carry the risk of masking symptoms. Originally developed from work carried out on X-ray contrast media, its physical action has been conclusively demonstrated in clinical trials.




'Gaviscon' prevents pain caused by acid reflux

Demulcent 'Gaviscon' covers the cardia suppressing acid reflux but should intra-abdominal pressure increase 'Gaviscon' moves up into the oesophagus soothing and protecting the mucosa in the hernial pouch. By physically preventing auto-digestion 'Gaviscon' effectively relieves the pain and allows the inflamed oesophagus to recover. In the form of 'Infant Gaviscon' the treatment has proved extremely useful in cases of gastric reflux in premature babies and infants.



■ Breaks the vicious circle of acid reflux—sphincter irritation and incompetence—acid reflux. ■ Simplifies the diagnosis of reflux oesophagitis. ■ Often postpones or avoids the need for surgery. ■ Ideal for long term prophylaxis.



GAVISCON

 Regd.

Gastric Reflux Suppressant

'Gaviscon': Supplied as chewable granules in single-dose (5 gramme) sachets containing Alginic Acid B.P.C. 0.521 g, Sodium Alginate B.P.C. 0.521 g, Magnesium Trisilicate B.P. 0.052 g, Dried Aluminium Hydroxide Gel B.P. 0.208 g, Sodium Bicarbonate B.P. 0.177 g, in a chocolate flavoured base containing 2.1 g sucrose. A single-dose is recommended after each meal and at bedtime.



Further information available on request from:
Reckitt & Colman Pharmaceutical Division, Hull

How many of the drugs you prescribe are as well documented as Alka-Seltzer?

TRADE MARK

In the booklet 'A Scientific Assessment of Alka-Seltzer' no less than forty-three sources of reference are quoted - and that represents only part of the documentation available on the product. Based on these references the facts about Alka-Seltzer are presented in explicit detail:

Physico-chemical studies—on pH, solubility, ionisation and fat-solubility, antacid and buffering action *in vitro*.

Pharmacological studies—on

antacid and buffering action *in vivo*, effect on gastric emptying, absorption, salicylate blood levels, excretion.


Therapeutic studies—on its antacid and analgesic properties.

Safety studies—on the difference between sodium acetylsalicylate and acetylsalicylic acid, gastroscopic observations, faecal blood loss, and special studies related to alcohol ingestion.

The total picture that emerges

is that of a safe and effective home remedy which is as well documented as many ethical prescription items.

For every doctor who is concerned about the subject of self-medication 'A Scientific Assessment of Alka-Seltzer' makes instructive reading. Copies available from: Dept G 11, Miles Laboratories Limited, Stoke Court, Stoke Poges, Bucks.



ORABOLIN (ethylestrenol) and phenformin

A NEW APPROACH TO THE PROPHYLAXIS OF POST-SURGICAL THROMBOEMBOLIC DISORDERS

August 1968

'phenformin when combined with ethylestrenol favourably influences four factors related to thrombosis, that is, fibrinolysis, plasma fibrinogen, platelet stickiness and serum cholesterol'.
Brit. M.J. (1968) 3, 498

February 1970

'about a third of all patients over the age of 40 undergoing an operation will develop thrombosis and (that) half of these are clinically undetectable.'
Editorial (1970) Lancet 7, 395

January 1971

'in patients who have recurrent deep vein thrombosis, treatment with phenformin and ethylestrenol which increases blood fibrinolytic activity may prove beneficial.'
Lancet (1971) 7, 242

NOW—1971

CONFIRMATION THAT ORABOLIN AND PHENFORMIN REDUCE POSTOPERATIVE FIBRINOLYTIC SHUT-DOWN

Lancet (1971) To be published.

Dosage: 4 mg Orabolin (ethylestrenol) b.d. and 50 mg Phenformin b.d. commencing three weeks before operation.

Contra-indications: Pregnancy or history of abnormal liver function tests.

Precautions: Anticoagulants, if administered, will need to be reduced in dosage, the amount of reduction being judged by the prothrombin time. In diabetes, careful control of diet and

frequent blood sugar estimations.

Side Effects: Phenformin may cause gastro-intestinal upsets; this complication is reduced by the use of delayed-action capsules. Long-term ethylestrenol therapy may cause amenorrhoea. Liver function is not affected in the recommended therapeutic dosage.

Form and Packs: 2 mg Orabolin (ethylestrenol) tablets in bottles of 100 and 500.



For further information on this interesting development contact:

Organon Laboratories Ltd · Crown House · Morden · Surrey

ACTA GASTRO-ENTEROLOGICA BELGICA

SOMMAIRE du FASCICULE 11-1970

Bibliographie.

P. ECTORS, S. VERBEUSTEL et J. DE GRAEF (Bruxelles). Effets de l'ictère obstructif sur la sécrétion gastrique d'acide, de pepsine, de polysaccharides sulfatés et de glycoprotéines par la muqueuse fundique chez le chien.

A. GERARD, J.-P. DEREUME, A. VAN MEERBEEK, R. PARMENTIER et A. BREMER (Bruxelles). Colite ischémique nécrosante. L. DELOYERS (Bruxelles). Colite ischémique chez un homme jeune. Forme sténosante rectosigmoïdienne, suivie après 28 mois, d'une nécrose segmentaire du côlon descendant. Rôle étiologique éventuel d'une toxicomanie à l'éphédrine.

E. JACOBS (Bruxelles). A propos de deux cas de nécrose étendue de la muqueuse colique. Rôle étiologique de perturbations circulatoires et de l'agression microbienne.

P. LÉONARD (Bruxelles). Le reflux dans le canal de Wirsung. Considérations à propos d'une expérience personnelle.

SOMMAIRE du FASCICULE 12-1970

Bibliographie.

M. SELIGMAN (Paris). La maladie des chaînes alpha.

H. BAZIN (Louvain). L'immunité humorale au niveau de l'intestin.

E. LEBACQ, B. FLASSE, J.-P. DEVOGELAERE, F. SMEETS and A. BOSLY (Jolimont). Immunological changes in Crohn's disease: attempts at devising a new diagnostic test.

J. WIBRAN (Bruxelles). Immunopathologie du tube digestif.

J. M. LIMBOSCH, Mme M. C. WOUSSEN-COLLE, A. GERARD et J. DE GRAEF (Bruxelles). La sécrétion acide et de pepsine en réponse à la gastrine et à la pentagastrine chez l'homme.

G. LIGNY et J. MANOUILLOFF (Charlevoix). Action chronique d'un inhibiteur de l'acétyl-cholinestérase sur les troubles cliniques et sur la sécrétion gastrique posthistaminique de patients hypochlorhydriques.

SOMMAIRE du FASCICULE 1-1971

Bibliographie.

A. RIBET, J. P. PASCAL, et N. VAYSSE (Toulouse). Etude comparative chez l'homme de l'activité des différentes sécrétines actuellement commercialisées.

M. VAGNE, L. DESCOS, C. TREPO, M. J. AULAGNIER et C. DESVIGNES (Lyon). La sécrétion hydrocarbonatée du pancréas après administration de sécrétine par injection unique et perfusion continue chez l'homme.

S. BONFILS, J. GILSON et J. N. JUNIEN (Paris). Premiers résultats de la céruléine dans l'exploration fonctionnelle pancréatique chez l'homme.

R. VAN DER HOEDEN, A. DELCOURT et P. ALGOET (Bruxelles). Le test sécrétine-pancréozymine. Recherche d'un paramètre significatif.

R. BOUVIER (Montpellier). Technique d'exploration du pancréas exocrine par dosage de l'activité enzymatique du liquide duodénal. Ses possibilités et ses limites.

Y. LE QUINTREC, M. THOMAS et A. LAPEGUE (Paris). Résultats du test d'hyperglyco-amylasémie provoquée dans les affections du pancréas exocrine.

M. CARBONI, N. PATRASSI, M. SIMI, E. LEZOCHÉ et C. MONTESANI (Rome). Le test glyco-amylasémique: son importance dans le diagnostic des pancréatites chroniques.

M. CARBONI, N. PATRASSI, M. SIMI, C. MONTESANI et G. PERROTTA (Rome). Pathologie pancréatique et amylase: étude clinique et expérimentale.

L. LEGER, P. FABIANI et J. P. LENRIOT (Paris). Dosages de lipase en pathologie pancréatique: valeur diagnostique et pronostique.

J. PARIJS, G. A. DE WEERDT et F. BARBIER (Gent). Lipase pancréatique et lipolyse des triglycérides; vitesse de formation d'acides gras libres saturés et non saturés en fonction de la température et de la concentration de calcium.

J. HUREAU, E. VAIREL, C. RABY et P. FORLOT (Paris). Les enzymes protéolytiques dans les pancréatites; présence et conséquences, valeur diagnostique.

R. KIEKENS, J. P. GOVAERTS et J. VAN GEERTRUYDEN (Bruxelles). Intérêt du dosage de la méthémalbumine dans le diagnostic et la thérapeutique de la pancréatite aiguë.

Y. GEFFROY, J. BOURREILLE et J. B. AUBET (Rouen). Les perturbations de l'homéostasie sérotoninique au cours de la pancréatite chronique récurrente. Rapports éventuels.

P. JACQUEMET (Lyon). Place de la duodénohypotonie dans le diagnostic des affections du pancréas exocrine.

CL. GUIEN et R. CAMATTE (Marseille). Valeur diagnostique et étude comparative des signes radiologiques des techniques standard dans les affections du pancréas.

R. CRISMER, M. DELFORGE et P. DEFRAANCE (Liège). Rendement clinique de la tomographie frontale du pancréas après pneumo-rétroperitoneumopéritoine.

H. PIETRI, J. BOURDE et G. HAGE (Marseille). L'exploration du pancréas par tomographie sagittale sous pneumopéritoine et rétroperitoneumopéritoine associés. Aspects normaux.

CL. GAUTIER-BENOIT et CL. L'HERMINE (Lille). Intérêt de l'artériographie bisélective coelio-mésentérique en chirurgie pancréatique.

J. PLESSIER (Paris). Confrontation entre la valeur de la parenchymatographie, la capacité exocrine du pancréas, l'insulinémie et la glucagonémie obtenues par l'injection intracoeliale de sécrétine purifiée.

R. STOPPA et J. PIETRO (Amiens). Lésions chirurgicales des artères viscérales au cours des pancréatites chroniques.


L. LEGER, J. C. ROUCAVROL et J. P. LENRIOT (Paris). Scintigraphies à la sélénométhionine dans les pancréatites aiguës et chroniques.

A. DELCOURT (Bruxelles). Conclusions du modérateur.

Acta Gastro-enterologica Belgica: 10 fascicules par an. Abonnements (Belgique et étranger: 1.200 Fr. b.-le fascicule isolé: 250 Fr. b. Secrétariat et rédaction: rue des Champs Elysées. 43. B 1050. Bruxelles. Belgique.

***The combative role
of Noxyflex against bacteria
is seen to be singularly effective
when compared
with the whole barrage
of alternative treatments***

When Noxyflex is applied directly to the site of existing or potential infection its effectiveness is undeniable. The latest Noxyflex literature contains a comparison of 'percentages' which points to the superiority of Noxyflex when compared with alternative treatments. Have you a copy?



Noxyflex
Geistlich Chester

2.5G. Noxytiolin and 10 mgm. Amethocaine HC1

RIFOCIN[®]-M


rifamide

LEPETIT

in BILIARY TRACT INFECTIONS

"... it is unique in its action in acute cholecystitis and other bacterial diseases of the biliary tract, for which it is the only specific antibiotic."¹

"A much higher proportion of the organisms found in cholangitis would be expected to be inhibited by this drug than by other antibiotics that achieve effective bile levels..."²



- * Active against penicillinase-producing organisms.
- * No cross-resistance or cross-allergenicity with other antibiotics has been demonstrated.
- * Well tolerated. No toxic reactions have been reported.
- * Dose: 150 mg intramuscularly every 8-12 hours.

RIFOCIN-M (rifamide) is a member of the RIFAMYCINS—a new group of antibiotics discovered and developed in the Research Laboratories of LEPETIT.

References: 1. *Med. J. Aust.*, 1966, **1**, 1-7. 2. *Brit. J. Pharmacol.*, 1967, **31**, 506-512.

Full information is available on request.



LEPETIT Pharmaceuticals Ltd.

Nicholson House, High Street, Maidenhead, Berks. Tel: Maidenhead 32277

Calm for the stomach of the ulcer prone.

For the peptic ulcer patient
hyperacidity and hypermotility can mean pain.

Nacton forte relieves this pain by its vagal blocking
action. Long-lasting pain relief is ensured,
and an environment which encourages healing.

There are no side effects at therapeutic doses.
Modern, effective treatment for peptic ulceration.

Nacton forte  **Bencard**

Full information on Nacton* forte (poldine methylsulphate B.P.)
is available on request from:

Bencard • Great West Road • Brentford • Middlesex.

*regd.



Selected Writings of Sir Arthur Hurst (1879-1944)

Wordsworth said of Coleridge that he was like a torch to set other men alight and the same could be truly said of Sir Arthur Hurst. His case demonstrations drew an audience, not merely from Guy's Hospital and London, but from all over the world. It was characteristic that his younger colleagues took an active part in these demonstrations. He inspired many to engage with him in physiological and clinical research work in gastroenterology.

The originality and solidity of his contributions to gastroenterology have sometimes been underestimated just because of his other brilliant gifts. The present volume, which contains only a fraction of his published work, will correct any such misapprehension. It is surprising how much of our present knowledge of gastroenterology goes back to Hurst, as a glance at the Bibliography will show. Moreover, he set an example of the continuous process of discovery and re-appraisal, without which a subject cannot flourish. His inspiration has been transmitted directly to the present generation of gastroenterologists in Great Britain but ultimately it will have to depend on the written word. This memorial volume reveals the reasons for the pride and affection with which the British Society of Gastroenterology looks back to its founder, and ensures, as far as is humanly possible, that his personality and accomplishments will not be forgotten.

The British Society of Gastroenterology is grateful to Dr. Thomas Hunt for having prepared this memorial volume on its founder while many of his friends are still alive and his image is still vivid in our minds. The task has been a considerable one and has been notably well completed.

Price £2.00 (U.S.A. \$5.00) net

Copies can be ordered now direct from

The Publishing Manager, GUT, BMA House, Tavistock Square,
London WC1H 9JR, or through any leading bookseller

TODAY'S DRUGS

The problem facing the prescribing doctor today is choice. Catalogues of drugs—no matter how detailed—are of little help unless accompanied by advice on which drug to give for what condition. So this edition of "Today's Drugs" not only gives a full account of the drugs but also includes clear accounts of the management of common conditions such as urinary infections and epilepsy.

This edition differs from the earlier one in other ways. It deals with only three subjects—*infections, vitamin deficiencies and anaemias, and disorders of the nervous system.* Further volumes will be published at intervals of about a year, and by the time the whole field of therapeutics has been covered this first volume will have become obsolete and due for replacement. In order to further emphasize the ephemeral nature of books on therapeutics—and to cut costs—the book is a paperback.

Demy Octavo

Paperback

224 pages

Price £1.00 (U.S.A. \$3.00) including postage

ORDER YOUR COPY NOW FROM

The Publisher, **BRITISH MEDICAL JOURNAL**, B.M.A. House,
Tavistock Square, London WC1H 9JR, *or through any leading bookseller*

Top prices offered for Back Runs of the following B.M.A. Journals:

Gut

Annals of the Rheumatic Diseases

Archives of Disease in Childhood

British Journal of Ophthalmology

British Journal of Venereal Diseases

Journal of Neurology, Neurosurgery and Psychiatry

Wm. DAWSON & SONS LTD.
BACK ISSUES DEPT.
CANNON HOUSE
PARK FARM ROAD
FOLKESTONE
KENT

Pathology of Trauma

The Proceedings of a Symposium
organized by the
Royal College of Pathologists

Edited by
S. Sevitt and H. B. Stoner

Price £2.00 (U.S.A. \$5.00)
including postage

The Publishing Manager, **JOURNAL OF CLINICAL PATHOLOGY**, B.M.A. House,
Tavistock Square, London WC1H 9JR



Digestion

Index
Vol. 3, No. 2
1970

Smoldas, J. (Olomouc):
Some Problems of the Ultrasonic Diagnosis of Bile Stones

Pirk, F. and Skála, I. (Prague):
Motility of the Digestive Tract After Administration of Medium Chain Triglycerides (MCT) as Compared with Long Chain Triglycerides (LCT)

Pedersen, S. A. (Odense):
Pentagastrin as a Stimulant in Routine Clinical Testing of Gastric Acid Secretion

Elder, J. B. (Glasgow):
Achalasia of the Cardia in Childhood

Beno, I.; Bučko, A.; Chorváthová, V. and Babala, J. (Bratislava):
Absorption of d-xylose and Fat and Histological Changes of the Small Intestine in Patients after Partial Gastrectomy

Mauer, I. and Levitan, R. (Chicago, IL):
Regional Enteritis and Celiac Artery Stenosis with Negative Small Bowel Studies. A Case Report

Bianchi Porro, G. and Maiolo, A. T. (Milano):
Cerebral Hemodynamics and Metabolism in Hepatic Cirrhosis. Effects of Portacaval Shunt Operation

EDITORIAL

Berndt, H. (Berlin):
Malabsorption in Skin Diseases

BOOK REVIEWS

Colombi, A. (Basel):
Early Diagnosis of Fatal Hepatitis

Steinitz, H.; Tals, B. and Stein, B. (Tel-Aviv):
Entamoeba histolytica and Dientamoeba fragilis and the Syndrome of Chronic Recurrent Intestinal Amoebiasis in Israel

Takada, A.; Kobayashi, K. and Takeuchi, J. (Kazanawa):
Gastroenteric Clearance of Albumin in Liver Cirrhosis; Relative Protein-losing Gastroenteropathy

Hoedemacker, Ph.J. (Groningen):
Heterotopic Gastric Mucosa in the Duodenum

Penzes, L. (Budapest):
Data of the Chemical Composition of the Ageing Intestine

CASE REPORT

Chowers, I.; Eyal, Z. and Saltz, N. J. (Jerusalem):
Cushing's Syndrome Associated with Polyps of the Colon

EDITORIAL

Berndt, H. (Berlin-Buch):
Is Appendectomy Followed by Increased Cancer Risk?

BOOK REVIEWS


NEWS

Ask for
specimen copies

All orders from the U.K. to be placed with
J. Wiley and Sons Ltd., Baffins Lane, Chichester, Sussex

S. Karger Basel (Switzerland) New York

Photograph by courtesy of B.O.A.C.



your journal by air mail

Many subscribers prefer to receive their copy of *GUT* within a few days of its publication by having it sent by air mail.

This is a service for all subscribers at the following annual rates which are charged in advance and in addition to the normal annual subscription rate.

If you reside in		Air Mail Charges
Area A (Examples)	Middle East, Cyprus, and Spanish N. & W. Africa etc.	£5.50
Area B (Examples)	Rest of World, including North America etc.	£7.25 (U.S.A. \$17.00)
Area C (Examples)	Australasia & Far East etc.	£8.50

Subscribers who wish to take advantage of this air mail service should write direct to The Subscription Manager, *GUT*, B.M.A. House, Tavistock Square, London WC1H 9JR, England.

ACTA GASTROENTEROLOGICA LATINOAMERICANA

VOL. 2 - No. 4

OCTUBRE-DICIEMBRE 1970

TRABAJOS ORIGINALES: ORIGINAL ARTICLES:

- Efectos Metabólicos de la norepinefrina sobre el hígado aislado de la rata. (Metabolic effects of norepinephrine on isolated perfused rat liver).* A. LEMBERG, R. WIKINSKI, E. M. IZURIETA, H. HALPERIN, M. PEREZ DE NEUMAN, P. LEMBERG, H. JAUREGUI, N. JAMARDO 179
- Actividad disacaridásica del intestino delgado de la rata: Su relación con la edad y la alimentación. (Disaccharidase activity of the rat small intestine: its relation with age and feeding).* R. LICASTRO, H. N. TOCCALINO 189
- Gastric Secretion Studies in Hypoxic rats. (Estudio de la secreción gástrica en la rata con hipoxia).* A. ALONSO, B. CANNATA, C. ZEOLITE, R. BORH, J. R. E. SUÁREZ 193
- Gastric Secretion study on dogs continuously perfused with histamine. (Estudio de la secreción gástrica en perros con perfusión continua de histamina).* B. CANNATA, A. ALONSO, C. ZEOLITE, R. BORN, E. FIGUEROA, J. R. E. SUÁREZ 197
- Actividad proteolítica en las gastritis atroficas. Evaluación del aporte terapéutico de un preparado digestivo. (Proteolytic activity in atrophic gastritis: Therapeutic evaluation of a digestive preparation).* M. SCHRAIER, P. A. MAZURE, J. N. COSEN 201

EDITORIAL

- Antígenos asociados a la hepatitis viral (Viral hepatitis associated antigens).* P. E. MORGANTE 207
- Actualización - Review
- Motilidad de las Vías Biliares. (Biliary tract motility).* M. ROYER, H. PARODI 211

LIBROS NUEVOS: BOOK REVIEWS

- Eponimic Syndroms and Disease and their Synonyms,* por S. JABLONSKI
- Actuelle Hepatologie,* por H. A. KÜHN, H. LIEHR (Edit.)
- Progress in Gastroenterology.* Vol. II, por G. B. Jersey Glass (Edit.)
- Bibliografía Gastroenterológica Latinoamericana. (Latinoamerican Gastroenterologic References)

NOTICIAS: NOTICES

Acta Gastroenterologica Latinoamericana is a tri-monthly publication sponsored by the *Asociación Interamericana de Gastroenterología*. It is edited by a non profit organization. The Editorial Board is formed by relevant gastroenterologists from all the Latin American countries.

Papers are published in Spanish, Portuguese, English and French languages. An English abstract gives precise statements of the studies done in each paper.

Subscription price \$9.00 USA per year.

Address: Instituto Nacional de la Salud, Martinez de Hoz y Perdriel, Haedo, Rep. Argentina.