in the treatment of peptic ulcer

ROTER

has <u>particular</u> advantages





ROTER achieves a high cure rate and often gives effective control of even refractory chronic cases which have resisted other treatments.

ROTER in many cases obviates the need for hospitalization or surgical intervention.

ROTER helps patients to keep ambulant and symptom-free with a minimum of medication and with few, if any, dietary restrictions.

ROTER in general practice enables the physician to take adequate care of his gastric cases without frequent or prolonged supervision.

ROTER has no known side effects.

ROTER requires no supplementary medication such as sedatives or anti-spasmodics.

ROTER shows a saving in cost when compared with conventional methods of treatment.

REFERENCES

Therapie der Gegenwart, 1953, 3, 108 Gazz. Med. Ital., 1954, 113, 343 Die Medizinische, 1955, 4, 163 British Medical Journal, 1955, 2, 827 C.R.S. Biol., 1956, 150, 1019 Am J. Gastroenterology, 1957, 28, 439 Wiener Med. W., 1958, 108, 218 Ärztliche Praxis, 1958, 10, 766 Journal Lancet, 1960, 80, 37

FORMULA

Each Roter tablet contains:
Bismuth Subnitrate Roter (350 mg.), Magnesium
Carbonate (400 mg.), Sodium Bicarbonate
(200 mg.) and Cortex Rhamni Frangulae (25 mg.).

PACKINGS OF ROTER TABLETS

Tins of 40 and 120, also dispensing sizes, 360 and 720 (P.T. exempt). Basic N.H.S. price: Under 1/- for 10 tablets.

ROTER TABLETS are in Category "S". Literature on Request



"I cannot eat but lytel mete My stomach is not goode..."

In diarrhoeic conditions due to infection, clinical experience has shown 'Sulphamezathine' to be far superior to the non-absorbable sulphonamides.

'Sulphamezathine' can be given with safety

to patients of all ages, and renal complications are most unlikely to occur.*

Furthermore, 'Sulphamezathine' dispenses with the need for using antibiotics—especially those of the costly "broad-spectrum" group.

* Brit. Nat. Form. (Stand. Edn.) 1960, p.45.

Sulphamezathine (Sulphadimidine B.P.)

Basic N.H.S. cost: 1/5d. per 25 tablets (0.5 g.). Suspension (100 ml.) 3/4d.





SURGILAR

ATRAUMATIC **NEEDLE SUTURES**

New! Improved!

IMPROVED ATTACHMENT!

New process gives greater attachment strength between needle and suture-eliminates ridge at point of closure ensuring a smoother line of flow reducing tissue trauma.

IMPROVED NEEDLE

Stronger, more evenly tempered to retain its shape during use. Point and cutting edges kept perfectly sharp in protective, plastic envelope.

IMPROVED CATGUT

Minimum diameter variation within and between strands of gut. Tensile strength considerably in excess of required standards. Unkinked-far more flexible.

... now with smoother line of flow

> Contained in envelope ready for immediate use. Non-irritant 'Hibitane'† chlorhexidine jar solution ensures complete

Full details of range, sizes and prices are available on application to The Sole U.K. **Distributors**





THE OLD MEDICAL SCHOOL, PARK STREET, LEEDS 1. Tel: 20085 (5 lines) Cables: 'Aseptic, Leeds 1'

38 Welbeck Street, London W.1. Tel: WELbeck 8152/3. Also at Johannesburg, Cape Town and Durban

*Regd. Trademark

** Trademark † Trademark of Imperial Chemical Industries Ltd.

MADE IN UNITED KINGDOM by CYANAMID of GREAT BRITAIN LTD

NOW IN

STERILIZED SURGICAL SURVERED ST. LICENCE F.S. WAY

JUST TEAR OPEN!

FOR CONFIDENT DIAGNOSIS IN GASTROENTEROLOGY

SECRETIN and PANCREOZYMIN

These two preparations can play important roles as diagnostic agents in gastroenterology.

They may be used with confidence in the serum enzyme test and the duodenal aspiration test.

SECRETIN

is supplied in 10 cc rubber-capped vials each containing 80-100 units. The potency is declared on each vial following biological assay.

PANCREOZYMIN

is supplied in 20 cc rubber-capped vials each containing 100 units.

Please write for further information and supplies to:

BOOTS PURE DRUG COMPANY LIMITED STATION STREET • NOTTINGHAM • ENGLAND



ASPIRIN IS SAFER WHEN IT IS SOLUBLE

Soluble aspirin is safer when it is stable



N.H.S. Basic Price 12/6 for 500 tablets in foils.

SOLPRIN provides soluble aspirin in its most stable form and dissolves rapidly in water to form a solution of calcium acetylsalicylate.


SOLPRIN is easy to take and palatable. It has proved rapid and reliable in its analgesic effect and is extremely well tolerated, even in prolonged and massive dosage.

SOLPRIN is not advertised to the public and is available only on prescription. Clinical samples and literature will be supplied on request.

SOLPRIN

Soluble and substantially neutral

RECKITT & SONS LIMITED, PHARMACEUTICAL DEPARTMENT, HULL.



Pregnancy is just one of the times when Complan can be a boon. Perhaps anorexia and nausea have prevented a normal food intake, or possibly the diet is inadequate. Complan can provide extra protein, vitamins and minerals in a most pleasant way. It may be mixed hot or cold and blends with most fruit juices and soups.

COMPLAN complete planned food to drink



1 LB. CARTONS The name Complan is a Glazo trade mar



GLAXO LABORATORIES LIMITEI

ULCERATIVE COLITIS



In Ulcerative Colitis extensive clinical experience confirms

SALAZOPYRIN

BRAND OF SULPHASALAZINE (SALICYLAZOSULPHAPYRIDINE)

to be a drug of significance in to-day's management of this distressing disease.

The present investigation suggests that salazopyrin is a useful drug Gut 1:217 (1960)

Outstanding characteristics of Salazopyrin treatment are

- Prompt remission of symptoms
- Bleeding and frequency of stools sharply reduced
- In many cases rectal mucosa heals within a month
- Abdominal cramp controlled
- Fever subsides, appetite improves
- Can be given safely over long periods of time

SALAZOPYRIN is also a safe and effective product in other torms of coliris and regional enteritis.

(R) PHARMACIA

Directional by Smer; & Moore Lide, 60'61 Williast Street, Leader, W. L.



PRODEXIN

CARRIES THE PEPTIC ULCER PATIENT SAFELY THROUGH

HIS DAY PRODEXIN is a product of BEECHAM RESEARCH LABORATORIES LTD.

BRENTFORD - ENGLAND - TELEPHONE ISLEWORTH 4111

PORTYN

brings new peace to the peptic ulcer patient



A potent new anticholinergic agent, Portyn * produces marked inhibition of gastric secretion, gastrointestinal spasm and hypermotility. It thus forms an excellent adjunct in the management of peptic ulcer and functional gastrointestinal disease.

Portyn Kapseals each contain 10 mg. benzilonium bromide. Initial adult dose—1 Kapseal three times daily. PARKE-DAVIS

Parke, Davis, Hounslow, Middlesex

*trade mark



'Streptotriad'

TRI-SULPHONAMIDE/STREPTOMYCIN

against bacillary dysentery

for the treatment of the acute phase and during the convalescent or chronic stage to prevent the patient becoming a carrier.

Each 'Streptotriad' tablet and 3.6 ml. teaspoonful of prepared suspension contains: Streptomycin (as sulphate B.P.) 65 mg.

Sulphamerazine B.P. 65 mg.


Sulphadiazine B.P. 100 mg.

Sulphathiazole B.P.C. 100 mg.



An M&B brand Medical Product

MANUFACTURED BY MAY & BAKER LTD



Pro-Banthine with Dartalan

Control of ulcer and anxiety symptoms with a single preparation


There are many situations in which patients receiving anticholinergic therapy for peptic ulcer also require an agent that will *effectively* treat associated emotional overlay. Clinical reports have shown Pro-Banthine with Dartalan to be outstandingly effective in relieving

the pain and distress characteristic of peptic ulcer and functional dyspepsias.

Packs, formula and basic NHS costs Pro-Banthine 15 mg with Dartalan 3 mg in bottles of 50, 100 and 500. Basic NHS costs 16/8, 31/8 and 146/8 respectively.

G.D.Searle & Co Ltd

High Wycombe England SR 55



Using the ENEMA in this day and age?

In the preparation of patients for operation, particularly operations on the abdomen, Dulcolax can effectively replace the enema as a method of bowel clearance. Two tablets in the evening followed by one suppository on the morning of the operation are normally sufficient. Unpleasantness for both patient and nurse is thus avoided.

Dulcolax is equally effective as a treatment for post-operative constipation. Being free from side-effects it can safely be used in pregnancy and for young and old patients alike.

If you are still using the enema, you are invited to compare it with the safety, the efficacy, the convenience of Dulcolax.

DULCOLAX*

(4,41-diacetoxy-diphenyl)-(pyridyl-2)-methane



For pre-operative intestinal sterilisation

SOFRAMYCIN TABLETS

250 mg. oral tablets

Effective against most organisms especially staphylococci, enterococci, enterobacteria including Proteus and particularly *Escherichia coli*.

"In several instances there was no growth of organisms at all in the stool after total dosages of 2 or 3 g."

"Undesirable side-effects from the drug were not found" Gut, 1960, 1, 323.



250 mg. Soframycin Oral Tablets Bottles of 20



London, N.W.10. Telephone: LADbroke 6611

STELABID


FOR THE MAN WHOSE JOB CAUSES WORRY AND GASTRIC DISTRESS




Many signalmen worry about their exacting job and are, consequently, prone to gastric disorders. For the worried signalman—and others—with a stress-induced gastro-intestinal complaint, 'Stelabid' is indicated. 'Stelabid' calms whilst it controls local gastric symptoms.


Each tablet of 'Stelabid' contains 1 mg. 'Stelazine' (brand of trifluoperazine) and 5 mg. 'Tyrimide' (brand of isopropamide).

SMITH KLINE & FRENCH LABORATORIES LIMITED WELWYN GARDEN CITY, HERTS





GABTRIC ANALYSIS 1 Superimposed gruel fractional testmeal curves of five cases of duodenal ulcer.



GASTRIC ANALYSIS 2 Same patients as in first graph, two days later, showing the striking neutralizing effect of days later, showing the striking neutralizing effect of acking Nulacin tablets (3 an hour). Note the return of ackity when Nulacin is discontinued.

"Nulacin has revolutionized the treatment of duodenal ulcer" Proc. R. Soc. Med.. 1958 \$1,1063

Nulacin tablets, dissolved slowly in the mouth, provide intragastric milk-alkali drip therapy, obviating the necessity for a tube. Treatment is easy and convenient for the ambulant as well as the bedridden patient. Nulacin may be relied on to provide an effective treatment for peptic ulcer as well as other conditions associated with gastric hyperacidity. No undesirable

side effects occur and no cases of alkalosis or acid rebound due to Nulacin have ever been reported. The safety and effectiveness of Nulacin therapy has been proven by *in vivo* clinical work carried out in many countries.


Nulacin tablets have no B.P. equivalent. The basic N.H.S. price of the 25 tablet tube is 2/-.

NULACIN is a Horlicks pharmaceutical product

For further information please write to:

HORLICKS LIMITED

PHARMACEUTICAL DIVISION, SLOUGH, BUCKS



IN ONE BOTTLE

Not really, because Aminosol is a casein hydrolysate. It does, however, provide adequate protein parenterally where oral nutrition is impracticable.

Supplied as Aminosol Glucose Vitrum (Aminosol G.V.) 307 calories per litre and

Aminosol Glucose Ethanol Vitrum (Aminosol G.E.V.)
662 calories per litre

AMINOSOL

In 500 and 1000 ml. bottles. Aminosol Glucose Vitrum also provided in 100 ml. paediatric size. Further information on request

Vitrum

PAINES & BYRNE LTD., PABYRN LABORATORIES, GREENFORD, MIDDX.

Uniquely effective in antibacterial and antiamoebic therapy

KAPSEALS

HUMATIN* is unique among broad-spectrum antibiotics in that it is highly effective against intestinal amoebiasis as well as many types of bacterial enteric infections. HUMATIN (Paromomycin, P.D. & CO.). has also been found useful in the management of hepatic coma and in suppressing, pre-operatively, the normal intestinal microflora.

Available in Kapseals* each containing the sulphate equivalent to 250 mg. paromomycin base. In bottles of 16 and 100.

Now also available as Humatin Syrup in bottles of 60 ml. and 500 ml.; each 5 ml. containing paromomycin sulphate equivalent to 125 mg. paromomycin base.

*TRADE MARK



STAPHYLOCOCCAL SUPERINFECTIONS

"Patients in hospital who have had an operation, and a course of 'broad-spectrum' antibiotics, sometimes develop diarrhoea which may be severe and even fatal."

Lancet Annotations 25th February, 1961



LACTOBACILLUS ACIDOPHILUS PREPARATION

For the prophylaxis and treatment of staphylococcal super infections.

"It was found that the staphylococci increased in both groups at the beginning of antibiotic treatment. Staphylococci continued to increase in the patients taking the antibiotic alone, but in those taking 'ENPAC' there was a pronounced drop in numbers."

Gordon D., Macrae J., & Wheater D. M. W. Lancet-May 4th, 1957. pp. 899-901

Samples and literature will be sent to the medical profession on request to:-

WILTS UNITED DAIRIES LIMITED · NUTRITION DEPARTMENT · TROWBRIDGE · WILTS



"This book will take its place as the reference volume on the surgery of the anus, rectum, and colon, both at home and overseas"—British Journal of Surgery.

Surgery of the Anus, Rectum and Colon

J. C. GOLIGHER, Ch.M., F.R.C.S.

Professor J. C. Goligher holds the chair of Surgery at the University of Leeds, and is Surgeon, General Infirmary at Leeds: formerly Surgeon St. Mary's Hospital and St. Mark's Hospital for Diseases of the Rectum and Colon, London.

This book, written especially for Surgeons in training and for young Consultants, comprehensively surveys contemporary knowledge of diseases of the rectum and colon, and at the same time provides a clear and detailed practical guide to the surgical management of these conditions. A special feature is the two chapters on developmental abnormalities of the anus and rectum and on Hirschsprung's Disease and related abnormalities by Mr. Harold Nixon, F.R.C.S., of the Hospital for Sick Children, Great Ormond Street, London. 896 pp. Crown 4to 5 colour plates 276 half-tone illustrations. 28 8s. net.

CASSELL MEDICAL BOOKS



ALUHYDE

for prompt, safe symptomatic medication in hyperchlorhydria and peptic ulcer

'ALUHYDE' tablets present a new combination with these advantages.

'ALUHYDE' effects sale reliable, nonsystemic neutralization by means of two of the most preferred modern antacid adsorbents...



ALUHYDE permits of quick mental sedative-effect by its inclusion of a barbiturate recognized for minimal cumulative potential...



'ALUHYDE' provides antispasmodic action through its content of an agent long known for its favourable influence against hypermotility...



Formula-

Each tablet: Dried Aluminium

Hydroxide Gel. B.P.C. - 3.75 gr.

Magnesium trisilicate B.P. - 3.75 gr.

Quinalbarbitone sodium B.P. - 0.50 gr.

Tinct. Belladonnæ B.P. - 3 m.

Packings—Regular packs of 50; bottles of 500 (dispensing)

Samples available on personally signed request of qualified physicians only

MULTIPAX CHEMICALS LTD

142. 144. 146 LARKHALL LANE, LONDON, S.W.4

Phones: MACaulay 8493/6821

MCL

IAST