

ROTER

In cases of dyspepsia with or without a radiologically proven ulcer Roter will normally provide prompt symptomatic relief and lead to rapid healing of the lesion. The reputation of Roter is world wide. It is a complete treatment in itself and requires no additional antispasmodics or anticholinergics.




FORMULA: Each Roter tablet contains: Bismuth Subnitrate Roter (350 mg.), Magnesium Carbonate (400 mg.), Sodium Bicarbonate (200 mg.) and Cortex Rhamni Frangulae (25 mg.).

PACKINGS OF ROTER TABLETS: Tins of 40 and 120, also dispensing sizes, 360 and 720 (P.T. exempt).

ROTER IN PEPTIC ULCER

A remarkably simple treatment in pancreatic enzyme replacement



Hitherto there has not been any reliable method of determining lipase activity in pancreatic extracts — none that ensured adequate control of fat digestion. This is now possible with Cotazym.

Cotazym (concentrated hog pancreas extract) enables known amounts of lipase to be administered easily and control to be established rapidly and accurately.

Cotazym is effective at lower dosage than has previously been possible and has, moreover, greater therapeutic activity. It has no offensive odour or taste and this, together with the simple method of its administration, makes it extremely acceptable to patients.

INDICATIONS

Fibrocystic disease
Pancreatic deficiency states
Post-gastrectomy malabsorption
Steatorrhoea

COTAZYM




Each Cotazym capsule contains:

Lipase — sufficient to digest 17G. dietary fat
Trypsin — sufficient to digest 34G. dietary protein
Amylase — sufficient to digest 40G. dietary starch

Bottles of 100 capsules — Basic N.H.S. Price 33/4d

Dosage — 1-3 capsules with each meal or as directed by the physician.



Organon Laboratories Ltd., Crown House, London Road, Morden, Surrey

**In peptic ulcer and functional gastro-intestinal disease,
NEW sustained-release anticholinergic PORTYN TEMPLETS
inhibit gastric secretion, gastro-intestinal spasm or hypermotility**

The sustained-release feature of PORTYN* TEMPLETS* provides the advantages of less frequent dosage and fewer and more mild side effects than with the immediate release form. Only one PORTYN TEMPLET twice daily at 12-hour intervals provides sustained inhibition of gastric secretion and gastro-intestinal hypermotility.

Available in bottles of 15 and 100.

PORTYN TEMPLETS

**Trade mark*



PORTYN TEMPLETS each contain 35 mg. benzilium bromide, Parke-Davis, (5 mg. for immediate release and 30 mg. for gradual release)

PARKE-DAVIS HOUNSLOW · LONDON · TELEPHONE: HOUNSLOW 2361

The success of **BIOGASTRONE** (carbenoxolone sodium) in gastric ulcer

further proof in 1965



“It is the only drug therapy for gastric ulcer which has been demonstrated conclusively to accelerate healing,

which can be achieved with patients remaining up and about, often at work, and with minimum change in diet”

from editorial synopsis: GUT (1965) 6:19



In 1959, the first clinical trial of Biogastrone* was begun in patients with gastric ulcer. The results indicated that Biogastrone significantly reduced the size of ulcer niche.

“It is suggested that Biogastrone promotes the healing of gastric ulcers.”

Lancet (1962) ii: 793

Now, in a further trial, the same author proves beyond any doubt that Biogastrone is successful in assisting gastric ulcers to heal. The average degree of healing was similar in both trials—72 and 78% in the Biogastrone groups and 35 and 39% for the control.

“It is concluded that carbenoxolone sodium [Biogastrone] facilitates the healing of gastric ulcers.”

Gut (1965) 6: 19

More detailed information from
**Berk Pharmaceuticals Limited, Catteshall Lane
Godalming, Surrey.**

* trade mark BR 3775/J817

VELACTIN

WANDER

A readily assimilated milk substitute
of vegetable origin for Milk Allergies



1111

Velactin contains no gluten or lactose. The biological value of the soya protein has been improved by the addition of DL-Methionine to make it equivalent to that of first-class animal protein.

Literature gladly sent on request

A. Wander Limited,
42 Upper Grosvenor Street,
London W.1.

Indications: Allergy or intolerance to human or cow's milk.¹ As basic feeding pending identification of specific food allergens.¹ Galactosaemia.^{2,3} Gluten enteropathy. Velactin may also be tried in cases of undetermined metabolic or non-specific intolerance.

References: (1) *Nutritio et Dieta* 1961; **3**, 89.

(2) *Lancet* 1959, **2** 849; 1960 **1**, 336.

(3) *Ugeskrift f. Læger* 1960, **122**, 1229.

Presentation: Tin containing 1 lb. net.



LONGER-LASTING DRIP THERAPY

Gastrils—first ever antacid pastilles. Gastrils do not break up in the mouth like tablets; they dissolve slowly and smoothly, so give sustained drip therapy for the treatment of peptic ulcer.

Gastrils contain a co-dried gel of aluminium hydroxide and magnesium carbonate, a buffer antacid comparable in activity to freshly prepared liquid aluminium hydroxide gel. Gastrils rapidly buffer gastric pH to an optimum level, and keep it there, without risk of acid rebound.

To relieve the tedium of constant antacid therapy, every Gastrils prescription is dispensed in equal quantities of two delicious flavours—mint and fruit. Few antacids are as effective as Gastrils—none is as good to take!

GASTRILS

ANTACID PASTILLES

INDICATIONS Hyperacidity, peptic ulcer, gastritis, heartburn, oesophagitis and hiatus hernia.

FORMULA Each pastille contains 500 mg. co-dried gel of aluminium hydroxide and magnesium carbonate.

BASIC N.H.S. COST 24 individually wrapped Gastrils in carton 3/-.
24 individually wrapped Gastrils from dispensing pack: 2/6½d. Gastrils are prescribable on form E.C.10

'Gastrils' is a registered trade mark of T. J. Smith & Nephew Ltd.



**Smith & Nephew
Pharmaceuticals Limited**

WELWYN GARDEN CITY · HERTFORDSHIRE · ENGLAND

NEW BLACKWELL BOOKS

The Small Intestine

Edited by A. C. THACKRAY M.D. and F. AVERY JONES M.D., F.R.C.P.

September 1965. 144 pages, 18 illustrations.

22s. 6d.

These papers were given at a symposium of the Fifth Congress of the International Academy of Pathology: they are intended to bring up to date our knowledge of various aspects of this once neglected part of the body.

Contributors: J. BADENOCH, B. CREAMER, R. C. CURRAN, J. L. EMERY, F. A. FAIRWEATHER, P. FARNAN, J. FINE, A. P. GOFFE, C. J. HODSON, B. C. MORSON, J. W. PAULLEY, W. S. PEART, W. SANDRITTER, MARGOT SHINER, J. A. WILLIAMS.

Therapeutic Agents and the Liver

Edited by NEIL MCINTYRE B.Sc., M.B., M.R.C.P. and SHEILA SHERLOCK M.D., F.R.C.P., F.R.C.P.E.

1965. 204 pages, 73 illustrations.

45s.

Thirty-six experts discuss this subject under three main headings: Effect of drugs on hepatic transport, conjugation and biliary secretion; Drug reactions in the human liver; and Alcohol and the liver.

Radiology of the Digestive System

K. LUMSDEN M.A., M.B., B.Chir., D.M.R.E. and S. C. TRUELOVE M.A., M.D., F.R.C.P.

September 1965. 548 pages, 359 illustrations.

£6

This radiological companion to Truelove and Reynell's *Diseases of the Digestive System* is primarily intended for physicians and surgeons, but will also be of value to the radiologist.

The Biliary System

A Symposium of the NATO Advanced Study Institute, edited by W. TAYLOR B.Sc., PH.D.

1965. 736 pages, 291 illustrations.

£6 15s.

The authors include anatomists, physiologists, biochemists, physicians, surgeons and radiologists who describe the present state of knowledge and the research trends of particular aspects of the biliary system.

Detailed leaflets available for all books on request.

BLACKWELL SCIENTIFIC PUBLICATIONS · OXFORD

ANTIBIOTICS AND THE INTESTINAL FLORA

Treatment with broad spectrum antibiotics suppresses the normal flora, which may be replaced by abnormal resistant organisms causing diarrhoea.

This change can be prevented or reversed by the implantation of *Lactobacillus acidophilus* which has been rendered antibiotic-resistant, and is thus able to multiply freely under these abnormal conditions. The products of growth of this organism are inimical to that of many other bacteria^{1,2} and thus contribute to its rapid preponderance over other species.

Oral treatment with antibiotic-resistant *L. acidophilus* in the form of ENPAC has been shown to eliminate *Candida* from the faeces of infants with antibiotic-induced diarrhoea, with relief of symptoms succeeded by rapid gain in weight^{3,4} and greatly to reduce the numbers of staphylococci in the faeces of patients treated with tetracyclines.⁵

¹(1958) *Ann. Inst. Pasteur* **95**, 194.

²(1959) *J. Bact.* **78**, 477.

³(1957) *Klin. Wschr.* **35**, 198.

⁴(1959) *Medizinische* **7**, 296.

⁵(1957) *Lancet* (i), 899.

ENPAC

LACTOBACILLUS ACIDOPHILUS PREPARATION

Samples and literature will be sent to the medical profession on request to:—

APLIN and BARRETT LIMITED · BIOLOGICALS DIVISION · TROWBRIDGE · WILTS



FIBER OPTIC COOL ILLUMINATED


PHOTOGRAPHY

provides for recording excellent views of the entire esophagus and cardio-esophageal junction with maximum patient comfort. Detailed pathology easily identified with colored, still or motion pictures through the Lo Presti Fiber Optic Foroblique Esophagoscope



American Cystoscope Makers, Inc.

8 Pelham Parkway, Pelham Manor (Pelham), N.Y. 10803



What is SURGICEL*?

SURGICEL is a haemostatic gauze made from oxidised regenerated cellulose. In the presence of blood, SURGICEL swells and turns into a gelatinous mass, forming an aggregate in the mouth of blood vessels and so assisting the normal process of clotting.

Where is SURGICEL used?

One of the most dramatic fields of usefulness for oxidised regenerated cellulose is found in Cardio-vascular Surgery. To arrest haemorrhage in surgery; wounds of the liver, spleen and kidney; Haemophiliac conditions and other blood clotting defects.

The Advantages of SURGICEL

rapid, reliable action—controls capillary bleeding. Effective in minimal amounts.

easy to handle—no preparation required

before use; easily sutures in place; conforms to contours of the viscera. SURGICEL does not adhere to gloves or instruments.

complete absorption—trials show that there is no gross evidence of SURGICEL 44 days after use and only minimal tissue reaction. SURGICEL causes no toxicity, intolerance or sensitivity reactions; it does not deteriorate on storage.


Presentation

SURGICEL is supplied sterile in Cartons of 3 vials available in the following sizes: strips of 2" x 14"; 4" x 8"; 2" x 3". For Dental Surgery—cartons of 12 sterile vials ½" x 2".

*trademark

ETHICON
LIMITED

Bankhead Avenue, Sighthill, Edinburgh 11



Whether you call it 'mucous colitis', 'spastic colon' or 'irritable colon' the clinical picture is the same – a chronic colonic disturbance with spasm/pain, disordered bowel habit (constipation or diarrhoea) and sometimes the passage of mucus per rectum.

Merbentyl (dicyclomine hydrochloride tablets B.N.F.) can relieve spasm/pain and help restore normal bowel activity in a very short space of time.

Merbentyl

tablets of dicyclomine BNF

Dosage

See British National Formulary 1963, Tablets – page 171.

Low basic N.H.S. Cost

An average week's treatment – 2s. 8d.

Also available with phenobarbitone and as Merbentyl Syrup.

Full clinical information available on request.

Merrell-National (Laboratories) Ltd.
London W.1

Trademarks:

Merbentyl, Merrell-National



A12068



a masked disease

Pancreatitis may be difficult to differentiate from a coronary thrombosis, perforated peptic ulcer, a dissecting aneurysm of the aorta or acute cholecystitis, but when the diagnosis is established, TRASYLOL the proteinase inactivator is indicated.

TRASYLOL specifically inhibits the proteolytic enzymes of the pancreas, and administered in adequate dosage arrests the autodigestive processes. A recent analysis of all cases of pancreatitis, those treated with TRASYLOL showed a reduction in mortality from 22% to 9%.


TRASYLOL is therefore indicated in the treatment of acute pancreatitis, pancreatic fistulae, and pancreatitis accompanying disorders of the gall bladder and bile ducts.

TRASYLOL®

Proteinase Inactivator



® Registered trade mark of **BAYER GERMANY**
U.K. Subsidiary **FBA PHARMACEUTICALS LTD**, Haywards Heath, Sussex



**2 TABLETS TWICE A DAY—
PROVEN MAINTENANCE THERAPY**

In Ulcerative Colitis

available as a 0.5 g. tablet, either plain or enteric coated

FIRST DEMONSTRATION . . . RESULTS SIGNIFICANT

"This is the first demonstration in a formal trial that any treatment reduces the relapse-rate in ulcerative colitis, 24 out of 34 patients taking 2 g. of sulphasalazine daily remained free of symptoms for a year. This result is significantly different from that obtained in a comparable group of patients who received a placebo."

PREFERABLE TO SYSTEMIC CORTICOSTEROIDS

"We have shown that sulphasalazine, 0.5 g. four times daily, is often effective in maintaining remission . . . and it therefore appears preferable to systemic corticosteroid drugs for this purpose."

REMISSION FOR ONE YEAR

"24 patients remained in symptomatic remission for a year while taking 2 g. of sulphasalazine daily, whereas only 8 remained symptom-free in the placebo group."

NORMAL MUCOSA

"2 of the 24 patients who remained symptom-free for a year while on sulphasalazine had a haemorrhagic mucosa at the end of the trial. The remaining 22 patients had a non-haemorrhagic mucosa which in many cases appeared normal."

MINIMAL SIDE EFFECTS

"In the present trial, using a dose of 2 g. daily, only 3 patients out of 34 had to discontinue treatment because of side-effects."

WHITE CELL COUNT

"In the patients treated with sulphasalazine there was no difference in the haemoglobin level before and after treatment; but the mean white-cell count was lower after six months or a year than at the start of treatment, though in no patient was it less than 4500 per c.mm."

The Lancet Jan 23rd 1965 • Vol I • Pages 185-188

SALAZOPYRIN




PHARMACIA
(GREAT BRITAIN) LTD

Literature and further information are supplied on request

SINCLAIR HOUSE • THE AVENUE • WEST EALING • LONDON W13

Distributed by Martindale Samore Ltd • 60-61 Welbeck St • London W1

AMINO SOL 10% + INTRALIPID (FAT EMULSION)



FOR COMPLETE PARENTERAL NUTRITION

... without risk of Thrombophlebitis



literature and suggested regimes on request

PAINES & BYRNE LTD., GREENFORD, MIDDLESEX
pre-eminent in Parenteral Nutrition

BRITISH MEDICAL JOURNAL



A comprehensive general medical journal. Essential for every medical man or student who wishes to keep informed of research in medical science, of trends in clinical medicine and of professional activities.

Weekly. Annual Subscription, United Kingdom, Irish Republic, £9 9s.; Abroad £5 5s. (\$15.00). and its associated specialist journals

BRITISH JOURNAL OF OPHTHALMOLOGY

Designed for the clinician and research worker and includes original articles, case notes, information on new appliances, book reviews.

Monthly. Annual Subscription, Inland £6 6s.; Overseas £7 (\$20.00). (Combined subscription for the two Ophthalmic journals, Inland £11 11s.; Overseas £13 (\$37.00)).

OPHTHALMIC LITERATURE

Abstracts of articles on the wide field of ophthalmology published throughout the world. Each volume contains some 5,000 abstracts—covering one year.

Six issues a year plus index. Annual Subscription, Inland £6 6s.; Overseas £7 (\$20.00). (For combined subscriptions see above.)

JOURNAL OF CLINICAL PATHOLOGY

Original and specially commissioned articles on each branch of pathology, with prominence given to its clinical application. Technical methods, notes on new or improved apparatus, book reviews.

Annual Subscription (six issues), Inland £4 4s.; Overseas £4 15s. (\$14.00).

BRITISH HEART JOURNAL

Original work on the heart and circulation—*anatomical, physiological and pathological.*

Annual Subscription (six issues), Inland £4 4s.; Overseas £4 15s. (\$14.00).

ARCHIVES OF DISEASE IN CHILDHOOD

Devoted to child health and disease, with emphasis on clinical paediatrics. Original articles by paediatricians and other specialists.

Annual Subscription (six issues), Inland £4 4s.; Overseas £4 15s. (\$14.00).

BRITISH JOURNAL OF PHARMACOLOGY AND CHEMOTHERAPY

Reports on original work in all branches of the subject, including experimental chemotherapy. Biochemical and pathological studies.

Annual Subscription (six issues), Inland £10 10s.; Overseas £11 (\$32.00).

JOURNAL OF NEUROLOGY, NEUROSURGERY AND PSYCHIATRY

Original articles on current research, throwing light on practice in neurology, neurosurgery and psychiatry.

Annual Subscription (six issues), Inland £4 4s.; Overseas £4 15s. (\$14.00).

ANNALS OF THE RHEUMATIC DISEASES

Original articles by authorities in the U.K., U.S.A. and elsewhere on every manifestation of the rheumatic diseases. Summaries in English, French and Spanish.

Annual Subscription (six issues), Inland £4 4s.; Overseas £4 15s. (\$14.00).

THORAX

Original work on the anatomy, physiology and pathology of the chest and heart. Descriptions of techniques and apparatus used in modern cardiac surgery.

Annual Subscription (six issues), Inland £4 4s.; Overseas £4 15s. (\$14.00).

GUT

Journal of the British Society of Gastroenterology. A journal for physicians, surgeons, radiologists, and pathologists, carrying authoritative articles on all aspects of gastroenterology.

Annual Subscription (six issues), Inland £4 4s.; Overseas £4 15s. (\$14.00).

JOURNAL OF MEDICAL GENETICS

A new journal publishing articles on clinical, statistical, pathological, cytological and biochemical aspects of medical genetics.

Quarterly. Annual Subscription, Inland £3 3s.; Overseas £3 10s. (\$10.00).

BRITISH JOURNAL OF VENEREAL DISEASES

Designed to disseminate, exchange and increase knowledge among consultants, practitioners, research workers, teachers and others concerned with the problem of venereal diseases.

Quarterly. Annual Subscription, Inland £3 3s.; Overseas £3 10s. (\$10.00).

BRITISH JOURNAL OF INDUSTRIAL MEDICINE

Authoritative papers on all aspects of occupational medicine, e.g., the pneumoconioses, occupational cancers, industrial toxicology and psychology.

Quarterly. Annual Subscription, Inland £3 3s.; Overseas £3 10s. (\$10.00).

BRITISH JOURNAL OF PREVENTIVE AND SOCIAL MEDICINE

Original work in the field, the emphasis on epidemiological methods and medical research.

Quarterly. Annual Subscription, Inland £3 3s.; Overseas £3 10s. (\$10.00).

MEDICAL AND BIOLOGICAL ILLUSTRATION

Illustration and its application to the requirements of the medical and biological sciences.


Quarterly. Annual Subscription, Inland £3 3s.; Overseas £3 10s. (\$10.00).

ABSTRACTS OF WORLD MEDICINE

Critical abstracts of important articles appearing in medical periodicals in every part of the world. Accent on what is new.

Monthly. Annual Subscription, Inland and Overseas £6 6s. (\$18.00).

Write to The Subscription Manager, BRITISH MEDICAL JOURNAL,
B.M.A. House, Tavistock Square, London W.C.1
or order through any leading subscription agent or bookseller
(All dollar rates quoted refer to the U.S. dollar)



for prompt safe
symptomatic
medication in
hyperchlorhydria
and peptic ulcer

• ALUHYDE •

Tablets present a combination with these advantages...

• ALUHYDE •

Effects safe, reliable, non-systemic neutralization by means of two of the most referred antacid adsorbents...



• ALUHYDE •

Permits a favourable enteric effect by its inclusion of a barbiturate recognized for minimal cumulative potential...



• ALUHYDE •

Provides antispasmodic action through its content of an agent long known for its favourable influence against hypermotility...



Formula—

Each tablet: Dried Aluminium

Hydroxide Gel, B.P.C. — 3.75 gr.

Magnesium trisilicate B.P. — 3.75 gr.

Quinalbarbitone sodium B.P.—0.50 gr.

Tinct. Belladonnae B.P. — 3.m.

Packings—Regular packs of 50,
bottles of 500 (dispensing)

Samples available on personally signed request
of qualified physicians only

Multipax Chemicals Ltd

121 NORWOOD ROAD HERNE HILL LONDON S.E.24. Phone: TULse Hill 3276

