

# ROTER

In cases of dyspepsia with or without a radiologically proven ulcer Roter will normally provide prompt symptomatic relief and lead to rapid healing of the lesion. The reputation of Roter is world wide. It is a complete treatment in itself and requires no additional antispasmodics or anticholinergics.




**FORMULA:** Each Roter tablet contains: Bismuth Subnitrate Roter (350 mg.), Magnesium Carbonate (400 mg.), Sodium Bicarbonate (200 mg.) and Cortex Rhamni Frangulae (25 mg.).

**PACKINGS OF ROTER TABLETS:** Tins of 40 and 120, also dispensing sizes, 360 and 720 (P.T. exempt).

## ROTER IN PEPTIC ULCER

## A remarkably simple treatment in pancreatic enzyme replacement



Hitherto there has not been any reliable method of determining lipase activity in pancreatic extracts — none that ensured adequate control of fat digestion. This is now possible with Cotazym.

Cotazym (concentrated hog pancreas extract) enables known amounts of lipase to be administered easily and control to be established rapidly and accurately.

Cotazym is effective at lower dosage than has previously been possible and has, moreover, greater therapeutic activity. It has no offensive odour or taste and this, together with the simple method of its administration, makes it extremely acceptable to patients.

### INDICATIONS

**Fibrocystic disease**  
**Pancreatic deficiency states**  
**Post-gastrectomy malabsorption**  
**Steatorrhoea**

# COTAZYM




Each Cotazym capsule contains:

Lipase — sufficient to digest 17G. dietary fat  
Trypsin — sufficient to digest 34G. dietary protein  
Amylase — sufficient to digest 40G. dietary starch

### **Bottles of 100 capsules**

Dosage — 1-3 capsules with each meal or as directed by the physician.




Organon Laboratories Ltd., Crown House, London Road, Morden, Surrey

# MERSUTURES\*

## Eyeless Needled Suture

Designed to  
give greater  
needle control  
in suturing



### FOR THE SURGEON

A new sharp needle swaged to each suture, in packs specially designed to open more easily, saving time and space. There is an ETHICON\* needle suture combination for virtually every operative procedure . . . and they are ready for immediate use.


The majority of the MERSUTURES Eyeless Needled Suture range are available in foil overwrap presentation, as well as in the traditional Sterile Foil pack in jars.

# ETHICON, LTD.

Bankhead Avenue Sighthill Edinburgh 11

\* TRADEMARKS

5690ET



**2 TABLETS TWICE A DAY—  
PROVEN MAINTENANCE THERAPY**

**In Ulcerative Colitis**

---

available as a 0.5 g. tablet, either plain or enteric coated

**FIRST DEMONSTRATION . . . RESULTS SIGNIFICANT**

*"This is the first demonstration in a formal trial that any treatment reduces the relapse-rate in ulcerative colitis, 24 out of 34 patients taking 2 g. of sulphasalazine daily remained free of symptoms for a year. This result is significantly different from that obtained in a comparable group of patients who received a placebo."*

**PREFERABLE TO SYSTEMIC CORTICOSTEROIDS**

*"We have shown that sulphasalazine, 0.5 g. four times daily, is often effective in maintaining remission . . . and it therefore appears preferable to systemic corticosteroid drugs for this purpose."*

**REMISSION FOR ONE YEAR**

*"24 patients remained in symptomatic remission for a year while taking 2 g. of sulphasalazine daily, whereas only 8 remained symptom-free in the placebo group."*

**NORMAL MUCOSA**

*"2 of the 24 patients who remained symptom-free for a year while on sulphasalazine had a haemorrhagic mucosa at the end of the trial. The remaining 22 patients had a non-haemorrhagic mucosa which in many cases appeared normal."*

**MINIMAL SIDE EFFECTS**

*"In the present trial, using a dose of 2 g. daily, only 3 patients out of 34 had to discontinue treatment because of side-effects."*

**WHITE CELL COUNT**

*"In the patients treated with sulphasalazine there was no difference in the haemoglobin level before and after treatment; but the mean white-cell count was lower after six months or a year than at the start of treatment, though in no patient was it less than 4500 per c.mm."*

The Lancet Jan 23rd 1965 • Vol I • Pages 185-188

# SALAZOPYRIN




**PHARMACIA**  
(GREAT BRITAIN) LTD

Literature and further information are supplied on request  
**SINCLAIR HOUSE • THE AVENUE • WEST EALING • LONDON W13**  
Distributed by Martindale Samoores Ltd

## No nourishment here . . .

. . . unless an adequate supply of pancreatic enzymes is available to complete digestion.




Pancrex V is the most effective preparation for replacing pancreatic secretion in the gut. It is indicated in cystic fibrosis of the pancreas, chronic pancreatitis, pancreatectomy, post-cholecystectomy and in all digestive disorders where a relative deficiency of pancreatic enzymes is a contributory factor. Pancrex V is activated, whole dried pancreas

having a tryptic activity approximately equal to 5 times that of Pancreatin B.P.

# Pancrex V

References: *Diseases of Children* (1964), Blackwell, Oxford. *Diseases of Infancy and Childhood*. 8th Edn. (1962), Churchill, London. *Lancet* (1960), 1, 365. *Brit. Med. J.* (1958), 2, 1039





# **PRODEXIN<sup>reg.</sup>** **keeps acid level at placid level**

Prodexin puts acidity firmly in its place—right in the middle of the safe zone of pH 3-5. And keeps it there. Prodexin tablets give continuous and prolonged relief from pain caused by hyperacidity, dyspepsia and peptic ulcer.

Prodexin contains aluminium glycinate, a fast acting buffer antacid. Left to disperse slowly in the mouth, the tablet releases a steady trickle of antacid into the stomach: it simulates an intragastric drip in a simple and most convenient way.

Prodexin tablets are conveniently small. They can be carried about and taken wherever and whenever necessary. This is particularly useful to the ambulant patient.

**Dosage:** One or more tablets an hour when necessary. For continuous therapeutic effect, the tablet should be sucked slowly or left to disperse in the mouth. For rapid relief of pain, the tablet may be chewed and then swallowed.

**Content:** aluminium glycinate 900mg., light magnesium carbonate 100mg.



**Bencard**  
Brentford · England

# PEPTIC ULCER PROBLEMS? CAVED - (S) TABLETS

**A  
V  
E  
D  
-**

Munch med. Wochr. No. 23, 1952

Munch med. Wochr. No. 23, 1952  
Swiss Ther Rev. XX.7. (a963)  
Press Med., No. No. 35, 1963

**(S)**

Med. Welt., No 18, 1963

Patients with duodenal and gastric ulcers have been successfully treated with CAVED - (S) for over 15 years. CAVED - (S) contains a standardized liquorice by the exclusive CEDONA process by which the side effects of liquorice are eliminated. Oedema, and headaches have never been reported.

In vivo tests show good acid control with very small antacid content.

Formula: Cedona Block Liquorice 380 mgm., Bism. subnit. 100 mgm., Alum Hydrox. 100 mgm.; Magnes Carb. Levis 200 mgm.; Sod. Bicarb. 100 mgm.; Frangula 30 mgm.

Dosage. 2 tabs. t.d.s.: p.c. chewed and swallowed with water.


Cost Basic N.H.S. cost 5/5 per week.





**the antistaphylococcal  
antibiotic that penetrates tissues  
at these levels...**

**bears the FUCIDIN label**



The clinical significance of these levels has been confirmed in many publications during the past five years. Over 100 authors have endorsed the value of

Fucidin in the treatment of staphylococcal infection. Fucidin (Sodium Fusidate B.P.) 250 mgm capsules.  
Leo Laboratories Ltd Hayes Middx



# VELACTIN

**WANDER**

A readily assimilated milk substitute  
of vegetable origin for Milk Allergies



M441

Velactin contains no gluten or lactose. The biological value of the soya protein has been improved by the addition of DL-Methionine to make it equivalent to that of first-class animal protein.


*Literature gladly sent on request*

A. Wander Limited,  
42 Upper Grosvenor Street,  
London W.1.

**Indications:** Allergy or intolerance to human or cow's milk.<sup>1</sup> As basic feeding pending identification of specific food allergens.<sup>1</sup> Galactosaemia.<sup>2,3</sup> Gluten enteropathy. Velactin may also be tried in cases of undetermined metabolic or non-specific intolerance.

**References:** (1) *Nutritio et Dieta* 1961; **3**, 89.  
(2) *Lancet* 1959, **2** 849; 1960 **1**, 336.  
(3) *Ugeskrift f. Læger* 1960, **122**, 1229.

**Presentation:** Tin containing 1 lb. net.



## **Sustained relief of peptic ulcer all through the night**

Sustained relief of peptic ulcer is now possible by adjunctive treatment with Portyn\* Templets\*, *the new and highly potent sustained-release anticholinergic*. Each tablet contains 5 mg. benzilonium bromide, Parke-

Davis, for immediate release and 30 mg for the gradual release which provides all-night inhibition of gastric activity. Available in bottles of 15 and 100.

\*Trade mark

# **PORTYN TEMPLETS**

**PARKE-DAVIS**

HOUNSLOW · LONDON · TELEPHONE: HOUNSLOW 2361

O-1269

# BIOGASTRONE®

'It is the only drug therapy for gastric ulcer which has been demonstrated conclusively to accelerate healing, which can be achieved with patients remaining up and about, often at work, and with minimum change in diet'. *Gut*, 1965, 6:19

'Carbenoxolone has a specific effect, facilitating the healing of gastric ulcer, and is not merely yet another symptomatic measure'. *Brit med J.* 1965 2:1274


**heals gastric  
ulcers in  
ambulant  
patients**

Tablets of 50mg carbenoxolone sodium.  
Please write for full prescribing information



Berk Pharmaceuticals Limited Godalming Surrey England

® Regd Trade Mark



Whether you call it 'spastic colon', 'mucous colitis', or irritable colon' the clinical picture is the same—a chronic colonic disturbance with spasm/pain, disordered bowel habit (constipation or diarrhoea) and sometimes the passage of mucus per rectum.

Merbentyl (dicyclomine hydrochloride tablets B.N.F.) can relieve spasm/pain and help restore normal bowel activity in a very short space of time.

# Merbentyl

tablets of dicyclomine BNF

**Usual adult dose**  
2 tablets, three times daily

**Low basic N.H.S. Cost**  
An average week's treatment—2s. 8d.

Also available with phenobarbitone and as Merbentyl Syrup.

Full clinical information available on request.

Merrell-National (Laboratories) Ltd. London W.1

Trademarks:  
Merbentyl, Merrell-National



## ANTIBIOTICS AND THE INTESTINAL FLORA

Treatment with broad spectrum antibiotics suppresses the normal flora, which may be replaced by abnormal resistant organisms causing diarrhoea.

This change can be prevented or reversed by the implantation of *Lactobacillus acidophilus* which has been rendered antibiotic-resistant, and is thus able to multiply freely under these abnormal conditions. The products of growth of this organism are inimical to that of many other bacteria<sup>1,2</sup> and thus contribute to its rapid preponderance over other species.

Oral treatment with antibiotic-resistant *L. acidophilus* in the form of ENPAC has been shown to eliminate *Candida* from the faeces of infants with antibiotic-induced diarrhoea, with relief of symptoms succeeded by rapid gain in weight<sup>3,4</sup> and greatly to reduce the numbers of staphylococci in the faeces of patients treated with tetracyclines.<sup>5</sup>

<sup>1</sup> (1958) *Ann. Inst. Pasteur* 95, 194.      <sup>2</sup> (1959) *J. Bact.* 78, 477.      <sup>3</sup> (1957) *Klin. Wschr.* 35, 198.  
<sup>4</sup> (1959) *Medizinische* 7, 296.      <sup>5</sup> (1957) *Lancet* (i), 899.



# ENPAC

## LACTOBACILLUS ACIDOPHILUS PREPARATION

Samples and literature will be sent to the medical profession on request to:—

**APLIN and BARRETT LIMITED · BIOLOGICALS DIVISION · TROWBRIDGE · WILTS**  
**TEL. TRO. 4035**



A NEW JOURNAL

## British Journal of Medical Education

*The Journal of the Association for the Study of Medical Education*

The aim of the *British Journal of Medical Education* is to act as a medium for interchange of information on medical education—Undergraduate, Postgraduate, and Continuing—in the United Kingdom and Overseas. It hopes to encourage the objective study of medical education both in the United Kingdom and throughout the world. The journal gives preference to the reporting of fact rather than opinion. The first issue was published in December, 1966. Subsequent issues will be published quarterly in March, June, September and December.

*Annual Subscription, Inland £3 3s.; Abroad £3 10s. (U.S.A. \$10.00)*

***The first issue of this new journal was published in December, 1966***

### ORDER YOUR SUBSCRIPTION NOW

The Subscription Manager, BRITISH JOURNAL OF MEDICAL EDUCATION, B.M.A. House, Tavistock Square, London, W.C.1, or through any leading subscription agent or bookseller.



## LONGER-LASTING DRIP THERAPY

**Gastrils**—first ever antacid pastilles. Gastrils do not break up in the mouth like tablets; they dissolve slowly and smoothly, so give sustained drip therapy for the treatment of peptic ulcer.

Gastrils contain a co-dried gel of aluminium hydroxide and magnesium carbonate, a buffer antacid comparable in activity to freshly prepared liquid aluminium hydroxide gel. Gastrils rapidly buffer gastric pH to an optimum level, and keep it there, without risk of acid rebound.

To relieve the tedium of constant antacid therapy, every Gastrils prescription is dispensed in equal quantities of two delicious flavours—mint and fruit. Few antacids are as effective as Gastrils—none is as good to take!

# GASTRILS

## ANTACID PASTILLES

**INDICATIONS** Hyperacidity, peptic ulcer, gastritis, heartburn, oesophagitis and hiatus hernia.

**FORMULA** Each pastille contains 500 mg. co-dried gel of aluminium hydroxide and magnesium carbonate.

**BASIC N.H.S.** 24 individually wrapped Gastrils in carton 3/-.

**COST** 24 individually wrapped Gastrils from dispensing pack: 2/6½d. Gastrils are prescribable on form E.C.10

*'Gastrils' is a registered trade mark of T. J. Smith & Nephew Ltd.*



**Smith & Nephew  
Pharmaceuticals Limited**

WELWYN GARDEN CITY · HERTFORDSHIRE · ENGLAND

**The first issue of**

# **CARDIOVASCULAR RESEARCH**

**was published January, 1967**

**This new journal is published in association with the British Cardiac Society.**

*Cardiovascular Research* is mainly for the publication of basic research. The range of subjects covered by the journal includes physiological, pathological, pharmacological, biochemical, haemodynamic, surgical, and similar advances in the study of the heart and circulation.

The January issue has now been published and subsequent issues will be published quarterly in April, July and October. Each issue will contain approximately 80 pages. The title page, contents list and volume index will be contained in the October issue.

*Annual Subscription, Inland £3 3s.; Abroad £3 10s. (U.S.A. \$10.00)*

## **ORDER YOUR SUBSCRIPTION NOW**

Subscriptions can be ordered direct from the Subscription Manager, *Cardiovascular Research*, B.M.A. House, Tavistock Square, London W.C.1, or through any leading subscription agent or bookseller throughout the world.

## SCANDINAVIAN JOURNAL OF GASTROENTEROLOGY

Four numbers will make up one annual volume,  
costing N. Kr. 88.00, \$14.75, or £4.8s., post free

Volume 1, No. 2, 1966

### Contents

M. Siurala, H. Julkunen & B. A. Lamberg: <i>Gastrointestinal Tract in Hyperthyroidism before and after Treatment</i> .. .. .	79
P. M. Christiansen & A. Johansen: <i>Single Gastric Biopsy in Subjects with Low Acid Secretion after Maximal Histamine Stimulation</i> .. ..	86
R. Gamklou & Y. Edlund: <i>Ductal Factors in the Pathogenesis of Acute Pancreatitis in the Rat</i> .. .. .	94
E. Gjone, J. Myren & O. M. Orning: <i>Crohn's Disease in Norway—Clinical Features</i> .. .. .	101
S. Bengmark, B. Ekman, R. Olsson & B. Rehnström: <i>Further Studies on the Effect of Testosterone on Liver Healing after Partial Hepatectomy in Male Rats</i> .. .. .	106
H. Worning, E. Amdrup & F. W. Hendriksen: <i>Experimental Studies on the Intestinal Flow Using <sup>51</sup>Cr as Indicator</i> .. .. .	111
B. Fretheim, H. Rustad & K. Sandnes: <i>Pre-scalene Lymph Node Biopsy in the Evaluation of Gastric Carcinoma</i> .. .. .	121
M. Mignon, J. M. Limbosch, J. H. Wyllie, M. J. Rheault & L. M. Nyhus: <i>Effects of Calcium Infusion on Gastric Secretion in Heidenhain Pouch Dogs</i> .. .. .	124
J. Myren, S. Gjeruldsen & B. Fretheim: <i>Gastric Secretion before and after Graded Partial Gastrectomy for Duodenal Ulcer</i> .. ..	132
P. Bach-Nielsen, H. Baden & A. M. Christensen: <i>Controlled Radiological Evaluation of an Intestinal Pacemaker</i> .. .. .	138
P. G. Burhol & J. Myren: <i>Dehydrogenase Activity and Mucosal Measurements in Jejunal Biopsies from Patients with Gastrointestinal Diseases</i> .. ..	142
Ilkka Häkkinen & Heikki A. Salmi: <i>Gastric Juice Sulphomucoprotein and Vitamin B<sub>12</sub> Binding</i> .. .. .	149
O. Hänninen, K. Alanen & K. Hartiala: <i>Levels of Uridine Diphosphate Glucose Dehydrogenase and UDPG in the Gastrointestinal Mucous Membrane</i> .. .. .	152

Issued 14 November 1966


UNIVERSITETSFÖRLAGET:

Publishers to the Norwegian Universities.  
P.O. Box 307, Oslo 3, Norway.

P.O. Box 142, Boston, Mass. 02113, U.S.A.

Postgiro 155 28, Oslo.

Postcheck 400 181, Hamburg.



for prompt safe  
symptomatic  
medication in  
hyperchlorhydria  
and peptic ulcer

**'ALUHYDE'**

Tablets—present a combination with these attributes...

**'ALUHYDE'**

Effectively safe, reliable, non-systemic neutralization by means of two of the most preferred antacid adsorbents...

$\pm \pm \pm$

**'ALUHYDE'**

Presents a quick mental sedative effect by its inclusion of an alkali paraffinate—recognized for minimal sedative potential...

~~~~~

**'ALUHYDE'**

Provides antispasmodic action through its content of an ingredient long known for its favourable influence against hypermotility...

=====

**Formula—**

Each tablet: Dried Aluminium

Hydroxide Gel, B.P.C. — 3.75 gr.

Magnesium trisilicate B.P. — 3.75 gr.

Quinalbarbitone sodium B.P.—0.50 gr.

Tinct. Belladonnae B.P. — 3 m.

Packings—Regular packs of 50;

Dispensing packs of 500.

Samples available on personally signed request  
of qualified physicians only

**Multipax Chemicals Ltd**

121 NORWOOD ROAD HERNE HILL LONDON S.E.24. Phone: TULse Hill 3276

