

ROTER

IN PEPTIC DISORDERS

does more than just relieve pain




Each tablet contains: Bismuth
Subnitrate Roter (350 mg.),
Magnesium Carbonate (400 mg.),
Sodium Bicarbonate (200 mg.) and
Cortex Rhamni Frangulae (25 mg.).

Packings: Tins of 40 and 120, also
dispensing sizes, 360 and 720
(P.T. exempt).

ROTER tablets quickly overcome the pain and discomfort of minor gastric disturbances, but their true value is most evident in their control and frequent healing of peptic ulcer in both the early and the advanced or chronic stages. With Roter therapy, hospitalisation or surgery is mostly avoided; the patient remains ambulant and uncommitted to a restricted or prescribed diet; and recurring medical attention is usually obviated. Roter is worth remembering for your "gastric" patients.

Literature will be supplied on request from:—

F.A.I.R. LABORATORIES LTD., TWICKENHAM, MIDDLESEX.



When acid turns peptic ulcer into peptic agony NACTON^{*} stems it at source

Halves Gastric Acid Secretion and Controls Pain. Recently published results of trials over periods of up to 7 years have shown that Nacton (poldine tablets B.P.) is remarkably effective in the treatment of patients suffering from peptic ulcer. It is of considerable help whenever hyper-acidity is causing pain.

Specific · Certain · Continuous. Nacton consistently reduces acid secretion by half, alleviates ulcer pain and gives the ulcer a chance to heal. It is easy to take, causes no side-effects in therapeutic doses, does not lead to tolerance.

The patient may eat as he chooses, will rely less and less on antacids—and is released, through the continuing action of Nacton, from the fear of being woken by pain. Above all, he will be restored to a full working life: *in one stringent trial, absenteeism fell from an average of 28 days a year to 2.*

Full details on product information, dosage instructions and references will be sent promptly on request.



Bencard


Brentford, England. Isleworth 4111

^{*}regd.

MERSUTURES*

Eyeless Needled Suture

Designed to
give greater
needle control
in suturing



FOR THE SURGEON

A new sharp needle swaged to each suture, in packs specially designed to open more easily, saving time and space. There is an ETHICON* needle suture combination for virtually every operative procedure . . . and they are ready for immediate use.

The majority of the MERSUTURES Eyeless Needled Suture range are available in foil overwrap presentation, as well as in the traditional Sterile Foil pack in jars.

ETHICON, LTD.

Bankhead Avenue Sighthill Edinburgh 11

* TRADEMARKS

5690ET

NOW YOU CAN INVESTIGATE AND PHOTOGRAPH THE STOMACH WITHOUT DISTORTION

The Gastro Camera is a miniature instrument built into a flexible tube. It enables you to photograph all the main anatomical features of the stomach, gives you permanent records in colour and speeds diagnosis. The Gastro Camera is simplicity itself to use—no photographic knowledge is required.


FOUR models are available, two of them incorporating fibre-optics enabling you to investigate the stomach lining visually before taking photographs. FOUR other fibre-optic models are also available with *external* photographic facilities. They are for taking biopsies; for cytological examination; a gastro fiberscope with angle device; oesophageal

forward viewing fiberscope. All models are supplied complete with all necessary accessories. Projector and recording facilities are also available.

No service problems. All models are serviced in U.K. quickly, and by fully trained Air-Shields personnel. The Gastro Camera is one of the most important diagnostic developments in the field of gastro-enterology. You should find out about the range of models available. Phone us now for a demonstration or detailed technical literature.

**These photographs are reproduced in black and white from colour transparencies. Transparencies are available on request.*

SEE WHAT WE MEAN?





THE OLYMPUS OPTICAL RANGE OF GASTRO CAMERAS

Sole agents in the United Kingdom:

AIR SHIELDS (U.K.) LIMITED

Towerfield Road, Shoeburyness, Southend-on-Sea, Essex. Shoeburyness 2294/5.

Nausea and vomiting are effectively controlled by Maxolon



- ☐ Maxolon rapidly controls vomiting, whether due to local or blood-borne irritants.
- ☐ Maxolon may be given orally or by injection, according to the condition of the patient.
- ☐ Maxolon does not sedate at the recommended dose.
- ☐ Maxolon neither affects gastric secretions nor causes dryness of the mouth.

Indications:

Nausea and vomiting due to:

Gastro-intestinal disorders.

Intolerance to essential drugs.

Malignant disease.

Uraemic conditions.

Post-operative conditions.

Deep X-ray or Cobalt therapy.

N.B. Maxolon is not so effective in vomiting of labyrinthine origin such as travel sickness, vertigo and Ménière's disease.

Contra-indications: Although animal tests in several mammalian species have shown no teratogenic effects, Maxolon, like all new drugs,

is not recommended during the early weeks of pregnancy.

Precaution: As both Maxolon and the phenothiazines may occasionally cause benign transient dystonia, care should be exercised in the event of both drugs being prescribed concurrently.

Dosage:

Adult:

Oral: 1 tablet (10mg.) or two teaspoonfuls (10mg.) of syrup, three times daily.

I.M.: 1 ampoule (10mg.) 1-3 times daily depending upon the severity of the condition.

I.V.: 1 ampoule (10mg.) when required.

Children:

5-14 years, $\frac{1}{2}$ -1 teaspoonful of syrup (2.5-5mg.), three times daily.

Information on other dosages is available on request from the Company's Medical Department.

Availability:



Tablets: containing 10mg. metoclopramide, in canisters of 20.



Syrup: containing 5mg. metoclopramide in 5ml., in bottles of 100ml.



Ampoules for injection: 2ml. ampoules containing 10mg. metoclopramide, in boxes of 10.




Maxolon* (metoclopramide) is a product of

Beecham Research Laboratories Brentford, England.

*Regd.





Sustained relief of peptic ulcer all through the night

Sustained relief of peptic ulcer is now possible by adjunctive treatment with Portyn* Templets*, *the new and highly potent sustained-release anticholinergic*. Each tablet contains 5 mg. benzilonium bromide, Parke-

Davis, for immediate release and 30 mg for the gradual release which provides all-night inhibition of gastric activity. Available in bottles of 15 and 100.

*Trade mark

PORTYN TEMPLETS

PARKE-DAVIS HOUNSLOW · LONDON · TELEPHONE 01-570 2361

O-1269



SPRINGER-VERLAG
BERLIN · HEIDELBERG · NEW YORK

Experimentelle Medizin, Pathologie und Klinik

Herausgegeben von R. Hegglin, F. Leuthardt, R. Schoen,
H. Schwiegk, A. Studer, H. U. Zollinger

Band 19

Kähler: Das Karzinoid

**Klinik, Endokrinologie, pathologische Anatomie, Pathogenese
und Therapie**

Von Dr.
Hans Joachim Kähler
Wiesbaden

Mit 32 Abbildungen
XII, 281 Seiten Gr.-8°
1967
Gebunden DM 78,—
US \$ 19.50

■ Bitte Gesamt-
prospekt der Reihe
anfordern!

Aus dem Geleitwort von Professor Heilmeyer: Der Autor legt in der vorliegenden Monographie eine umfassende Darstellung der Klinik, der pathologischen Physiologie, der Pathologie und Therapie der Karzinoide mit und ohne endokrine Semiotik vor. Darin werden die Karzinoide des Dünndarms, des Magens, der Bronchien und des Rektums eingehend besprochen. Einen größeren Raum nimmt das typische Karzinoid-Syndrom ein. In einem Anhangskapitel wird die Physiologie, Pathophysiologie und Pharmakodynamik des Serotonins besprochen. Der Autor hat sich viele Jahre mit mir gemeinsam mit dem Karzinoidproblem an meiner Freiburger Klinik befaßt. Seine besondere Begabung, ein umfassendes Problem unter Berücksichtigung der gesamten Literatur souverän und klar zur Darstellung zu bringen, kommt auch der vorliegenden Arbeit zugute. Jeder Arzt und Forscher, der sich mit dem Karzinoid-Problem befaßt, wird darin über alle Fragen auf das Beste beraten.


**Recent Results in
Cancer Research**

**Fortschritte der
Krebsforschung**

**Progrès dans les
recherches
sur le cancer**

**Volume 10: Nelson, Radioactive Phosphorus in the Diagnosis
of Gastrointestinal Cancer**

By Robert S. Nelson, M.D., Professor of Medicine, General
Faculty, University of Texas; Internist, Chief,
Gastroenterology Service, Department of Medicine, The University
of Texas M.D. Anderson Hospital and Tumor Institute,
Houston, Texas
With 61 figures. Approx. 60 pages 8vo. 1967. Cloth DM 15,—
US \$ 3.75



**2 TABLETS TWICE A DAY—
PROVEN MAINTENANCE THERAPY**

In Ulcerative Colitis

available as a 0.5 g. tablet, either plain or enteric coated

FIRST DEMONSTRATION . . . RESULTS SIGNIFICANT

"This is the first demonstration in a formal trial that any treatment reduces the relapse-rate in ulcerative colitis, 24 out of 34 patients taking 2 g. of sulphasalazine daily remained free of symptoms for a year. This result is significantly different from that obtained in a comparable group of patients who received a placebo."

PREFERABLE TO SYSTEMIC CORTICOSTEROIDS

"We have shown that sulphasalazine, 0.5 g. four times daily, is often effective in maintaining remission . . . and it therefore appears preferable to systemic corticosteroid drugs for this purpose."

REMISSION FOR ONE YEAR

"24 patients remained in symptomatic remission for a year while taking 2 g. of sulphasalazine daily, whereas only 8 remained symptom-free in the placebo group."

NORMAL MUCOSA

"2 of the 24 patients who remained symptom-free for a year while on sulphasalazine had a haemorrhagic mucosa at the end of the trial. The remaining 22 patients had a non-haemorrhagic mucosa which in many cases appeared normal."

MINIMAL SIDE EFFECTS

"In the present trial, using a dose of 2 g. daily, only 3 patients out of 34 had to discontinue treatment because of side-effects."

WHITE CELL COUNT

"In the patients treated with sulphasalazine there was no difference in the haemoglobin level before and after treatment; but the mean white-cell count was lower after six months or a year than at the start of treatment, though in no patient was it less than 4500 per c.mm."

The Lancet Jan 23rd 1965 • Vol I • Pages 185-188

SALAZOPYRIN




PHARMACIA
(GREAT BRITAIN) LTD


Literature and further information are supplied on request
SINCLAIR HOUSE • THE AVENUE • WEST EALING • LONDON W13
Distributed by Martindale Samoore Ltd



Strength in hand



with London Hospital Sutures



**Strong, sure, supple — packaged for speed and safety.
Available in a full range of catgut and non-absorbable
materials with or without eyeless needles**



**London Hospital (Ligature Department) Ltd • Faringdon Avenue • Harold Hill • Romford • Essex
Telephone Ingrebourne 41101-4 • Telegrams "Ultratan Romford"**

PENBRITIN KS

3 way action stops Bacterial Diarrhoea



Penbritin KS contains Penbritin, Kaolin and Sulphadimidine, and has been specifically formulated to provide both symptomatic and bacterial treatment for gastro-intestinal infections.


PENBRITIN KS

eradicates infection in the wall of the intestine
kills organisms in the lumen

consolidates stools, adsorbs toxins and exerts a demulcent effect on inflamed mucosa.

Indications

Bacterial Diarrhoea
Bacterial Food Poisoning
Bacillary Dysentery
Intestinal infections due to sensitive organisms.



Wide Bacterial Cover

Penbritin KS is active against :
Salmonellae, Shigellae
E.coli, Strep. faecalis
(enterococci), Clostridia,
Staphylococci (other than
penicillinase-producing
strains)

Accepted Antibacterial Agents

Penbritin and Sulphadimidine are well-established B.P. preparations. Both are well absorbed and provide antibacterial activity not only in the lumen but also in the gut wall, the most important site of infection.



Rapid Action


Clinical experience has shown that relief is given often within 24 hours.

Dosage : Adults 20 ml. 6-hourly;
Children 10 ml. 6-hourly.

Availability : Bottles of 60 ml.
(Basic N.H.S. cost 8/9 per bottle).

Side-Effects Unlikely

Penbritin KS combines ampicillin and sulphadimidine, two of the safest and most widely used antibacterials. Side effects are unlikely, due to the short duration of treatment. Contra-indications : Allergy to penicillin or intolerance to sulphonamides.



Penbritin KS (regd.) is a product of

Beecham Research Laboratories
Brentford, England.



BIOGASTRONE®

carbenoxolone sodium

Specific healing effect

'Carbenoxolone has a specific effect, facilitating healing of gastric ulcer, and is not merely yet another symptomatic measure'. *Brit med J*, 1965, 2: 1274

Patients remain up and about

'It is the only drug therapy for gastric ulcer which has been demonstrated conclusively to accelerate healing, which can be achieved with patients remaining up and about, often at work, with minimum change in diet'. *Gut*, 1965, 5: 19

**the only drug
conclusively shown to
heal gastric ulcers**

without bed rest or dietary restrictions

Tablets of 50 mg carbenoxolone sodium.
Please write for full prescribing information.





Berk Pharmaceuticals Limited Godalming Surrey England

® Regd trade mark

No nourishment here . . .

. . . unless an adequate supply of pancreatic enzymes is available to complete digestion.




Pancrex V is the most effective preparation for replacing pancreatic secretion in the gut. It is indicated in cystic fibrosis of the pancreas, chronic pancreatitis, pancreatectomy, post-cholecystectomy and in all digestive disorders where a relative deficiency of pancreatic enzymes

is a contributory factor. Pancrex V is activated, whole dried pancreas

having a tryptic activity approximately equal to 5 times that of Pancreatin B.P.

Pancrex V



References: *Diseases of Children* (1964), Blackwell, Oxford. *Diseases of Infancy and Childhood*. 8th Edn. (1962), Churchill, London. *Lancet* (1960), 1, 365. *Brit. Med. J.* (1958), 2, 1039

SCANDINAVIAN JOURNAL OF GASTROENTEROLOGY

Four numbers will make up one annual volume,
costing N. Kr. 88.00, \$14.75, or £4.8s., post free

Contents

H. Peterson, J. Myren & O. P. Foss: <i>Total Nitrogen Content of Human Duodenal Juice before and after Intravenous Injection of Secretin</i> ..	1
J. Myren, R. Ceballos & G. C. Luketic: <i>Colchicine Effects on the Mouse Liver</i>	7
N. Helsingen Jr. & K. Nygaard: <i>A Jejunal Pouch as Substitute Stomach after Total Gastrectomy</i>	18
H. Worning, S. Müllertz, E. Hess Thaysen & H. O. Bang: <i>pH and Concentration of Pancreatic Enzymes in Aspirates from the Human Duodenum during Digestion of a Standard Meal</i>	23
I. P. T. Häkkinen: <i>Differentiation of Antigenic Gastric Cancer Sulphopolysaccharides from Metaplastic 'Intestinal' type Sulphopolysaccharides</i>	39
K. J. Öbrink & M. Waller: <i>Potassium in Gastric Juice under Non-Steady-State Conditions</i>	44
T. Scherstén: <i>The Synthesis of Taurocholic and Glycocholic Acids in Human Liver Homogenates and Subcellular Fractions in Obstructive Jaundice</i>	49
P. R. F. Bell: <i>Gastric Arterio-venous Shunts</i>	59
B. S. Bedi, J-P. Govaerts, S. P. Master & I. E. Gillespie: <i>Inhibition of Gastric Acid Secretion by Intravenous Cholecystokinin Extract</i> ..	68
J. Kyle, A. O. Adesola, L. F. Tinckler & J. De Beaux: <i>Incidence of Diverticulitis</i>	77

Issued 1 June 1967

UNIVERSITETSFORLAGET
Oslo and Bergen

Publishers to the Norwegian Universities.
P.O. Box 307, Oslo 3, Norway.

P.O. Box 142, Boston, Mass. 02113, U.S.A.

Postgiro 155 28, Oslo.

Postcheck 400 181, Hamburg.



The patient with an ulcer-round-the-corner

Hyperacidic and ulcer-prone— one of the many dyspeptic patients who are always coming back to your surgery for an antacid. His is a long-term problem, which calls for a long-term antacid. Not only an antacid with prolonged action but one he can live with. Handy and unobtrusive, easy to take and with a texture and flavour he doesn't easily tire of. Gastrils, in fact.

Gastrils are slow-dissolving acacia gum base pastilles each containing 500 mg. co-dried gel of aluminium hydroxide and magnesium carbonate. Sucking produces an even flow of antacid and demulcent that continues regardless of stomach emptying.

Presentation: to avoid monotony, two flavours, mint and fruit. Individually wrapped: in cartons of 24 or from the dispensing pack (always equal quantities of each flavour).

Basic N.H.S. cost of 24 pastilles from the dispensing pack, 2/6½d.

'Gastrils' is a registered trade mark of T. J. Smith & Nephew Ltd.

PRESCRIBABLE ON FORM E.C.10.

GASTRILS

give prolonged relief from acid pain



**Smith & Nephew
Pharmaceuticals Ltd.**

WELWYN GARDEN CITY · HERTFORDSHIRE

VELACTIN

WANDER

A readily assimilated milk substitute
of vegetable origin for Milk Allergies



M441

Velactin contains no gluten or lactose. The biological value of the soya protein has been improved by the addition of DL-Methionine to make it equivalent to that of first-class animal protein.

Literature gladly sent on request

A. Wander Limited,
42 Upper Grosvenor Street,
London W.1.

Indications: Allergy or intolerance to human or cow's milk.¹ As basic feeding pending identification of specific food allergens.¹ Galactosaemia.^{2,3} Gluten enteropathy. Velactin may also be tried in cases of undetermined metabolic or non-specific intolerance.


References: (1) *Nutritio et Dieta* 1961; **3**, 89.
(2) *Lancet* 1959, **2** 849; 1960 **1**, 336.
(3) *Ugeskrift f. Laeger* 1960, **122**, 1229.

Presentation: Tin containing 1 lb. net.



**the antistaphylococcal
antibiotic that penetrates tissues
at these levels...**

bears the FUCIDIN label



The clinical significance of these levels has been confirmed in many publications during the past five years. Over 100 authors have endorsed the value of

Fucidin in the treatment of staphylococcal infection. Fucidin (Sodium Fusidate B.P.) 250 mgm capsules.
Leo Laboratories Ltd Hayes Middx



ANTIBIOTICS AND THE INTESTINAL FLORA

Treatment with broad spectrum antibiotics suppresses the normal flora, which may be replaced by abnormal resistant organisms causing diarrhoea.

This change can be prevented or reversed by the implantation of *Lactobacillus acidophilus* which has been rendered antibiotic-resistant, and is thus able to multiply freely under these abnormal conditions. The products of growth of this organism are inimical to that of many other bacteria^{1,2} and thus contribute to its rapid preponderance over other species.

Oral treatment with antibiotic-resistant *L. acidophilus* in the form of ENPAC has been shown to eliminate *Candida* from the faeces of infants with antibiotic-induced diarrhoea, with relief of symptoms succeeded by rapid gain in weight^{3,4} and greatly to reduce the numbers of staphylococci in the faeces of patients treated with tetracyclines.⁵

¹ (1958) *Ann. Inst. Pasteur* **95**, 194.

² (1959) *J. Bact.* **78**, 477.

³ (1957) *Klin. Wschr.* **35**, 198.

⁴ (1959) *Medizinische* **7**, 296.

⁵ (1957) *Lancet* (i), 899.



LACTOBACILLUS ACIDOPHILUS PREPARATION

Samples and literature will be sent to the medical profession on request to:—

APLIN and BARRETT LIMITED · BIOLOGICALS DIVISION · TROWBRIDGE · WILTS
TEL. TRO. 4035

Published 1 June, 1967

TISSUE AND ORGAN TRANSPLANTATION

The Proceedings of a Symposium organised by the College of Pathologists

CONTENTS: Genetic background and transplantation antigens · Action of lymphocytes in transplantation · The role of the macrophage in transplantation immunity · Humoral and cell-mediated effector mechanisms in tissue transplantation · Immunological tolerance · The response of draining lymph nodes to immunological stimulation in intact and thymectomized animals · Immunological properties of antilymphocytic serum · Immunosuppressive agents and allogeneic transplantation · Immunosuppressive action of ribonucleases · The relevance of leucocyte antigens for allogeneic renal transplantation · The transplantation of cartilage · Rejection in treated renal allografts · Grafts of haemopoietic tissue: the nature of haemopoietic stem cells · Bone transplantation.

PRICE 30s. (U.S.A. \$4.50) including postage

Copies can be ordered now from: The Subscription Manager

JOURNAL OF CLINICAL PATHOLOGY

B.M.A. House, Tavistock Square, London, W.C.1., or through any leading bookseller

EDIZIONI MINERVA MEDICA

Quarterly Index

**MINERVA
MEDICO-
BIBLIOGRAFICA**

A list of the world's medical,
surgical and specialist
literature.

Annual subscription rate \$ 25.00

Italian Medical Journal in
English

**PANMINERVA
MEDICA**

The Journal of the Italian Medical
Association Monthly Review of
Italian Medicine.

Annual subscription rate \$ 13.30

INTERNATIONAL MEDICAL JOURNALS IN ENGLISH

**THE JOURNAL OF
SPORTS MEDICINE
AND PHYSICAL
FITNESS**

Official Journal of the "Fédération
Internationale de Médecine
Sportive". Quarterly publication.

Annual subscription rate \$ 10.00


**THE JOURNAL OF
CARDIOVASCULAR
SURGERY**

Official Journal of the International
Cardiovascular Society.
Bi-monthly publication.

Annual subscription rate \$ 13.30

EDIZIONI MINERVA MEDICA

corso Bramante 83-85 - Torino - Italy



**for prompt safe
symptomatic
medication in
hyperchlorhydria
and peptic ulcer**

• ALUHYDE •

Tablets present a combination with these advantages...

• ALUHYDE •

Effects safe, reliable, nonsystemic neutralization by means of two of the most preferred antacid adsorbents...



• ALUHYDE •

Permits of quick mental sedative effect by its inclusion of a parbiturate recognized for minimal cumulative potential...



• ALUHYDE •

Provides antispasmodic action through its content of an agent long known for its favourable influence against hypermotility...



Formula—

Each tablet: Dried Aluminium
Hydroxide Gel, B.P.C. — 3.75 gr.
Magnesium trisilicate B.P. — 3.75 gr.
Quinalbarbitone sodium B.P.—0.50 gr.
Tinct. Belladonnae B.P. — 3.m.

**Packings—Regular packs of 50;
Dispensing packs of 500.**

**Samples available on personally signed request
of qualified physicians only**

Multipax Chemicals Ltd

121 NORWOOD ROAD HERNE HILL LONDON S.E.24. Phone: TULse Hill 3276

