

ROTER

IN PEPTIC DISORDERS

does more than just relieve pain



Each tablet contains: Bismuth
Subnitrate Roter (350 mg.),
Magnesium Carbonate (400 mg.),
Sodium Bicarbonate (200 mg.) and
Cortex Rhamni Frangulae (25 mg.).

Packings: Tins of 40 and 120, also
dispensing sizes, 360 and 720
(P.T. exempt).

ROTER tablets quickly overcome the pain and discomfort of minor gastric disturbances, but their true value is most evident in their control and frequent healing of peptic ulcer in both the early and the advanced or chronic stages. With Roter therapy, hospitalisation or surgery is mostly avoided; the patient remains ambulant and uncommitted to a restricted or prescribed diet; and recurring medical attention is usually obviated. Roter is worth remembering for your "gastric" patients.

Literature will be supplied on request from:-


F.A.I.R. LABORATORIES LTD., TWICKENHAM, MIDDLESEX.



new-

NE[®]

introduces direct
healing for
duodenal
ulcers



DUOGASTRONE[®]

introduces specific therapy for duodenal ulcers

DUOGASTRONE has a direct, rapid healing action which can obviate the need for symptomatic medication, bed rest and dietary restriction in the treatment of duodenal ulcers.

The active constituent, carbenoxolone sodium, has been described as the only drug to heal gastric ulcers.¹ Now, by means of a specially designed 'positioned release' capsule, DUOGASTRONE delivers this unique active healing agent directly to the site of duodenal ulceration. Absorbing moisture from the stomach, the DUOGASTRONE capsule swells to a size which prevents its passage through the pyloric canal. At the pyloric antrum – the pressure of the sphincter and the intracapsular pressure combine to break open the capsule – releasing a concentrated solution of carbenoxolone sodium directly into the duodenum.²

Patients feel the benefit of DUOGASTRONE therapy within a few days. Freed from distressing symptoms they go about their normal activities whilst healing proceeds. Objective improvement – as evidenced by disappearance of the ulcer crater – is often seen within four weeks.²

Including the published results of a double-blind trial² the over-all success rate with DUOGASTRONE was 87%.

DUOGASTRONE capsules, each containing 50 mg carbenoxolone sodium. Containers of 28 capsules.

References

1 Gut (1965) 6:19.

2 Practitioner (1967) 199:109.



Additional information on request from:



Berk Pharmaceuticals Limited, Godalming, Surrey

No nourishment here . . .

. . . unless an adequate supply of pancreatic enzymes is available to complete digestion.





Pancrex V is the most effective preparation for replacing pancreatic secretion in the gut. It is indicated in cystic fibrosis of the pancreas, chronic pancreatitis, pancreatectomy, post-cholecystectomy and in all digestive disorders where a relative deficiency of pancreatic enzymes is a contributory factor. Pancrex V is activated, whole dried pancreas

having a tryptic activity approximately equal to 5 times that of Pancreatin B.P.

Pancrex V

References: *Diseases of Children* (1964), Blackwell, Oxford. *Diseases of Infancy and Childhood*. 8th Edn. (1962), Churchill, London. *Lancet* (1960), 1, 365. *Brit. Med. J.* (1958), 2, 1039





Early warning campaign page vii

"Ulcerative colitis is one of the greatly misunderstood diseases of today. It is mistakenly thought to be rare only because it is not recognised often in its early stages. This is unfortunate, as many patients respond well to medical treatment if the disease is recognised in time. With proper management, most of them can avoid surgery."¹

"It is also felt that the incidence is increasing."²

As with any disease, early diagnosis and immediate initiation of definitive therapy is essential. Prompt action will reduce the number of fulminating cases and benefit both the patient and the physician. To secure earlier hospital referrals general practitioners are now being alerted by an extensive information campaign detailing the symptoms of the disease.

The acute attack page vi


"Ulcerative colitis is a formidable disease which carries a greatly increased risk of death throughout the whole period of follow up."³

The rational approach to therapy of ulcerative colitis is to use a drug with a specific affinity for connective tissue especially the colonic submucosa. Salazopyrin is such a drug. ⁴ Results of Salazopyrin treatment are often dramatically successful. ⁵

The relapse page viii

Details of "the first demonstration in a formal trial that any treatment reduces the relapse-rate in ulcerative colitis."⁶

1. Postgrad. Med., 1960, 28, 157
2. Proc. roy. Soc. Med., 1966, 59, 369
3. Gut, 1963, 4, 299
4. Acta Med. scand., 1963, 173, 61 and 391
5. From Gastroenterology Vol II. W. B. Saunders Company Philadelphia 1964, p. 863
6. Lancet, 1965, i, 185



The acute attack

Achieving Remission

Salazopyrin has a marked affinity for connective tissues. Very high concentrations of Salazopyrin are found in the intestines and intestinal lumen, and unlike ordinary sulphonamides, Salazopyrin exhibits a prolonged retention in the colonic submucosa. ¹

The results of Salazopyrin treatment of ulcerative colitis are often dramatic. Within two or three days, the number of stools decreases, abdominal cramps disappear, the fever subsides and the appetite improves. ²

The auxiliary rôle of steroids in fulminating cases must not be overlooked, but their action is essentially suppressive rather than curative. ³ Successful results are reported with the combined use of Salazopyrin and steroids. ^{4, 5}

However, the mild and moderate acute attacks are best treated with Salazopyrin alone. ^{6, 7}

A high percentage of patients respond to medical treatment but more than 80% will have a relapse within one year unless treatment is continued beyond the acute attack. ⁸ (see following pages)

Dosage for the Acute Attack

Two to four tablets (1 g to 2 g) four to six times daily. The dosage should be adjusted according to the patient's needs. This is decreased to the maintenance dose (2 g daily) as the patient improves. At any indication of a relapse, however, the dosage should be increased to the maximum tolerated level.

1. Acta Med. scand., 1963, 173, 61 and 391
2. From Gastroenterology Vol II. W. B. Saunders Company Philadelphia 1964, p.863
3. New Engl. J. Med., 1964, 271, 891
4. Brit. med. J., 1962, ii, 1708
5. Dallas Med. J., 1964, 50, 240
6. Proc. roy. Soc. Med., 1960, 53, 647
7. Gut, 1960, 1, 217
8. Gut, 1963, 4, 299

The patient with ulcerative colitis is usually pale, tense, depressed and exceedingly disturbed emotionally.

He is weak and beset by many problems (some real and some imaginary).


His intestines are inflamed and infected. He passes very frequent bloody and pus-ridden stools. One bowel action often precedes another. He may exhibit rectal tenesmus.

The condition will also manifest itself by loss of weight, anorexia, anaemia, abdominal cramps and fever.

Examination reveals malnutrition, dehydration, abdominal tenderness and increased bowel sounds. The rectal mucosa has a granular feel.

Mild and moderate cases may only have some of these symptoms.

**THIS NOTICE IS CURRENTLY
BEING BROUGHT TO THE ATTENTION
OF GENERAL PRACTITIONERS
WITH THE OBJECT OF CREATING
GREATER AWARENESS OF
ULCERATIVE COLITIS**





The relapse

Maintaining Remission

Most ulcerative colitis patients will respond to medical treatment but more than 80% will have a relapse within 1 year unless the treatment is continued beyond the acute attack.

Salazopyrin is the *only* preparation which is suitable for the long-term treatment of the ulcerative colitis patient.

"This is the first demonstration in a formal trial that any treatment reduces the relapse rate in ulcerative colitis . . . it therefore appears preferable to systemic corticosteroids, for this purpose.

24 (out of 34) patients remained in symptomatic remission for a year while taking 2 g of Salazopyrin daily whereas only 8 (out of 33) remained symptom free in the placebo group.

22 out of the 24 patients on Salazopyrin who remained in remission at the end of the trial had a non-haemorrhagic mucosa which, in many cases, appeared normal.

. . . only 3 patients out of 34 had to discontinue treatment because of side-effects.

In the patients treated with Salazopyrin, there was no difference in the haemoglobin level before and after treatment; but the mean white-cell count was lower after 6 months or a year than at the start of the treatment, though in no patient was it less than 4500 per c. mm."

Lancet, 1965, i. 185

SALAZOPYRIN[®]

Proven Maintenance Therapy: 2 tablets twice a day.
Salazopyrin (sulphasalazine) is available as both the plain 0.5g tablet or as the 0.5g En-tablet. The latter is enteric coated and has been specifically designed for use in the patient who may exhibit gastro-intestinal intolerance to the plain tablet.
Literature and detailed information on Salazopyrin are available on request.



PHARMACIA
(GREAT BRITAIN) LTD

Paramount House, 75 Uxbridge Road, Ealing, London W.5. Telephone 01-579 0102/7.

Distributed by Martindale Samoore Ltd., and in the Republic of Ireland by Goodbody Ltd.

VELACTIN

WANDER

A readily assimilated milk substitute
of vegetable origin for Milk Allergies



M441

Velactin contains no gluten or lactose. The biological value of the soya protein has been improved by the addition of DL-Methionine to make it equivalent to that of first-class animal protein.

Literature gladly sent on request

A. Wander Limited,
42 Upper Grosvenor Street,
London W.1.

Indications: Allergy or intolerance to human or cow's milk.¹ As basic feeding pending identification of specific food allergens.¹ Galactosaemia.^{2,3} Gluten enteropathy. Velactin may also be tried in cases of undetermined metabolic or non-specific intolerance.

References: (1) *Nutritio et Dieta* 1961; **3**, 89.
(2) *Lancet* 1959, **2** 849; 1960 **1**, 336.
(3) *Ugeskrift f. Laeger* 1960, **122**, 1229.

Presentation: Tin containing 1 lb. net.

NOW YOU CAN INVESTIGATE AND PHOTOGRAPH THE STOMACH WITHOUT DISTORTION

The Gastro Camera is a miniature instrument built into a flexible tube. It enables you to photograph all the main anatomical features of the stomach, gives you permanent records in colour and speeds diagnosis. The Gastro Camera is simplicity itself to use—no photographic knowledge is required.


FOUR models are available, two of them incorporating fibre-optics enabling you to investigate the stomach lining visually before taking photographs. FOUR other fibre-optic models are also available with *external* photographic facilities. They are for taking biopsies; for cytological examination; a gastro fiberscope with angle device; oesophageal

forward viewing fiberscope. All models are supplied complete with all necessary accessories. Projector and recording facilities are also available.

No service problems. All models are serviced in U.K. quickly, and by fully trained Air-Shields personnel. The Gastro Camera is one of the most important diagnostic developments in the field of gastro-enterology. You should find out about the range of models available. Phone us now for a demonstration or detailed technical literature.

**These photographs are reproduced in black and white from colour transparencies. Transparencies are available on request.*

SEE WHAT WE MEAN?



THE OLYMPUS OPTICAL RANGE OF GASTRO CAMERAS

Sole agents in the United Kingdom:


AIR SHIELDS (U.K.) LIMITED

Towerfield Road, Shoeburyness, Southend-on-Sea, Essex. Shoeburyness 2294/5.

MERSUTURES* **General Closure Range**

Eyeless
Needled
Sutures

Introducing
hand held needles



For the surgeon who prefers Hand Held Needles, Ethicon Ltd offer a complete range of needled sutures attached to catgut, Black Braided Silks and Nylon to meet most surgical interventions.

DESCRIPTIVE LITERATURE AVAILABLE ON REQUEST

ETHICON, LTD.

Bankhead Avenue, Sighthill, Edinburgh 11

ANTIBIOTICS AND THE INTESTINAL FLORA

Treatment with broad spectrum antibiotics suppresses the normal flora, which may be replaced by abnormal resistant organisms causing diarrhoea.

This change can be prevented or reversed by the implantation of *Lactobacillus acidophilus* which has been rendered antibiotic-resistant, and is thus able to multiply freely under these abnormal conditions. The products of growth of this organism are inimical to that of many other bacteria^{1,2} and thus contribute to its rapid preponderance over other species.

Oral treatment with antibiotic-resistant *L. acidophilus* in the form of ENPAC has been shown to eliminate *Candida* from the faeces of infants with antibiotic-induced diarrhoea, with relief of symptoms succeeded by rapid gain in weight^{3,4} and greatly to reduce the numbers of staphylococci in the faeces of patients treated with tetracyclines.⁵

¹ (1958) *Ann. Inst. Pasteur* **95**, 194. ² (1959) *J. Bact.* **78**, 477. ³ (1957) *Klin. Wschr.* **35**, 198.
⁴ (1959) *Medizinische* **7**, 296. ⁵ (1957) *Lancet* (i), 899.



LACTOBACILLUS ACIDOPHILUS PREPARATION

Samples and literature will be sent to the medical profession on request to:—

APLIN and BARRETT LIMITED · BIOLOGICALS DIVISION · TROWBRIDGE · WILTS
TEL. TRO. 4035



Pergamon Press Books



GASTRIC SECRETION— Mechanisms and Control

Proceedings of a Symposium

Held at the University of Alberta,
Edmonton, Canada.

Edited by **T. K. Shnikta,**
J. A. L. Gilbert and **R. C. Harrison,**
University of Alberta.

Partial Contents:

Structure and Function: The Fine Structure of the Gastric Mucosa—**S. Ito**; Observations on the Regulation of the Growth of the Gastric Mucosa—**G. P. Crean**; *Neuro-Endocrine Control*: The Nerve Supply of the Stomach and its Relationship to Secretion—**J. G. Murray**; Vagal Effects on the Gastric Secretion—**L. Olbe**; Magnitude of the Antrum's Role in the Cephalic Phase of Gastric Secretion—**P. H. Jordan Jnr.** and **C. De La Rosa**; *Mucus*: Regulation of Gastric Mucus Secretion—**R. Menguy**; Stimuli for Gastric Mucus Secretion—**F. Hollander**; Intrinsic Factor Secretion—**G. H. Jefferies**; *Secretion: Stimulation*: Gastric Secretion and Duodenal Ulcer—**L. R. Dragstedt**; The Role

of Intestinal Digestion in the Control of Acid Secretion; **D. W. Elliott** and **G. Kerr**; *Gastrin*: Some Physiological Actions of Gastrin—**M. I. Grossman**; Action of Purified Gastrin Polypeptide on the Isolated Gastric Mucosa of the Bullfrog—**W. D. Davidson, C. A. E. Lemmi** and **J. C. Thompson**; *Secretion: Inhibition*: The Recognition and Assay of Gastrone—**C. F. Code**; Endocrine and Exocrine Studies on Homotransplanted Gastric Pouches—**J. C. Thompson, I. A. Davies, W. D. Davidson** and **J. H. Miller**
516 pages £7/\$21.50*

*Sterling prices applicable in U.K. only.

Pergamon Press Ltd.
Headington Hill Hall, Oxford

Pergamon Press Inc.
44-01 21st Street,
Long Island City,
New York 11101, U.S.A.

PRECLINICAL CARCINOMA OF THE CERVIX UTERI

Its nature, origin and development.
An Interdisciplinary Approach

M. Coppleson, *King George V Memorial Hospital, Royal Prince Alfred Hospital, Sydney* and **B. Reid,** *Queen Elizabeth Research Institute, The University of Sydney.*

A lavishly illustrated record of the work for which four authors were awarded the Edgar Gentilli Prize by the Royal College of Obstetricians and Gynaecologists. The book deals with the origin, nature and management of preclinical squamous carcinoma of the cervix, including carcinoma *in situ* and early invasive carcinoma which is undetected by the traditional methods of speculum examination, palpitation, probing and history taking.

Section Headings:

Introduction; Clinical Studies; Epidemiological Studies; Laboratory Studies; General Cancer Theory.
320 pages £5/\$15.00*

CAVED - (S)

TABLETS FOR PEPTIC ULCERS



**CAVED-(S) THE LIQUORICE THERAPY,
NO SIDE EFFECTS EVER REPORTED i.e., OEDEMA,
HEADACHES, ETC.**

More than 10 written up Hospital Trials, including four
British double blind. 15 years use in 44 countries.

DUODENAL AND GASTRIC ULCERS SUCCESSFULLY TREATED

Good results obtained with all age groups and chronic cases.

**Offers safe ambulant treatment without diet,
restrictions or additional rest, etc.**

Very good in vivo acid control. Powerful spasmolytic action.

Very low cost. 7 days full treatment dose N.H.S. cost 5/5d.

DOSE: 2 tabs. t.d.s., p.c., chewed and swallowed with a drink of water


Formula: Cedona Block Liquorice, Bism. subnit., Alum Hydrox.,
Magnes. Carb. Levis., Sod. Bicarb. Frangula.

Requests for Representative to call, or Samples and Literature to:

TILLOTTS LABORATORIES, 94 Marsham Street, London S.W.1



Strength in hand



with London Hospital Sutures




**Strong, sure, supple – packaged for speed and safety.
Available in a full range of catgut and non-absorbable
materials with or without eyeless needles**



**London Hospital (Ligature Department) Ltd • Faringdon Avenue • Harold Hill • Romford • Essex
Telephone Ingrebourne 41101-4 • Telegrams "Ultratan Romford"**

Nausea and vomiting are effectively controlled by Maxolon



- Maxolon rapidly controls vomiting, whether due to local or blood-borne irritants.
- Maxolon may be given orally or by injection, according to the condition of the patient.
- Maxolon does not sedate at the recommended dose.
- Maxolon neither affects gastric secretions nor causes dryness of the mouth.

Indications:

Nausea and vomiting due to:

Gastro-intestinal disorders. _____
 Intolerance to essential drugs. _____
 Malignant disease. _____
 Uraemic conditions. _____
 Post-operative conditions. _____
 Deep X-ray or Cobalt therapy. _____

N.B. Maxolon is not so effective in vomiting of labyrinthine origin such as travel sickness, vertigo and Ménière's disease.

Contra-indications: Although animal tests in several mammalian species have shown no teratogenic effects, Maxolon, like all new drugs,

is not recommended during the early weeks of pregnancy.

Precaution: As both Maxolon and the phenothiazines may occasionally cause benign transient dystonia, care should be exercised in the event of both drugs being prescribed concurrently.

Dosage:

Adult:

Oral: 1 tablet (10mg.) or two teaspoonfuls (10mg.) of syrup, three times daily.

I.M.: 1 ampoule (10mg.) 1-3 times daily depending upon the severity of the condition.

I.V.: 1 ampoule (10mg.) when required.

Children:

5-14 years, $\frac{1}{2}$ -1 teaspoonful of syrup (2.5-5mg.), three times daily.

Information on other dosages is available on request from the Company's Medical Department.

Availability:

Tablets: containing 10mg. metoclopramide, in canisters of 20.

Syrup: containing 5mg. metoclopramide in 5ml., in bottles of 100ml.

Ampoules for injection: 2ml. ampoules containing 10mg. metoclopramide, in boxes of 10.




Maxolon* (metoclopramide) is a product of

Beecham Research Laboratories Brentford, England.

*Regd.



in staphylococcal infection of the gastro-intestinal tract



powerful · bactericidal · antistaphylococcal

The potential severity and the rapidity of staphylococcal enterocolitis demand prompt and specific antibacterial therapy—often well before sensitivity reports become available. FUCIDIN is rapidly and easily absorbed; its unique molecular structure and metabolism lead to biliary concentrations up to 12 times higher than serum levels and bactericidal distribution throughout the alimentary tract. FUCIDIN exhibits remarkable ability to reach the nidus of infection even in the presence of pus, and does not disturb normal bowel flora. For early control of infection FUCIDIN has proved a reliable and economical choice—in over five years of clinical use.

For full literature and prescribing information please write to the Medical Information Department.

fucidin

Capsules, 250 mg. sodium fusidate B.P., in bottles of 36 and 250; also as Paediatric Suspension equivalent to 35 mg. sodium fusidate per ml., in bottles of 60 ml.



LEO LABORATORIES LIMITED · HAYES
LEO MIDDLESEX · TELEPHONE 01-573 6224

BIOGASTRONE®

carbenoxolone sodium

Specific healing effect

'Carbenoxolone has a specific effect, facilitating healing of gastric ulcer, and is not merely yet another symptomatic measure'. *Brit med J*, 1965, 2: 1274

Patients remain up and about

'It is the only drug therapy for gastric ulcer which has been demonstrated conclusively to accelerate healing, which can be achieved with patients remaining up and about, often at work, with minimum change in diet'. *Gut*, 1965, 5: 19

**the only drug
conclusively shown to
heal gastric ulcers**

without bed rest or dietary restrictions

Tablets of 50 mg carbenoxolone sodium.
Please write for full prescribing information.



Berk Pharmaceuticals Limited Godalming Surrey England

© Regd trade mark

Gastroenterologia

International Journal of Gastroenterology

The journal publishes original papers and surveys in the field of gastroenterology by scientists of all countries. Articles are published in English, German and French at the choice of the author and they are followed by a summary in all three languages.

Two volumes of six numbers each are issued annually. Subscription price per volume 95s. or S.Frs. 56 (postage included).

Some recently published contributions in English:

E. D. PALMER (East Orange, N.J.): Course of Esophageal Varices in Cirrhosis, in the Absence of Portal Decompression—O. HARTKOPP and P. AAGAARD, (Copenhagen): Studies on Uropepsin Excretion, Maximal and Basal Acid Output in Recurrent Peptic Ulcer—E. VENTURA; and S. CALANDRA (Modena) M. L. ZENEROLI and S. CALANDRA (Modena): Serum

Ceruloplasmin in Liver Cirrhosis—G. F. GARUSI and L. NOTO (Verona): Inflammatory Pseudo-Tumours of the Caecum—J. TICHÝ and M. HRADSKÝ (Hradec Králové): Ultrastructure of the Gastric Mucosa in Healthy Human Subjects and in Patients with Pernicious Anaemia.

Please ask for specimen copies.

S. Karger


4000 Basel (Switzerland)

New York

Arnold-Böcklin-Strasse 25

**All orders from the U.K. to be placed with
J. Wiley and Sons, Baffins Lane, Chichester, Sussex**

The patient with an ulcer-round-the-corner



Hyperacidic and ulcer-prone—one of the many dyspeptic patients who are always coming back to your surgery for an antacid. His is a long-term problem, which calls for a long-term antacid. Not only an antacid with prolonged action but one he can live with. Handy and unobtrusive, easy to take and with a texture and flavour he doesn't easily tire of. Gastrils, in fact.

Gastrils are slow-dissolving acacia gum base pastilles each containing 500 mg. co-dried gel of aluminium hydroxide and magnesium carbonate. Sucking produces an even flow of antacid and demulcent that continues regardless of stomach emptying.

Presentation: to avoid monotony, two flavours, mint and fruit. Individually wrapped: in cartons of 24 or from the dispensing pack (always equal quantities of each flavour).

Basic N.H.S. cost of 24 pastilles from the dispensing pack, 2/6½d.

'Gastrils' is a registered trade mark of T. J. Smith & Nephew Ltd.

PRESCRIBABLE ON FORM E.C.10.

GASTRILS

give prolonged relief from acid pain

Full information is available on request.

SNP Smith & Nephew
Pharmaceuticals Ltd.

WELWYN GARDEN CITY · HERTFORDSHIRE

Published 1 June 1967

TISSUE AND ORGAN TRANSPLANTATION

The Proceedings of a Symposium organised by the College of Pathologists

Genetic background and transplantation antigens	J. R. BATCHELOR and B. A. CHAPMAN
Action of lymphocytes in transplantation	J. L. TURK
The role of the macrophage in transplantation immunity	D. C. DUMONDE
Humoral and cell-mediated effector mechanisms in tissue transplantation	GÖRAN MÖLLER and ERNA MÖLLER
Immunological tolerance	N. A. MITCHISON
The response of draining lymph nodes to immunological stimulation in intact and thymectomized animals	DELPHINE M. V. PARROTT
Immunological properties of antilymphocytic serum	M. F. A. WOODRUFF
Immunosuppressive agents and allogeneic transplantation	M. C. BERENBAUM
Immunosuppressive action of ribonucleases	J. F. MOWBRAY
The relevance of leucocyte antigens for allogeneic renal transplantation	J. J. VAN ROOD, A. VAN LEEUWEN and J. W. BRUNING
The transplantation of cartilage	THOMAS GIBSON
Rejection in treated renal allografts	K. A. PORTER
Grafts of haemopoietic tissue: the nature of haemopoietic stem cells	J. F. LOUTIT
Bone transplantation	J. CHALMERS

PRICE 30s. (U.S.A. \$4.50) including postage

Copies can be ordered now from: The Subscription Manager

JOURNAL OF CLINICAL PATHOLOGY

B.M.A. House, Tavistock Square, London, W.C.1. or through any leading bookseller

Deutsche Zeitschrift für Verdauungs- und Stoffwechselkrankheiten

einschließlich Theorie und Praxis der Krankenernährung

Organ der Deutschen Gesellschaft für Verdauungs- und Stoffwechselkrankheiten

Unter Mitarbeit von

H.-W. BANSI—Hamburg, H. H. BERG—Hamburg,
H. BREDT—Mainz, R. EMMRICH—Leipzig,
G. FANCONI—Zürich, N. HENNING—Erlangen,
E. KLENK—Köln, K. KLINKE—Düsseldorf,
R. MANCKE—Rendsburg, N. G. MARKOFF—Chur,
A. PEIPER—Leipzig, K. SEIDEL—Jena,
W. SIEDE—Frankfurt/Main, K. THOMAS—Göttingen,
J. WALDENSTRÖM—Malmö und
A. WALLGREN—Stockholm

Herausgegeben von

Professor Dr. M. GÜLZOW, Rostock, und
Professor Dr. D. LOHMANN, Leipzig

The "Deutsche Zeitschrift für Verdauungs- und Stoffwechselkrankheiten" (German Journal for Diseases of the Digestive and Metabolic System, including the Theory and Practice of Invalid Diet) aims to inform specialists, whether scientists or practitioners, of internal medicine, surgery, pediatry, radiography etc., about the latest discoveries. It continues the tradition of the oldest German-language journal of gastroenterology, from the archives of diseases of the digestive system. As well as original contributions, which form the major part of the contents, it contains current surveys, reports of conferences at home and abroad, book reviews and summaries of original contributions in this field published in other journals.

Yearly, 1 volume of 6 issues, each volume MDN 31, 80. The periodical may be obtained through the local book trade as well as directly from the publishing house.

J. A. BARTH

DDR 701 Leipzig, Salomonstr. 18 b, POB 109

JOHANN AMBROSIUS BARTH · LEIPZIG



• ALUHYDE •

Tablets present a combination with three active ingredients...

• ALUHYDE •

For relief of hyperchlorhydria and peptic ulceration... 

• ALUHYDE •

Points of experimental activity affected by its mechanism of action... 

• ALUHYDE •

Provides prompt symptomatic relief through its combination of... 

Formula—

- Each tablet: Dried Aluminium Hydroxide Gel, B.P.C. — 3.75 gr
- Magnesium trisilicate B.P. — 3.75 gr
- Quinalbarbitone sodium B.P.—0.50 gr
- Tinct. Belladonnae B.P. — 3 m.

Packings—Regular packs of 50;
Dispensing packs of 500.

Samples available on personally signed request of qualified physicians only

Multipax Chemicals Ltd

121 NORWOOD ROAD HERNE HILL LONDON S.E.24. Phone: TULse Hill 3276

