Amylase activity in human bile

L. A. DONALDSON, S. N. JOFFE¹, W. MCINTOSH, AND M. J. BRODIE

From the University Department of Surgery, Royal Infirmary, Glasgow, and Department of Biochemistry, Stobhill General Hospital, Glasgow

SUMMARY The mean amylase level in 42 human bile samples was 154 IU/l and there was no significant difference in the amylase activity of 32 paired serum and bile samples. Estimation of the amylase thermolability of bile showed it to be similar to that of serum. This suggests that the amylase activity in bile may have filtered through the liver from the hepatic circulation rather than refluxed from the pancreatic duct. The presence of amylase in human bile provides further evidence that the liver might have a role in the regulation of serum amylase.

While there is evidence that the amylase activity in human serum is related to hepatocellular dysfunction (Bhutta and Rahman, 1971) it is controversial whether there is a hepatic production of amylase (Janowitz and Dreiling, 1959; Nothman and Callow, 1971; Gamkhou and Scherstén, 1972; Taussig *et al.*, 1974; Skude, 1975; Otsuki *et al.*, 1976). In dogs (Nothman and Callow, 1971) and rats (Arnold and Rutter, 1963) the liver has been shown to play an important role in amylase production with canine bile containing approximately one quarter of the activity of normal serum (Singh *et al.*, 1970). Until recently the main difficulty in confirming or refuting a hepatic origin for amylase in man has been the difficulty in identification of amylase isoenzymes (Warshaw *et al.*, 1976).

Using the technique of thermolability (Donaldson *et al.*, 1977), it has been suggested that only 44% of the amylase activity in normal human serum appears to have originated from pancreas and salivary glands. The tissue(s) of origin of the remaining 56%, which is a heat stable amylase, is not established. However, intestinal mucosal amylase in the rat is known to be thermostable (Alpers and Solin, 1970). Human bile was examined to see if it contained amylase activity, and whether it had a pattern of thermolability similar to normal serum.

Methods

Bile samples (2-3 ml) were collected from patients

undergoing elective biliary surgery, who had no clinical evidence of hepatic disease and with normal liver function tests (bilirubin, AST, ALT, LDH, total protein, albumin, alkaline phosphatase, and γ GT). At laparotomy the liver was macroscopically normal. The samples were collected by direct needle puncture and aspiration, after opening the peritoneal cavity before manipulating the gallbladder or common bile duct. Patients with a fibrosed and contracted gallbladder were excluded, as were patients with an empyema or mucocoele of the gallbladder. The gallbladder samples were taken to facilitate removal of the organ and the common bile duct samples collected at needle aspiration for identification of the duct before performing operative duct cholangiography. Before collection, the common bile duct was occluded distally by digital pressure to avoid possible contamination by pancreatic duct and duodenal contents. Bile was collected from 42 patients and a preoperative serum sample was available in 32 of these patients. The amylase activity was estimated using the Phadebas amylase test (Pharmacia Ltd.) with a normal serum range of 80-300 IU/l. In 20 of the samples of bile, the amylase thermolability was also assessed using a previously described method (Donaldson *et al.*, 1977).

Results

Amylase activity was present in all samples of bile. The mean amylase content of the 42 samples of bile was 154 (SEM 16) IU/l. The mean preoperative serum amylase in the 32 patients was 179 (SEM 15) IU/l. There was no significant difference between the amylase activity in the 32 paired samples of bile and serum ($P > 0.1$), assessed either by parametric or non-parametric tests. The mean amylase thermo-

¹Address for correspondence: Mr S. N. Joffe, University Department of Surgery, Royal Infirmary, Glasgow G4 0SF, Scotland.

lability in bile was 47% (SEM 4.5), which was similar to the thermolability of normal serum.

Discussion

The possible relationship between the liver and amylase found in the serum has been extensively investigated in animals. Arnold and Rutter (1963) showed that the liver synthesised most of the amylase found in serum. Deficiency of the essential amino acid tryptophan, diminished amylase production, and administration of puromycin, a specific inhibitor of protein synthesis, caused a cessation of amylase production by the liver. In the dog, the liver appears to produce amylase (Nothman and Callow, 1971), and, in the baboon, Duane *et al.* (1972) showed that the liver was a major site of amylase destruction.

In contrast, there is little direct evidence for the existence of a hepatic amylase in man, although it has been suggested for many years that the serum amylase level is altered in patients with liver disease (Janowitz and Dreiling, 1959; Bhutta and Rahman, 1971). The identification of a hepatic isoamylase in man would give conclusive evidence for its existence.

Many workers have attempted to identify the isoenzymes of human amylase using electrophoresis, electrofocusing, or immunodiffusion techniques. In some cases only zones of activity consistent with salivary origin were identified in the serum (Benjamin and Kenny, 1974; Taussig *et al.*, 1974; Otsuki *et al.*, 1976). Other workers found additional zones of amylase activity, the origin of which was unclear (Harada *et al.*, 1974; Skude, 1975; Takeuchi *et al.*, 1975). Joseph *et al.* (1966) isolated a possible hepatic isoamylase and Warshaw *et al.* (1976) found an additional faint zone of amylase activity in one-third of normal people, which was prominent in 67% of patients with hyperamylasaemia and liver disease. The pancreas may also be able to manufacture an S-type isoamylase (Shimamura *et al.*, 1975).

Bile performs both an excretory and digestive function and, although human bile has been extensively investigated (Chenderovitch, 1973), little reference has been made to its amylase activity. Previous investigations with regard to the amylase content of bile have involved small numbers of samples with no details regarding the types of patient studied, methods of collection, or whether safeguards were taken to avoid reflux of pancreatic juice up the common bile duct. One report (Skude, 1975) examined only a single sample of bile and another (Warshaw *et al.*, 1976) investigated eight samples of bile, three of which contained no amylase activity.

In the 42 samples of bile examined, there was a similar amount of amylase activity found in human bile and normal serum. Furthermore, the amylase activity was 47% thermolabile, which is almost identical with the percentage of serum activity. In this study particular care was taken during the bile collection to prevent reflux of amylase from pancreatic duct. In view of the very high amylase activity of 120 000 IU/l in pure pancreatic juice, which is >99% thermolabile (Donaldson *et al.*, 1977) any reflux of minute amounts of pancreatic juice would considerably increase the total amount of amylase activity in the bile and its percentage thermolability.

The similarity in the total and percentage amylase thermolability in both bile and serum might suggest that the amylase activity in the bile had freely filtered from the blood flowing through the liver rather than refluxing from the pancreatic duct.

References

- Alpers, D. H., and Solin, M. (1970). The characterization of rat intestinal amylase. *Gastroenterology*, **58**, 833-842.
- Arnold, M., and Rutter, W. J. (1963). Liver amylase. III. Synthesis by the perfused liver and secretion into the perfusion medium. *Journal of Biological Chemistry*, **238**, 2760-2765.
- Benjamin, D. R., and Kenny, M. A. (1974). Clinical value of amylase isoenzyme determinations. *American Journal of Clinical Pathology*, **62**, 752-758.
- Bhutta, I. H., and Rahman, M. A. (1971). Serum amylase activity in liver disease. *Clinical Chemistry*, **17**, 1147-1149.
- Chenderovitch, J. (1973). Bile secretion. *Clinics in Gastroenterology*, **2**, 31-47.
- Donaldson, L. A., McIntosh, W. B., and Brodie, M. J. (1977). Amylase thermolability in body fluids. *Scandinavian Journal of Gastroenterology*, **12**, 637-639.
- Duane, W. C., Frerichs, R., and Levitt, M. D. (1972). Simultaneous study of the metabolic turnover and renal excretion of saliva amylase-¹²⁵I and pancreatic amylase-¹³¹I in the baboon. *Journal of Clinical Investigation*, **51**, 1504-1513.
- Gamklou, R., and Scherstén, T. (1972). Activity of α -amylase and α -1, 4-glucosidase in human liver tissue. *Scandinavian Journal of Clinical and Laboratory Investigation*, **30**, 201-207.
- Harada, K., Kitamura, M., and Ikenaga, T. (1974). Isoenzyme study of postoperative transient hyperamylasaemia. *American Journal of Gastroenterology*, **61**, 212-216.
- Janowitz, H. D., and Dreiling, D. A. (1959). The plasma amylase. Source, regulation and diagnostic significance. *American Journal of Medicine*, **27**, 924-935.
- Joseph, R. R., Olivero, E., and Ressler, N. (1966). Electrophoretic study of human isoamylases. *Gastroenterology*, **51**, 377-382.
- Nothman, M. M., and Callow, A. D. (1971). Investigations on the origin of amylase in serum and urine. *Gastroenterology*, **60**, 82-89.
- Otsuki, M., Saeki, S., Yuu, H., Maeda, M., and Baba, S. (1976). Electrophoretic pattern of amylase isoenzymes in serum and urine of normal persons. *Clinical Chemistry*, **22**, 439-444.

- Shimamura, J., Fridhandler, L., and Berk, J. E. (1975). Does human pancreas contain salivary-type isoamylase? *Gut*, **16**, 1006-1009.
- Singh, H., Pepin, J., Appert, H. E., Páirent, F. W., and Howard, J. M. (1970). Amylase and lipase secretion in the hepatic and intestinal lymph. *Archives of Surgery*, **100**, 253-256.
- Skude, G. (1975). Sources of the serum isoamylases and their normal ranges of variation with age. *Scandinavian Journal of Gastroenterology*, **10**, 577-584.
- Takeuchi, T., Matsushima, T., and Sugimura, T. (1975). Separation of human α -amylase isoenzymes by electrofocusing and their immunological properties. *Clinica Chimica Acta*, **60**, 207-213.
- Taussig, L. M., Wolf, R. O., Woods, R. E., and Deckelbaum, R. J. (1974). Use of serum amylase isoenzymes in evaluation of pancreatic function. *Pediatrics*, **54**, 229-235.
- Warshaw, A. L., Bellini, C. A., and Lee, K. H. (1976). Electrophoretic identification of an isoenzyme of amylase which increases in serum in liver diseases. *Gastroenterology*, **70**, 572-576.

The February 1979 Issue

THE FEBRUARY 1979 ISSUE CONTAINS THE FOLLOWING PAPERS

- Pure pancreatic juice studies in normal subjects and patients with chronic pancreatitis M. E. DENYER AND P. B. COTTON
- Impaired pancreatic polypeptide release in chronic pancreatitis with steatorrhoea T. E. ADRIAN, H. S. BESTERMAN, C. N. MALLINSON, C. GARALOTIS, AND S. R. BLOOM
- Release of motilin by oral and intravenous nutrients in man N. D. CHRISTOFIDES, S. R. BLOOM, H. S. BESTERMAN, T. E. ADRIAN, AND M. A. GHATEI
- Intestinal absorption under the influence of vasopressin: studies in unanaesthetised rats R. DENNHARDT, B. LINGELBACH, AND F. J. HABERICH
- Amino acid and peptide absorption after proximal small intestinal resection in the rat A. B. GARRIDO, JR., H. J. FREEMAN, Y. C. CHUNG, AND Y. S. KIM
- Absorption of antigens after oral immunisation and the simultaneous induction of specific systemic tolerance E. T. SWARBRICK, C. R. STOKES, AND J. F. SOOTHILL
- Variability of gluten intolerance in treated childhood coeliac disease B. MCNICHOLL, B. EGANMITCHELL, AND P. F. FOTTRELL
- Treatment of osteomalacia associated with primary biliary cirrhosis with parenteral vitamin D₂ or oral 25-hydroxyvitamin D₃ JULIET E. COMPSTON, L. W. L. HORTON, AND R. P. H. THOMPSON
- Serum bilirubin: a prognostic factor in primary biliary cirrhosis J. M. SHAPIRO, H. SMITH, AND F. SCHAFFNER
- Rectal biopsy in patients presenting to an infectious disease unit with diarrhoeal disease R. J. DICKINSON, H. M. GILMOUR, AND D. B. L. MCCLELLAND
- Rectal IgE cells in inflammatory bowel disease D. P. O'DONOGHUE AND PARVEEN KUMAR
- Gynaecomastia associated with cimetidine R. W. SPENCE AND L. R. CELESTIN
- Clinical trial*
Double-blind comparison of cimetidine and placebo in the maintenance of healing of chronic duodenal ulceration K. D. BARDHAN, D. M. SAUL, J. L. EDWARDS, P. M. SMITH, S. J. HAGGIE, J. H. WYLLIE, H. L. DUTHIE, AND I. V. FUSSEY
- Case reports*
Hypobetalipoproteinaemia—a variant of the Bassen-Kornzweig syndrome B. B. SCOTT, J. P. MILLER, AND M. S. LOSOWSKY
- Peutz-Jeghers syndrome associated with gastrointestinal carcinoma B. COCHET, J. CARREL, L. DESBAILLETS, AND S. WIDGREN
- Notes and activities; Books

Copies are still available and may be obtained from the PUBLISHING MANAGER, BRITISH MEDICAL ASSOCIATION, TAVISTOCK SQUARE, LONDON WC1H 9JR, price £3.50, including postage