Comparative studies of mononuclear phagocyte function in patients with Crohn's disease and colon neoplasms

W L BEEKEN, SUSAN ST ANDRE-UKENA, AND R MARY GUNDEL

From the Department of Medicine, University of Vermont, Burlington, Vermont, USA

SUMMARY Phagocytosis and cellular cytotoxicity by mononuclear phagocytes of blood and intestinal mucosa were studied in patients with Crohn's disease and large bowel neoplasms. Antibody coated sheep erythrocytes were used for phagocytic assays and cellular cytotoxicity *in vitro* was measured by 24 hour isotope release from ⁷⁵Selenium methionine-labelled RPMI 4788 human cancer cell cultures in the presence of mononuclear phagocyte-enriched effector populations. The mean percent of mononuclear phagocytes in Ficoll-Hypaque purified mononuclear cell suspensions of blood of healthy controls was 25.9 compared with 44.6 in patients with Crohn's disease, 45.6 in patients with colon neoplasms and 11.6 in intestinal mucosa. Phagocytic indices were similar in all groups, but the phagocytic capacity of mucosal macrophages was twice that of blood monocytes. Mean cytotoxicity of monocytes of patients with Crohn's disease was 12.8% compared with 22.9% for monocytes from normal controls, and 29.4% for patients with colon tumours. Mean cytotoxicity by mucosal macrophages was 18.0% compared with 13.2% by mucosal lymphocyte populations. Exposure of monocytes of Crohn's disease patients to bacterial lipopolysaccharide modestly increased cytotoxicity, but exposure did not alter phagocytosis by monocytes of patients or controls. The results indicate that monocytes of patients with Crohn's disease exhibit subnormal *in vitro* cytotoxicity. Mucosal macrophages from patients with various diseases show enhanced phagocytosis compared with blood monocytes, and they can mediate cellular cytotoxicity *in vitro*.

Mononuclear phagocytes are a prominent part of the inflammatory response in Crohn's disease and certain neoplasms, but little is known of their function in these disorders. In Crohn's disease monocyte numbers in peripheral blood are increased¹⁻⁴ and their turnover is faster.³ IgG and C₃ receptor activity,⁵ lysosomal enzyme concentrations,⁶⁻⁸ and secretion of plasminogen activator⁹ are also increased. Phagocytosis of *Candida albicans*¹⁰ and *Staphylococcus aureus*¹¹ is augmented in Crohn's disease but intracellular killing of ingested organisms is not increased. In addition, monocytes may act as suppressors of natural killer cells in Crohn's disease.¹²

Mononuclear phagocytes are also of potential

relevance to patients with colonic neoplasms from the standpoint of immune surveillance¹³ and resistance to infection.¹⁴ Compared with normal subjects, patients with certain cancers have raised numbers of circulating mononuclear phagocytes,¹⁵ and macrophages are found in significant numbers within many tumours.¹⁶ Tumour macrophages are essential for antitumour cytotoxic responses in certain *in vitro* systems.¹⁷ In human mammary tumours and melanomas, where mononuclear phagocytes may account for as much as 30% of the tumour cell mass, the percentage of macrophages correlates inversely with metastasis.¹⁸

To explore the functional status of mononuclear phagocytes in Crohn's disease and colon neoplasms further, we studied phagocytosis and cellular cytotoxicity by these cells in peripheral blood and intestinal mucosa from patients with these disorders.

Address for correspondence: W L Beeken, MD, C 311 Given Building, University of Vermont, Burlington VT, 05405 USA.

Received for publication 20 December 1982.

Methods

The subjects studied are listed in Table 1. Healthy controls had no evidence of gastrointestinal disease or intercurrent illness. Crohn's disease was documented by combined clinical, radiological, and histological criteria. According to Crohn's Disease Activity Indices,¹⁹ which ranged from 64 to 447, eight patients had inactive disease and 12 had active disease. Nine patients were on no medication, seven were taking corticosteroids in varying doses, two were taking sulphasalazine, and two were taking azathioprine. In the colon neoplasia group, five patients had Dukes A, seven had Dukes B, and four had Dukes C lesions. One patient had a large benign polyp. Intestinal mucosa was available for study from three patients with Crohn's disease, one with ulcerative colitis, one with angiodysplasia, and 10 with colonic neoplasms. All subjects gave informed consent according to the guidelines of the Human Experimentation Committee of the University of Vermont.

MONONUCLEAR PHAGOCYTE ISOLATION

Peripheral blood In patients having resections blood samples were removed immediately before surgery. Sixty millilitres of venous blood was collected in preservative free heparin (10 U/ml) and the mononuclear cells separated on Ficoll-Hypaque.²⁰ Mononuclear cells were washed three times in chilled Hank's balanced salt solution and resuspended in tissue culture medium consisting of

Eagle's minimum essential medium (MA Bioproducts, Walkersville, MD) containing 2 mM L-glutamine, penicillin (100 U/ml), streptomycin (100 mcg/ml), and gentamicin (50 mcg/ml) and 10% heat inactivated fetal calf serum or autologous serum or plasma. Mononuclear cells were counted in a haemocytometer and viability was determined by trypan blue exclusion. The percentage of mononuclear phagocytes in the mononuclear cell population was determined from alpha naphthyl-acetate esterase strains²¹ of cytocentrifuge preparations.

Monocytes were separated from lymphocytes by adherence in 75 cm² tissue culture flasks for 30–60 minutes, the period determined to be optimal by time course adherence studies. Lymphocytes were removed by vigorous washing three times, and adherent mononuclear phagocytes were chilled on ice and removed by gentle scraping with a rubber policeman. Cells in the suspension were counted and cytocentrifuge preparations were stained for esterase to determine the percentage of monocytes.

Mucosal mononuclear phagocytes A modification of the method of Bull and Bookman²² was used to separate mononuclear cells from the intestinal mucosa for subsequent isolation of mucosal macrophages. Immediately after resection representative full thickness segments of intestine were immersed in chilled Hank's balanced salt solution containing antibiotics. Using sterile technique, the mucosa was dissected from the muscularis, extensively washed, weighed, and incubated for five minutes in Hank's balanced salt solution containing 1 mM dithiothreitol and washed again in calcium-magnesium free Hank's balanced salt solution. Mucosa was then incubated for three 60 minute periods in 0.75 mM EDTA to remove epithelial cells from the lamina propria. The exposed lamina propria was next incubated overnight in tissue culture medium containing 0.075 mg/ml collagenase (Type IV, Sigma Chemical Co, St Louis, MO, or CLSPA, Worthington Biochemical Corp, Freehold, NJ) at 37°C with gentle stirring. Mononuclear cells were filtered through a siliconised mesh, number 300, and further purified on a Ficoll-Hypaque gradient. Cells were washed, aliquots removed for study, and the remaining mononuclear cells were transferred to tissue culture flasks for adherence of mononuclear phagocytes as detailed above. Viability of mucosal mononuclear cell suspensions routinely exceeded 85%, and were usually above 90%.

PHAGOCYTOSIS OF ANTIBODY COATED SHEEP ERYTHROCYTES

Phagocytosis by adhered cells 5×10^5 mononuclear

Table 1 Subjects studied

Peripheral blood					
	n	Age range	M:F	Phagocytosis n	Cytotoxicity n
Healthy controls	20	25–58	11:9	14	14
Crohn's disease	20	24–54	9:11	11	12
Colonic neoplasm	17	40–87	8:9	13	12
Intestinal mucosa					
	n	Segment		Phagocytosis n	Cytotoxicity n
Crohn's disease	3	Normal ileum (2)		1	3
		Abnormal ileum (1)			
Ulcerative colitis	1	Normal right colon		1	0
Angiodysplasia	1	Normal right colon		0	1
		Normal colon			
Colon neoplasm	10	Normal colon		8	4

phagocytes suspended in 0.5 ml tissue culture medium (fetal calf serum) were spread evenly on plastic cover slips (Thermanox 1.5, Lux Scientific Corp, Newbury Park, CA) and incubated for one hour at 37°C in 5% CO₂. Adhered cells were rinsed with phosphate buffered saline and incubated for another hour with 0.5 ml of IgG coated sheep erythrocytes (Cordis Laboratory, Miami, FL). Cover slips were immersed in distilled water for 20 seconds to lyse unphagocytosed erythrocytes, rinsed in phosphate buffered saline for five seconds, air dried, fixed, and stained for alpha naphthylacetate esterase.

Phagocytosis by cell suspensions 5 × 10⁵ cells per 0.5 ml tissue culture medium (fetal calf serum) were mixed with 0.5 ml of IgG coated erythrocytes in 15 ml test tubes and the suspensions were incubated at 37°C in CO₂ for 60 minutes with frequent shaking. The suspensions were then centrifuged, the supernatants removed, and the pellets resuspended in 5 ml ammonium chloride, 0.9%, to lyse erythrocytes. Suspensions were again centrifuged, resuspended in tissue culture medium (fetal calf serum) and cytocentrifuge preparations were stained for alpha naphthylacetate esterase.

Results of the phagocytosis studies were expressed as the percent of 100 or more mononuclear phagocytes showing erythrocyte ingestion (phagocytic index) and the mean number of erythrocytes ingested by 20 randomly selected mononuclear phagocytes (phagocytic capacity).

CELLULAR CYTOTOXICITY

Cytotoxicity assays were conducted with effector populations of mixed mononuclear cells (lymphocytes and mononuclear phagocytes, mononuclear cells) and adherent cells (mononuclear phagocyte-enriched, mononuclear phagocytes) in 10% fetal calf serum or autologous serum. Target cells were RPMI 4788 human colon cancer cells grown in tissue culture as adherent monolayers (kindly supplied by Dr M F Kagnoff, University of Californian, San Diego, La Jolla, CA). RPMI 4788 cells were labelled with ⁷⁵seleno-methionine (⁷⁵Se RPMI 4788) by washing in methionine-free Ham's solution, mixing with 50 to 100 µc of ⁷⁵Se and incubating in Falcon Microtiter II plates for a 24 hour period in humidified 5% CO₂ at 37°C. Labelled targets were washed three times and inoculated with medium alone and with suspensions of mononuclear cells and mononuclear phagocyte effector cell populations at effector:target ratios of 50 viable lymphocytes or mononuclear phagocytes per target cell. Cytotoxicity assays were done in quintuplicate except for several instances when a limited yield of

effectors required they be done in triplicate or quadruplicate. Microtitre plates were incubated for four and 24 hours at 37°C in 5% CO₂, and the radioactivity in the supernatant solutions and in the target cell monolayers was determined. Cytotoxicity was calculated by the following equations:

$$\text{Total percent release} = \frac{\text{CPM supernatant}}{\text{CPM supernatant} + \text{CPM cells}} \times 100$$

$$\text{Cytotoxicity} = \frac{\% \text{ release in test condition} - \% \text{ spontaneous release}}{100 - \% \text{ spontaneous release}} \times 100$$

Using cytospin values of % mononuclear phagocytes in mononuclear cells populations, the % mononuclear cell cytotoxicity due to mononuclear phagocyte effectors was calculated as follows:

$$\% \text{ MNC cytotoxicity due to MP} = \frac{\% \text{ MP cytotoxicity} \times \% \text{ MP in MNC}}{\% \text{ MNC cytotoxicity}}$$

These calculations assumed that non-mononuclear phagocyte adherent cells were not cytotoxic. Thus values of mononuclear phagocyte cytotoxicity, mononuclear cell cytotoxicity and percent mononuclear cell cytotoxicity due to mononuclear phagocytes were available for comparison.

The effect of lipopolysaccharide (S. Typhosa 0901, Difco Laboratories, Detroit, MI) was evaluated in phagocytosis studies by preincubating mononuclear phagocytes in 50 µg/ml for 1 hour before exposure to antibody coated erythrocytes. The effect on cytotoxicity was assessed by addition of 50 µg/ml lipopolysaccharide during cytotoxicity assays.

Results were statistically analysed for significant differences by paired and unpaired Student's *t* tests.

Results

The percentages of mononuclear phagocytes in whole mononuclear cell suspensions and adherent populations in various groups are given in Table 2. Mononuclear phagocytes comprised a significantly higher proportion of the mononuclear cell population in Crohn's disease and cancer patients compared with controls (*p* < 0.001), but mucosal suspensions had significantly lower proportions of mononuclear phagocytes compared with normal blood (*p* < 0.01). The percentage of mononuclear phagocytes in adherent populations from mucosa was significantly lower than in blood of controls and Crohn's disease (*p* < 0.001). The esterase negative

Table 2 Mean percentages $\pm$ SEM of mononuclear phagocytes (MP) in mononuclear cell suspensions (MNC) and adherent populations.

	Control blood	Crohn's disease blood	Colon cancer blood	Intestinal mucosa
Percent MP in MNC	25.9 $\pm$ 3.9 (13)	44.6 $\pm$ 4.7 (16)	45.6 $\pm$ 4.3 (14)	11.6 $\pm$ 2.5 (8)
Percent MP in adherent population	67.5 $\pm$ 4.2 (15)	77.9 $\pm$ 4.3 (15)	64.4 $\pm$ 5.5 (14)	41.1 $\pm$ 9.0 (8)

() = number of subjects

adherent cells were presumably B-lymphocytes.²³

The results of phagocytosis studies are given in Table 3. Phagocytic indices in suspensions were somewhat less than in adhered cells. The phagocytic capacity ranged from 2.7 to 4.1 in peripheral blood for all groups, and was nearly twice that for mucosal macrophages. Exposure of blood monocytes to lipopolysaccharide had no effect on either the phagocytic index or the phagocytic capacity in controls and patients with Crohn's disease.

Cytotoxicity of mononuclear phagocytes in four hour studies was minimal. Results of 24 hour assays are given in Tables 4 and 5. Cytotoxicity was lowest in patients with Crohn's disease and highest in patients with colonic neoplasms. Cytotoxicity by mucosal mononuclear phagocytes was similar to that of monocyte-enriched preparations of peripheral blood of healthy subjects, and was actually greater than cytotoxicity of mononuclear cells. Results in fetal calf serum and autologous serum were similar. Mean cytotoxicity by mixed mononuclear cell populations (lymphocytes and mononuclear phagocytes) was 42.2% for healthy controls, 22.3% for patients with Crohn's disease, 30.7% for patients with colon carcinoma and 13.3% for intestinal

mucosa. Assuming that the only cytotoxic cells in the adherent populations were mononuclear phagocytes, the remainder being non-cytotoxic B-cells,²³ the percentages of cytotoxicity of mixed mononuclear cells populations attributable to mononuclear phagocytes were 15, 27, 44 and 31 for controls, Crohn's disease, cancer and intestinal mucosa respectively. The percentage in cancer patients was significantly higher than for controls ($p < 0.001$).

Lipopolysaccharide had a modest stimulatory effect on cytotoxicity by blood monocytes of patients with Crohn's disease, but did not increase monocyte cytotoxicity of healthy controls. (Table 6.)

Discussion

These studies are consistent with previous observations of increased numbers of mononuclear phagocytes in patients with Crohn's disease and document similar changes in patients with colon neoplasms. We also confirm that mononuclear phagocytes comprise approximately 10% of the mononuclear population of the lamina propria of

Table 3 Results of phagocytic studies

	Control blood	Crohn's disease blood	Colon cancer blood	Intestinal mucosa
Phagocytic index - suspension	87.6 $\pm$ 2.0 (14)	85.5 $\pm$ 8.0 (11)	86.3 $\pm$ 2.4 (10)	67.6 $\pm$ 8.7 (6)
Phagocytic index - adhered	92.7 $\pm$ 1.4 (14)	89.8 $\pm$ 3.3 (11)	94.0 $\pm$ 1.0 (13)	87.6 $\pm$ 6.0 (10)
Phagocytic index - adhered + LPS	90.0 $\pm$ 3.1 (9)	91.1 $\pm$ 3.3 (7)		
Phagocytic capacity - suspension	3.1 $\pm$ 0.2 (14)	2.7 $\pm$ 0.3 (11)	3.6 $\pm$ 0.3 (13)	5.5 $\pm$ 1.0 (6)
Phagocytic capacity - adhered	3.7 $\pm$ 0.2 (14)	3.2 $\pm$ 0.2 (11)	4.11 $\pm$ 0.2 (13)	8.1 $\pm$ 0.7 (10)
Phagocytic capacity - adhered + LPS	3.3 $\pm$ 0.03 (9)	3.4 $\pm$ 0.4 (7)		

Phagocytic index = percent of mononuclear phagocytes ingesting erythrocytes. Phagocytic capacity = number of erythrocytes ingested per mononuclear phagocyte. LPS = lipopolysaccharide. () = number of subjects studied. Results are means $\pm$ SEM.

Table 4 Results of 24 hour cytotoxicity studies

	Control blood	Crohn's disease blood	Colon cancer blood	Intestinal mucosa
MP cytotoxicity	22.9±6.3 (14)	12.8±4.1 (14)	29.4±8.3 (12)	18.0±5.2 (8)
MNC cytotoxicity	42.2±10.7 (12)	22.3±4.3 (15)	30.7±5.5 (12)	13.2±4.2 (8)
Percent MNC cytotoxicity due to MP	15.0±3.0 (13)	27.0±6.8 (15)	43.9±7.7 (12)	31.0±16.2 (8)

Means ± standard errors of the means.

MP = mononuclear phagocyte; MNC = mononuclear cell. () = number studied.

the distal human intestine. In this study a significant number of adherent cells were esterase negative and therefore did not qualify as mononuclear phagocytes. Extrapolating from animal studies these cells were presumably B-lymphocytes, and thus were neither cytotoxic nor phagocytic. The consistently lower phagocytic index in suspensions, especially in mucosal suspensions, is unexplained, but could indicate altered Fc receptor function in a subgroup of the mononuclear phagocytes population. The phagocytic capacity of circulating monocytes of all study groups was similar, but a significantly higher phagocytic capacity was found in mucosal macrophages. These cells were apparently stimulated or activated within the intestinal mucosa. Among many potential mediators to which mucosal macrophages are exposed, lipopolysaccharide would be expected to be important. Despite contrary results in animal studies,²⁴ however, under the conditions of our studies, lipopolysaccharide had no effect on phagocytic function of blood monocytes of

controls and Crohn's disease patients.

The lowest cytotoxicity values occurred in mononuclear phagocytes and mononuclear cell effectors of blood from patients with Crohn's disease, and in mononuclear cells of intestinal mucosa. The highest values occurred in mononuclear phagocytes from patients with colonic neoplasms. The subnormal cytotoxicity of mononuclear phagocytes in Crohn's disease may be related to rapid monocyte turnover and resulting immaturity, but purified human monocytes reportedly lose their cytotoxic capability during maturation *in vitro*.²⁵ Cytotoxicity was minimal at four hours and unheated autologous serum was not cytotoxic, indicating that the observed cytotoxicity was not an antibody mediated process.²⁶⁻²⁸ Thus we surmise that cell lysis occurred by non-specific cell mediated cytotoxicity. It is known that human monocytes effect cytotoxicity by spontaneous cell mediated mechanisms,²⁹ as well as by antibody dependent cellular cytotoxicity,^{25 29} and by

Table 5 Percent cytotoxicity of mucosal and blood mononuclear cell (MNC) and mononuclear phagocyte (MP) effectors. Assays conducted in 10% fetal calf serum at effector:target ratios of 50 lymphocytes per target for MNC and 50 esterase positive macrophages per target for MP

Patient group	Source of mucosa	Blood			Mucosa		
		MNC	MP	% MNC due to MP	MNC	MP	% MNC due to MP
Crohn's disease	Abnormal ileum	18.3	3.2	6	12.9*	15.1*	7
	Normal ileum	21.6	12.4	42	0.3*	17.6*	100
	Normal ileum	11.3	17.5	6	21.2	15.5	4
Colon Ca	Normal colon (A)	60.5	16.5	20	31.3	47.6	20
	Normal colon (C)	5.5	15.8	100	1.4	23.0	100
	Normal colon (A)	32.1	16.0	28	6.0	4.1	4
	Normal colon (A)	19.2	22.0	26	9.9	3.5	3
Angiodysplasia	Normal colon	20.3	11.9	19	23.0	17.8	10
M		23.7	14.4	31	13.2	18.0	31
SEM		6.3	2.1	11.5	4.2	5.2	16.2

* Conducted in autologous serum.

M = mean; SEM = standard error of the mean. () = Dukes classification.

Table 6 Effect of lipopolysaccharide (LPS) on 24 hour cytotoxicity by mononuclear cells (MNC) and mononuclear phagocytes (MP) of patients with Crohn's disease and controls matched for age and sex

	Matched control			
	AS		AS + LPS	
	MNC	MP	MNC	MP
Subject				
1	13.1	11.4	53.7	2.0
2	117.3	70.6	89.1	54.8
3	73.9	18.8	85.1	18.0
4	36.8	11.6	67.4	16.6
5	83.0	72.6	99.0	77.0
M	64.8	37.0	78.9	33.7
SEM	20.4	15.9	9.1	15.5
Crohn's disease				
1	11.7	5.5	15.0	5.5
2	25.3	9.3	44.5	10.5
3	22.8	4.6	26.9	9.8
4	11.1	3.4	21.7	7.0
5	19.5	5.5	20.1	13.8
M	18.1	5.7	25.6	9.3
SEM	3.2	1.1	5.7	1.6

AS = autologous serum or plasma. M = mean. SEM = standard error of the mean.

augmentation of lymphocyte cytotoxicity for adherent targets.³⁰ Lymphokines³¹ and lipopolysaccharide³⁰⁻³³ enhance cytotoxicity by mononuclear phagocytes. In man, hydrocortisone and acetylsalicylic acid reportedly augment monocyte cytotoxicity,³⁴ but corticosteroids inhibit cytotoxicity of interferon stimulated peritoneal macrophages of mice.³⁵ Animal studies also indicate that spontaneous cytotoxicity by natural killer cells is inhibited by starvation,³⁶ but this important aspect has not been examined using human mononuclear phagocytes. All of these factors could be influencing the cytotoxicity observed in this study, and further investigations will be required to obtain a better understanding of these events.

These studies also show that adherent mucosal macrophages are capable of effecting cytotoxicity for colon cancer cells grown in culture. In both blood and mucosa mononuclear phagocytes were calculated to account for approximately 30% of the cytotoxic capacity of mixed mononuclear cell effectors. Adherent mucosal macrophages were often more cytotoxic than the corresponding mucosal mononuclear populations and the circulating monocytes of the same patient. The small numbers of resections from which macrophage

cytotoxicity data were available, and the diverse sources preclude any conclusions regarding altered functions in specific patient groups. Moreover, the mononuclear phagocyte populations contained esterase negative mononuclear cells, and the methods used did not permit direct assay of cytotoxicity by non-adherent mononuclear phagocytes of mucosa. New techniques for isolating purer populations of mononuclear phagocytes have been reported,²⁵ and the application of these methods will be of great advantage in future studies of macrophages of intestinal mucosa.

In conclusion, we confirm the increase of mononuclear phagocytes in peripheral blood of patients with Crohn's disease and document similar findings in colonic cancer. Mononuclear phagocytes comprise approximately 10% of the mononuclear cell population of human intestinal mucosa. This value is one-fourth that found in blood of patients with Crohn's disease and colon cancer and one-half that found in healthy subjects. Phagocytosis by circulating monocytes of the various study groups was similar, but mucosal macrophages ingested significantly more opsonised erythrocytes. Adherent mononuclear phagocytes of blood of normal subjects and colon cancer patients and adherent mucosal macrophages showed active cytotoxicity for neoplastic epithelial cells *in vitro*. Cytotoxicity by blood mononuclear phagocytes in patients with Crohn's disease was subnormal, coinciding with similar defects reported for natural killer cells in these patients.¹² Lipopolysaccharide had little effect on phagocytes of patients with Crohn's disease. Further studies of these functional differences in Crohn's disease and colon cancer are needed to determine if these changes could be involved in the pathogenesis of these disorders.

This study was supported by PHS Grants AM 21564 and S07 05429-20 and General Clinical Research Centers Grant RR 109. Jennifer Penhallow provided expert typing assistance.

References

- 1 Thayer WR, Charland C, Field C. Subpopulations of circulating white blood cells in inflammatory bowel disease. *Gastroenterology* 1976; **71**: 379-84.
- 2 Mee AS, Berney J, Jewell DP. Monocytes in inflammatory bowel disease: absolute monocyte counts. *J Clin Pathol* 1980; **33**: 917-20.
- 3 Meuret G, Bitzi A, Hammer B. Macrophage turnover in Crohn's disease and ulcerative colitis. *Gastroenterology* 1978; **74**: 501-3.

- 4 Auer IO, Wechsler W, Ziemer E, Malchow H, Sommer H. Immune status in Crohn's disease. I. Leukocyte and lymphocyte subpopulations in peripheral blood. *Scand J Gastroenterol* 1978; **13**: 561-71.
- 5 Schmidt M, Douglas S. Monocyte IgG receptor activity dynamics and modulation - normal individuals and patients with granulomatous diseases. *J Lab Clin Med* 1977; **89**: 332-40.
- 6 Mee AS, Jewell DP. Monocytes in inflammatory bowel disease: monocyte and serum lysosomal enzyme activity. *Clin Sci* 1980; **58**: 295-300.
- 7 Ganguly NK, Lloyd B, Price CP *et al*. Acid hydrolases in monocytes from patients with inflammatory bowel disease, chronic liver disease and rheumatoid arthritis. *Lancet* 1978; **1**: 1073-5.
- 8 Doe WF, Temple C, Dorsman B, Golder J. Monocyte function in chronic inflammatory bowel disease. I. Lysosomal enzyme activity. *Aust NZ J Med* 1980; **10**: 367.
- 9 Doe WF, Dorsman B, Temple C, Golder J. Monocyte function in chronic inflammatory bowel disease. *Aust NZ J Med* 1980; **10**: 368.
- 10 Whorwell PJ, Bennett P, Tanner AR, Wright R. Monocyte function in Crohn's disease and ulcerative colitis. *Digestion* 1981; **22**: 271-5.
- 11 Mee AS, Szawatakowski M, Jewell DP. Monocytes in inflammatory bowel disease: phagocytosis and intracellular killing. *J Clin Pathol* 1980; **33**: 921-5.
- 12 Auer IO, Ziemer E, Sommer H. Immune status in Crohn's disease. V. Decreased in vitro natural killer cell activity in peripheral blood. *Clin Exp Immunol* 1980; **42**: 41-9.
- 13 Hibbs JB, Chapman HA, Weinberg JB. The macrophage as an antineoplastic surveillance cell: biological perspectives. *J Reticuloendothelial Soc* 1978; **24**: 549-70.
- 14 Estevez M, Fen L, Bachman AE, Pavlovsky A. Effective function of peripheral blood monocytes in patients with Hodgkin's disease and non-Hodgkin's lymphoma. *Cancer* 1980; **46**: 299-302.
- 15 Wood GW, Neff JR. A reevaluation of B-lymphocyte levels in peripheral blood from cancer patients. *J Natl Cancer Inst* 1978; **61**: 715-8.
- 16 Wood GW, Gollahon KA. Detection and quantitation of macrophage infiltration into primary human tumors with use of cell surface markers. *J Natl Cancer Inst* 1977; **59**: 1081-7.
- 17 Woodward JG, Daynes RA. Cell mediated immune response to synergistic UV-induced tumors. *Cell Immunol* 1978; **41**: 304-19.
- 18 Alexander P, Eccles SA, Gauci CLL. The significance of macrophages in human and experimental tumors. *Ann NY Acad Sci* 1976; **276**: 124-33.
- 19 Best WR, Becktel JM, Singleton JW, Kern F. Development of a Crohn's disease activity index. *Gastroenterology* 1976; **70**: 439-44.
- 20 Boyum A. Isolation of lymphocytes, granulocytes and macrophages. *Scand J Immunol* 1976; **5**: (suppl) 15.
- 21 Li CY, Yam LT, Crosby WH. Histochemical characterization of cellular and structural element of human spleen. *J Histochem Cytochem* 1972; **20**: 1049-58.
- 22 Bull DM, Bookman MA. Isolation and functional characterization of human intestinal mucosal lymphoid cells. *J Clin Invest* 1977; **59**: 966-74.
- 23 Nathan CF, Asofsky R, Terry WD. Characterization of the nonphagocytic adherent cell from the peritoneal cavity of normal and BCG treated mice. *J Immunol* 1977; **118**: 1612-21.
- 24 Schubert RD, David JR. Stimulation of guinea pig macrophage pinocytosis by lipopolysaccharide (LPS): evidence that LPS acts directly on the macrophage. *Cell Immunol* 1980; **55**: 166-73.
- 25 Stevenson HC, Katz P, Wright DG *et al*. Human blood monocytes: characterization of negatively selected human monocytes and their suspension cell culture derivatives. *Scand J Immunol* 1981; **14**: 243-56.
- 26 Nelson DL, Bundy BM, Strober W. Spontaneous cell-mediated cytotoxicity by human peripheral blood lymphocytes *in vitro*. *J Immunol* 1977; **119**: 1401-5.
- 27 Russel JH, Masakowski VR, Dobos CB. Mechanisms of immune lysis. I. Physiological distinction between target cell death mediated by cytotoxic T lymphocytes and antibody plus complement. *J Immunol* 1980; **124**: 1100-5.
- 28 Sanderson CJ, Thomas JA. The mechanism of K cell (antibody-dependent) cell mediated cytotoxicity. I. The release of different cell components. *Proc Roy Soc Lond B* 1977; **197**: 407-15.
- 29 Horwitz DA, Kight N, Temple A, Allison AC. Spontaneous and induced cytotoxic properties of human adherent mononuclear cells. Killing of non-sensitized and antibody-coated non-erythroid cells. *Immunology* 1979; **36**: 221-8.
- 30 DeVries JE, Mendelsohn J, Bont WS. Requirement for monocytes in the spontaneous cytotoxic effects of human lymphocytes against non-lymphoid target cells. *Nature* 1980; **283**: 574-6.
- 31 Cameron D, Churchill WH. Cytotoxicity of human macrophages for tumor cells. *J Clin Invest* 1979; **63**: 977-84.
- 32 Doe WF, Henson PM. Macrophage stimulation by bacterial lipopolysaccharides. I. Cytolytic effect on tumor target cells. *J Exp Med* 1978; **148**: 544-56.
- 33 Taramelli D, Holden HT, Varesio L. In vitro induction of tumoricidal and suppressor macrophages by lymphokines: possible feedback regulation. *J Immunol* 1981; **126**: 2123-8.
- 34 Kleinerman ES, Lovie JS, Wahl LM, Muchmore AV. Pharmacology of human spontaneous monocyte mediated cytotoxicity. I. Enhancement by salicylates and steroids. *Arthritis Rheumatism* 1981; **24**: 774-80.
- 35 Schultz RM, Chirigos MA, Stoychkov JN, Pavlidis NA. Factors affecting macrophage cytotoxic activity with particular emphasis on corticosteroids and acute stress. *J Reticuloendothelial Soc* 1979; **26**: 83-92.
- 36 Saxena RK, Saxena QB, Adler WH. Regulation of natural killer activity during starvation. *Indian J Exp Biol* 1980; **18**: 1383-6.