Alpha₁-antitrypsin, a reliable endogenous marker for intestinal protein loss and its application in patients with Crohn's disease

U KARBACH, K EWE, AND H BODENSTEIN

From the I. Medizinische Klinik und Poliklinik, Mainz, FR Germany.

SUMMARY Intestinal protein loss is generally determined by radio-labelled macromolecules. Alpha₁-antitrypsin has been proposed as an endogenous marker for protein losing enteropathy, but different opinions exist about its reliability. In 25 patients with Crohn's disease faecal protein loss was studied with intestinal alpha₁-antitrypsin (α_1 AT) clearance. Simultaneously, in 10 patients α_1 AT clearance was compared with faecal ⁵¹Cr clearance after intravenous ⁵¹Cr-albumin injection. There was a linear relation ($p < 0.05$) between α_1 AT clearance and ⁵¹Cr clearance in these cases. In all patients α_1 AT clearance was raised above control values. α_1 AT clearance, however, did not correlate with the activity index of Crohn's disease.¹ This index does not contain direct criteria of intestinal inflammation, does not take into account localisation or extent of inflammation, and includes complications such as extraintestinal manifestations, fistuli, stenoses not necessarily related to actual mucosal involvement. It is concluded that α_1 AT is a reliable marker for intestinal protein loss and that the intestinal changes of Crohn's disease generally lead to an increased protein exudation into the gut.

Gastrointestinal loss of plasma proteins can be detected by a variety of labelled macromolecules: ⁵⁹Fe-labelled dextran² and ¹³¹I-PVP³ are not split by digestive enzymes; the radioactive isotopes ⁵¹Cr-albumin,⁴ ⁶⁷Cu-ceruloplasmin⁵ or ⁹⁵Nb-albumin⁶ are poorly absorbed by the alimentary tract. These methods are expensive and the patients are exposed to radioactivity and therefore cannot be used routinely as screening-tests or to monitor the course of illness. Crossley and Elliot⁷ showed that α_1 AT, a protease inhibitor synthesized by the liver and present in serum at a concentration of 2-5 g/l, is not degraded by digestive enzymes nor reabsorbed by the intestine. It has been shown to be a reliable endogenous marker of intestinal protein loss by Bernier and colleagues.⁸ These authors reported that both faecal α_1 AT concentration and α_1 AT clearance were significantly higher in patients with suspected exudative enteropathy than in controls. Their results were not confirmed by Haeney *et al*,⁹ who compared α_1 AT concentration and α_1 AT

clearance with the conventional ⁵¹Cr-albumin method. According to Keaney and Kelleher¹⁰ the contradictory results could be caused by a difference in methods and by comparison of different parameters.

We examined the validity of α_1 AT as a reliable indicator of intestinal protein loss in patients with Crohn's disease, comparing α_1 AT clearance with the clearance of ⁵¹Cr-albumin and studied problems of methodology.

Finally, we tried to correlate intestinal α_1 AT clearance with site, extent, and activity of intestinal inflammation or resection in Crohn's disease.

Methods

PATIENTS

Alpha₁-antitrypsin clearance was determined in 10 healthy controls and in 25 patients with Crohn's disease (mean age 31.6 ± 2.5 , range 18-63 years; 17 women, eight men). The site and extent of inflammation was documented radiologically and/or by endoscopy and histology (Table). Ten patients had undergone intestinal resection, five of them suffered from anastomosal relapse. Twelve patients had complications: intestinal stenosis occurred in six

Address for correspondence: Prof Dr med K Ewe, I Med Klinik und Poliklinik, Johannes Gutenberg Universität, Langenbeckstr 1, D 6500 Mainz, FR Germany.

Accepted for publication 22 October 1982

Table Morphological, clinical, and laboratory data of patients

Patient (no.)	Resection	Involvement	Complications	CDAI	α_1 AT clearance (ml/d)	ESR (mm/1 h)	Serum albumin (g%)	HCT (%)	Treatment
1. MD	Ileoascendost	0	0	27	20.7	15	3.6	41	0
2. RC	Ileoascendost	0	Anastomosal stenosis	354	46.2	4	4.2	42	0
3. GM	Ileoascendost	0	0	115	40.6	20	3.4	31	0
4. HE	Ileotransversostomy	0	0	35	44.7	8	4.8	38	0
5. BU	Subtotal colectomy	0	0	29	39.5	10	4.1	40	0
6. EH	Ileoascandost	AR	0	118	203.6	—	3.5	42	SASP, ST
7. WE	Ileotransversostomy	AR	Anastomosal stenosis	158	307.8	3	4.5	46	SASP, ST
8. GP	Ileotransversostomy	AR	Anal fistula	92	34.5	38	4.0	37	METRO, ST
9. FM	Ileotransversostomy	AR	0	136	105.9	13	3.7	39	SASP, ST
10. KP	Ileosigmoidostomy	AR	Anal fistula	431	40.6	22	3.2	44	METRO, ST
11. SH	0	tC	0	50	54.5	20	4.3	37	SASP
12. KH	0	TI (10 cm)	Stenosis (TI) anal fistula	227	23.0	9	4.8	43	SASP, ST
13. WG	0	TI (20 cm), tC	EIM	151	53.3	26	3.2	41	SASP, ST
14. HM	0	TI (20 cm)	0	64	105.7	14	4.3	38	SASP, ST
15. JM	0	TI (30 cm)	Stenosis (TI)	58	57.2	10	3.6	38	SASP
16. MW	0	TI (25 cm)	Stenosis (TI)	88	59.1	26	3.9	40	SASP, ST
17. SB	0	TI (30 cm), CA	Enterol fistula	124	25.3	42	3.7	38	SASP, ST
18. SU	0	TI (20 cm), CA	0	244	103.4	13	3.7	39	SASP, ST
19. RM	0	TI (10 cm) CT (20 cm)	Stenosis (TI)	242	26.7	26	3.8	35	SASP, ST
20. FG	0	TI (5 cm), tC	Enterol fistula EIM	485	80.9	18	3.7	41	SASP, ST
21. RP	0	TI (30 cm), tC	Enterovaginal fistula	261	123.0	115	2.7	33	SASP, ST
22. WL	0	TI (30 cm), tC	0	125	377.7	13	3.9	39	SASP, ST
23. DH	0	TI (30 cm), tC	0	182	168.6	66	3.1	41	SASP, ST
24. ZK	0	tC	0	48	42.1	24	4.4	38	SASP
25. ZI	0	tC	0	185	67.1	65	2.5	22	SASP, ST

Abbreviations: TI=terminal ileum, tC=total colon, CA=ascending colon, TC=transverse colon, CD=descending colon, AR=relapse at anastomosis, EIM=extraintestinal manifestation, SASP=salazopyridine, ST=methylprednisolone, METRO=metronidazole.

patients (one of them also having anal fistula), extraintestinal manifestations (arthritis and erythema nodosum) in two patients (one also having anal fistula), perianal complications only in two patients and internal fistulae occurred in two patients.

In all patients the CDAI¹ was established. In the Table data on ESR, serum albumin concentrations, haematocrit and therapy are also given.

In 10 patients α_1 AT clearance was compared with faecal excretion of intravenously injected ⁵¹Cr-albumin and with the intestinal ⁵¹Cr clearance.

DETERMINATION OF ALPHA₁-ANTITRYPSIN

Alpha₁-antitrypsin of serum and faeces was determined by radial immunodiffusion. The diameter of the precipitation ring of the plates

containing a monospecific antigen against α_1 AT (Behringwerke) was measured after 72 hours and compared with a reference curve established from the diameters of rings of a standard solution. Faeces and sera were kept at -20°C until used. The volume of the three day stool samples was calculated by weight and specific gravity. Specific gravity was determined by weighing a volume of 5 ml of the homogenised stool. An aliquot of 2 g of the homogenised stool was diluted 1:4 with normal saline or for comparison with aqua dest, and again homogenised for 10–60 minutes. After centrifugation at 1500–3000 g for 10–30 minutes 5 μ l of the supernatant was applied to the plates. The sera obtained at the beginning of stool sampling were diluted 1:5 in normal saline and 5 μ l of the samples were applied to the plates. In five samples the

determinations were repeated after storage of faeces and sera for eight months at -20°C . Faecal $\alpha_1\text{AT}$ loss was calculated from faecal $\alpha_1\text{AT}$ concentration and stool volume. $\alpha_1\text{AT}$ clearance was calculated by the formula

$$\alpha_1\text{AT clearance} = \frac{\text{faecal volume} \times \alpha_1\text{AT faecal concentration}}{\alpha_1\text{AT serum concentration}}$$

^{51}Cr -ALBUMIN METHOD

The radioactivity of the three day stool samples (Behringwerke) was measured, after intravenous injection of $10 \mu\text{Ci } ^{51}\text{Cr}$ -albumin, with a gamma counter. The ^{51}Cr -serum concentration was calculated from the administered ^{51}Cr -dose and from the total serum volume calculated by conventional formulas.¹¹ The formula for computing the ^{51}Cr clearance corresponded to that for $\alpha_1\text{AT}$.

Results

There was a good correlation between $\alpha_1\text{AT}$ clearance and ^{51}Cr excretion ($r=0.84$, $\alpha=5\%$) or ^{51}Cr clearance ($r=0.87$, $\alpha=5\%$). The relation between $\alpha_1\text{AT}$ and ^{51}Cr clearance as expressed by a linear equation was $y=0.73x+4.42$ (Fig. 1). The different methods of treating the faeces with normal saline or aqua dest, the duration of homogenisation, and the rpm of centrifugation, had no influence on the results. After storage of the stool and the serum at -20°C for eight months the recovery rate of $\alpha_1\text{AT}$ was more than 90%.

In our experiments the faecal loss of $\alpha_1\text{AT}$ in

Fig. 1 Relation between $\alpha_1\text{AT}$ clearance and ^{51}Cr -albumin clearance.

Fig. 2 Faecal $\alpha_1\text{AT}$ loss and intestinal $\alpha_1\text{AT}$ clearance in patients with Crohn's disease and controls.

patients with Crohn's disease (average 308 mg/d, range 22–785 mg/d) was higher than in controls (average 54 mg/d, range 0–136 mg/d) (Fig. 2). When the clearance of $\alpha_1\text{AT}$ was calculated there was a better separation (Fig. 2) of patients with Crohn's disease (average 93 ml/d, range 19–367 ml/d) from controls (5 ml/d on an average, range 0–22 ml/d) and with only one exception $\alpha_1\text{AT}$ clearance was raised above control values in all patients. As shown in Fig. 3 there was no difference between patients with involvement of the small intestine only, and the colon only. The clearance of $\alpha_1\text{AT}$ in patients with skip lesions of the large bowel was similar to those with total colonic inflammation and the same findings applied to the inflammation of the ileum. Three patients with inflammation of both colon and ileum had higher values than those who had only the ileum or the colon involved. On the other hand three patients with relapse of Crohn's disease over a small segment adjacent to the anastomosis had higher values than others with total colonic and/or ileal involvement. Even after curative resection

Fig. 3 Site and extent of Crohn's disease and α_1 AT clearance (rec=recurrence, term ileum=terminal ileum, part colon=partial colon).

faecal α_1 AT was raised. The increase of faecal α_1 AT in the resected patients was about twice above control values. There was no correlation between activity index and α_1 AT clearance (Fig. 4). Patients with inflammation in remission (activity index <150) had values of α_1 AT clearance similar to patients with a CDAI higher than 400.

In patients with complications such as fibrous

Fig. 4 Activity index (CDAI) in patients with Crohn's disease in comparison to intestinal α_1 AT clearance.

Fig. 5 Activity index (CDAI) in patients with Crohn's disease and under consideration of complications such as intestinal stenosis, anal fistula, and extraintestinal manifestations.

strictures, anorectal fistulation, or extraintestinal manifestations, the mean CDAI (206 ± 43) was significantly higher ($p < 0.02$) than in patients without complications (99 ± 19) (Fig. 5).

The mean α_1 AT clearance in patients without complications (104.9 ± 26.5 ml/d) was raised above

Fig. 6 Intestinal α_1 AT clearance in patients with Crohn's disease and under consideration of complications.

the values of patients with complications (73.1 ± 22.8 ml/d) (Fig. 6).

Discussion

Similar to the results of Florent and coworkers¹² we found a linear relationship of the intestinal protein loss measured by ^{51}Cr or by $\alpha_1\text{AT}$ clearance. The slope of the curve comparing $\alpha_1\text{AT}$ and ^{51}Cr -albumin clearance (Fig. 1) should be unity provided both proteins follow the same mechanism of trans-mucosal exudation, whereas a value of 0.73 was calculated from our results. There is some destruction of $\alpha_1\text{AT}$ in gastric juice of pH below 3,¹³ but from the results of Crossley and Elliot⁷ it is unlikely that $\alpha_1\text{AT}$ is degraded in the small or large bowel to any significant degree nor is ^{51}Cr absorbed to any appreciable extent.¹⁴ Moreover, the molecular weights of $\alpha_1\text{AT}$ ¹⁵ and albumin are comparable and no significant difference exists in exudative enteropathy in the loss of small or large proteins. Höfer *et al*¹⁶ found that chrominated albumin is present in a monomer fraction and in an aggregated ^{51}Cr -albumin fraction. The metabolic behaviour of the aggregate and monomer fraction is different. The aggregate fraction is stored in the liver in a greater amount than the monomer fraction followed by a smaller intestinal exudation, which could possibly explain the difference between $\alpha_1\text{AT}$ and ^{51}Cr clearance.

Differences in preparing stool such as dilution, homogenisation, or centrifugation, had no influence on the determination of $\alpha_1\text{AT}$. Therefore the contradictory results of Bernier *et al*⁸ and Haeney *et al*⁹ cannot be explained in that way.

Our results confirm those of Bernier *et al*⁸ in that no overlap of $\alpha_1\text{AT}$ clearance was seen in patients and controls, and all patients had raised $\alpha_1\text{AT}$ clearance levels (Fig. 2). If all patients with Crohn's disease lost serum proteins into the intestinal lumen one should expect a quantitative correlation of intestinal protein loss or serum albumin concentrations and extent and/or severity of disease.^{13 17} This was not so in our cases. Even patients with apparent curative resection had raised values while others with a high activity index or those with extensive involvement had relatively minor protein losses (Figs. 3, 4). An explanation for this paradox may be that normal macroscopic appearance of the mucosa does not exclude histological involvement with changes characteristic of Crohn's disease.^{3 18 19} Lymph drainage may be impaired and protein exudation increased if the mesenteric and lymph nodes are involved, features which cannot be diagnosed by radiology. On the other hand quiescent disease may still show impressive

radiological changes.

Florent *et al*¹³ found an exceptionally good correlation between the intestinal $\alpha_1\text{AT}$ clearance and the clinical activity assessed by the CDAI. From our results it is not possible to ascertain any coherence between CDAI and intestinal protein loss. The important objection to the CDAI is, that it is largely determined by subjective variables that may not generally be reliable indicators of the intestinal inflammatory activity but often result from complications or residual effects of the disease. It is obvious that considerable subjective complaints like fibrous strictures, anorectal fistulation, extra-intestinal manifestations, or conglomerate tumour may not necessarily be related to the severity of intestinal inflammation.

Accordingly, in patients with complications the mean CDAI was higher than in patients without complications (Fig. 5). In contrast the mean $\alpha_1\text{AT}$ clearance in patients without complications was raised above the values of patients with complications (Fig. 6). From this comparison it is obvious that in patients suffering from complications the CDAI reflects a much higher clinical activity as it is evident from the intestinal protein loss indicating severity of the local intestinal inflammation.¹⁴

Discrepancies between the quantity of intestinal protein loss and the activity of Crohn's disease estimated by the Best index may be explained by the absence of mucosal parameters as determining factors for this index. The same reason may apply for the six of 14 patients of Florent *et al*¹³ who had raised $\alpha_1\text{AT}$ clearances in spite of a normal Best index and developed a relapse on follow up.

Selection of patients may be another cause explaining differences of the studies. Twenty-eight out of the 30 patients of Florent *et al*¹³ with Crohn's disease had marked hypoalbuminemia (mean serum albumin concentration 2.73 ± 0.53 g%). Only nine out of 26 of our patients had albumin concentrations below 3.6 g% (mean value 3.80 ± 0.57 g%).^{17 20} Furthermore, these authors studied patients without previous intestinal surgery whereas 10 of our patients had undergone intestinal resection and complications were not documented.

Protein loss *via* the intestinal wall seems to be one of the most accurate parameters to measure the inflammatory activity in the intestine.¹⁴ So the determination of $\alpha_1\text{AT}$ by radioimmunodiffusion may be a convenient and reproducible objective method to assess severeness of intestinal inflammation.

Further investigations will show if the $\alpha_1\text{AT}$ clearance is to be of use in following the course of illness or the efficacy of therapy in patients with Crohn's disease.

References

- 1 Best W, Bechtel JM, Singleton JM, Kern J. Development of a Crohn's disease activity index. *Gastroenterology* 1978; **70**: 739–44.
- 2 Andersen SB, Jarnum S. Detection of abnormal gastrointestinal protein loss by means of ⁵⁹Fe-inferon. *Lancet* 1965; **1**: 1060–2.
- 3 Gordon RS. Exudative enteropathy. Abnormal permeability of the gastrointestinal tract demonstrable with PVP. *Lancet* 1959; **1**: 325–6.
- 4 Waldmann TA. Gastrointestinal protein loss demonstrated by ⁵¹Cr-labelled albumin. *Lancet* 1961; **2**: 121–3.
- 5 Waldmann TA, Morell GA, Wochner RD, Strober W, Sternlieb I. Measurement of gastrointestinal protein loss using ceruloplasmin labelled with ⁶⁷copper. *J Clin Invest* 1976; **46**: 10–20.
- 6 Jeejeehoy KN, Singh B, Mani RS, Sanjana SM. The use of ⁹⁵Nb-labelled albumin in the study of gastrointestinal protein loss. *Physiology and pathophysiology in plasma protein metabolism*. Bern: Huber-Verlag, 1965.
- 7 Crossley JR, Elliot RB. Simple method for diagnosing protein losing enteropathies. *Br Med J* 1977; **1**: 428–9.
- 8 Bernier JJ, Florent CH, Desmazes CH, Aymes CH, l'Hirondel CH. Diagnosis of protein-losing enteropathy by gastro-intestinal clearance of alpha₁-antitrypsin. *Lancet* 1978; **2**: 763–4.
- 9 Haeney MR, Fields J, Carter RA, Thompson RA, Asquith P. Is faecal alpha₁-antitrypsin excretion a reliable screening test for protein-losing enteropathy? *Lancet* 1979; **2**: 1161–2.
- 10 Keaney NP, Kelleher J. Faecal excretion of alpha₁-antitrypsin in protein-losing enteropathy. *Lancet* 1980; **1**: 711.
- 11 Schmitt RK, Thews G. *Einführung in die Physiologie des Menschen*. Berlin: Springer Verlag, 1976.
- 12 Florent C, l'Hirondel C, Desmazes C, Aymes C, Bernier JJ. Intestinal clearance of alpha₁-antitrypsin. *Gastroenterology* 1981; **81**: 777–80.
- 13 Florent C, l'Hirondel C, Desmazes C, Giraudeau V, Bernier JJ. Evaluation of ulcerative colitis and Crohn's disease activity by measurement of alpha₁-antitrypsin intestinal clearance. *Gastroenterol Clin Biol* 1981; **5**: 193–7.
- 14 van Tongeren HJ, Eekhout A. Criteria to assess the severity of Crohn's disease and the effect of treatment. In: Weterman IT, Pena AS, Booth CC, eds. *The management of Crohn's disease*. Proceedings of the workshop on Crohn's disease, Leyden 23–25 October 1975. Amsterdam: Elsevier, 1976: 153–8.
- 15 Schulze HE, Heide K, and Haupt H. Alpha₁-Antitrypsin aus Humanserum. *Klin Wochenschr* 1962; **40**: 427–9.
- 16 Höfer R, Schatz H, Thumb N. Untersuchungen über das Stoffwechselverhalten von ⁵¹Cr-Albumin. *Wien Klin Wochenschr* 1968; **31/32**: 587–9.
- 17 Beeken WL, Busch HJ, Silvester DL. Intestinal protein loss in Crohn's disease. *Gastroenterology* 1972; **62**: 207–15.
- 18 Dyer N, Stansfield A, Dawson A. The value of rectal biopsy in the diagnosis of Crohn's disease. *Scand J Gastroenterol* 1970; **5**: 491–6.
- 19 Ruddel W, Lovell D, Blendis L. Bowel mucosa in Crohn's disease. *Lancet* 1976; **1**: 539–40.
- 20 Brassine A. Étude du métabolisme de la sérum-albumine et mesure des pertes digestives des protéines plasmatiques dans la maladie de Crohn et la rectocolite hémorragique. *Acta Gastroenterol Belg* 1978; **41**: 7–17.
- 21 Goodman M, Skinner J, Truelove S. Abnormalities in the apparently normal bowel mucosa in Crohn's disease. *Lancet* 1976; **1**: 275–6.