Correspondence

Combined cow's milk protein and gluten induced enteropathy

Sir, – Watt *et al* (*Gut* 1983; **24**: 165–70) present a very interesting patient with combined cow's milk protein and gluten induced enteropathy. We have been following a very similar case for eight years, which has already been reported in a local publication.¹

The patient, a girl, was referred to the hospital at 10 months of age, with symptoms of recurrent diarrhoea and low weight gain since the neonatal period. She had received anti-diarrhoeal drugs in an irregular way.

At 6 months of age she showed poor appetite and failure to thrive, and at 9 months abdominal distention and restlessness became evident. She had been fed a cow's milk formula since birth and gluten was introduced into the diet at the age of 2½ months. On admission a malabsorption syndrome was apparent and a jejunal biopsy showed a subtotal villous atrophy. Suspecting the coexistence of cow's milk protein and gluten induced enteropathy, as described by Visakorpi and Immonen,² a cow's milk free and gluten free diet was introduced. The patient became asymptomatic and thrived well. The jejunal biopsy performed 10 months later showed partial villous atrophy.

Cow's milk protein was reintroduced at the age of 23 months and four months later a jejunal biopsy still exhibited a subtotal villous atrophy. Suspecting that the patient's diet was not being adhered to we again emphasised to the family the importance of a strict gluten free diet. Five months later, however, the histological appearance of the jejunal mucosa remained unchanged, despite the mother's assurance that no gluten had been administered. Withdrawal of cow's milk produced a completely normal jejunal biopsy 19 months later. Four months later a new diet containing cow's milk again caused severe partial villous atrophy of the intestine.

To clarify whether the enteropathy was induced by cow's milk protein or by a lax gluten free diet, at the age of 5½ years a cow's milk challenge was carried out, taking jejunal biopsies both immediately before, and 24 hours after ingestion according to methodology previously described.³ We found that the jejunal biopsy changed from a normal histological picture to a severe partial villous atrophy. Milk was therefore withdrawn from the diet and six months later we confirmed histologically that the intestinal mucosa had returned to normal.

At this time a gluten challenge was carried out, producing six months later severe partial villous atrophy of the intestine. In view of these results we diagnosed combined cow's milk protein and gluten induced enteropathy. As in the case of Watt and his colleagues, our patient remained completely asymptomatic during all the dietary manipulations.

At 8½ years of age a similar cow's milk challenge was repeated and a normal jejunal biopsy was then observed both before and after the provocation test. On the basis of this finding we consider that the milk induced enteropathy had been of transient nature.

In our experience, the frequency of asymptomatic cow's milk induced enteropathy is 33%, as previously reported,³ and corroborated by a further, wider range study. We did not find evidence of an allergic origin for this enteropathy, so we believe it should be differentiated from cow's milk allergy.

Finally, we too raise the question of whether this enteropathy might produce long term sequelae, and for this reason we recommend a very careful follow up of these patients.

J C VITORIA,
A SOJO, AND
C CAMARERO

*Department of Pediatrics,
Hospital Infantil de la Seguridad
Social Cruces, Bilbao,
Spain.*

References

- 1 Sojo A, Cerezo JM, Martinez-Ayucar M, López C, Abeijón JA, Fernandez de Pinedo R, Vitoria JC. Enteropatía inducida por gluten. *Bol S Vasco-Nav Ped* 1980; **15**: 119–34.
- 2 Visakorpi JK, Immonen P. Intolerance to cow's milk and wheat gluten in the primary malabsorption syndrome in infancy. *Acta Paediatr Scand* 1967; **56**: 49–56.
- 3 Vitoria JC, Camarero C, Solaguren R, Aranjuelo M, Oliveros R, Navajas A, Rodriguez-Soriano J. Cow's milk protein-sensitive enteropathy. Clinical and histological results of the cow's milk provocation test. *Helv Paediatr Acta* 1979; **34**: 309–18.

Reply

Sir, – We were most interested to read the letter of Vitoria *et al*, and are grateful for the opportunity to reply. They describe a case of combined cow's milk protein and gluten induced enteropathy similar to that which we reported in *Gut* 1983; **24**: 165–70.

At first sight the major difference between the

cases appears to be that in their case, the cow's milk protein induced element of the enteropathy resolved at the age of 8½ years, while our patient was still intolerant of both proteins at the time of publication. This situation, however, has changed in the last week.

In August 1982, the child was clinically well and had a normal jejunal biopsy while on a gluten free and cow's milk free diet. Cow's milk only was reintroduced at this time and after a symptom free year a further jejunal biopsy was performed (Figure). This showed a normal villous pattern and normal surface epithelium though there was still a

Figure Jejunal biopsy, PAS stained $\times 125$ (original magnification). The biopsy appearance has returned to almost complete normality on a gluten free, cow's milk protein containing diet at the age of 8½ years.

mild increase in the number of lymphocytes and plasma cells in the lamina propria. Although the biopsy is not completely normal there is a good reason to expect that histological normality will be achieved, and a further biopsy will be undertaken to confirm this. Our patient is now 8½ years of age.

It is of interest that in both these cases the cow's milk protein intolerance resolved, but this resolution was, at least histologically, delayed to the age of 8½ years. We feel sure that some of the other interesting questions posed by these two cases will only be answered when our immunological colleagues have finally elucidated the mechanisms involved in intolerance to dietary proteins.

J R PINCOTT
AND J WATT

*The Hospital for Sick Children,
Great Ormond Street,
London WC1N 3JH.*

Factors affecting mortality and morbidity after surgery for obstructive jaundice

Sir,
Following the recent publication of our paper in *Gut* 1983; 24: 845–52, we have had a number of enquiries requesting further information on the relationship of age and mortality. We reported that patients over 60 years of age had a significantly higher hospital mortality than those below 60 but that age was not of independent significance when other factors, such as bilirubin, haematocrit, and malignancy were taken into account. The complete data relating age and mortality (Table) show that in our series, hospital mortality does not increase significantly in any decade in patients over 60 years of age. If one looks at other conditions where one would expect

Table Relationship of age and hospital mortality for 373 patients undergoing surgery for obstructive jaundice

Age (years)	Patients (no)	Patients (no)	Hospital mortality* (%)
<50	66	0	0
50–59	55	3	5.5
60–69	96	11	11.5
70–79	123	15	12.2
80+	33	5	15.2
Overall group	373	34	9.1

* Hospital mortality – death within 30 days of operation or during the same hospital admission.

age to be a significant factor, such as pulmonary embolism, it has been shown from epidemiology surveys that the apparent increase in frequency with age is likely to be due to an increase in associated conditions such as malignancy and heart disease,¹ a conclusion similar to that reached for age and hospital mortality in our series.

It is thus our conclusion that age alone should not be a discriminant in the selection of patients for surgical treatment of their obstructive jaundice.

J MICHAEL DIXON,
C P ARMSTRONG
S W DUFFY
R A ELTON, AND
G C DAVIES

University Departments of Clinical Surgery, Royal Infirmary, Edinburgh Medical Computing and Statistics Units, University of Edinburgh

Reference

- 1 Coon WW, Collier FA. Some epidemiologic considerations of thromboembolism: *Surg Gynecol Obstet* 1959; 109: 487–97.