References

- 1 Hamilton I, Cobden I, Axon ATR. *In vitro* determination of small intestinal permeability. *Gut* 1984; **25**: 909–10.
- 2 Menzies IS. Transmucosal passage of inert molecules in health and disease. In: Skadhauge E, Heintze K, eds. *Intestinal absorption and secretion*. Falk Symposium 36 Lancaster: MTP Press, 1983: 527–43.
- 3 Menzies IS, Laker MF, Pounder R, Bull J, Heyer S, Wheeler PG, Creamer B. Abnormal intestinal permeability to sugars in villous atrophy. *Lancet* 1979; **ii**: 1107–9.
- 4 Jenkins RT, Lock CJL, Rooney PJ. ⁵¹Cr-EDTA in studies of intestinal permeability. *Lancet* 1984; **ii**: 1342.
- 5 Schutz SG, Solomon AK. Determination of effective hydrodynamic radii of small molecules by viscometry. *J Gen Physiol* 1961; **44**: 1189–99.
- 6 Killan RCG, Lawrence JL, Sharma VC. The refinement of the crystal structure of α -D-glucopyranose monohydrate. *Acta Cryst* 1971; **B27**: 1707–10.
- 7 Berman HM, Jeffrey GA, Rosenstein RD. The crystal structures of the α and β forms of D-mannitol. *Acta Cryst* 1968; **B24**: 442–49.
- 8 Wood RA, Jeffrey GA. The crystal structure and solid state NMR analysis of lactulose. *Am Cryst Assoc Newsletter* 1982; **10**: 18, and Allen FH, Bellard S, Brice MD *et al*. The Cambridge crystallographic data centre: computer-based search, retrieval, analysis and display of information. *Acta Cryst* 1979; **B35**: 2331–39.
- 9 Gerdorf LE, Baenziger NA, Goff HM. Crystal and molecular structure of a substitution-labile chromium (III) complex: aquo(ethylenediamine-triacetatoacetic acid)chromium(III). *Inorg Chem* 1981; **00**: 1606–8.
- 10 Ecknauer R, Buck B, Breitig D. An experimental model for measuring intestinal permeability. *Digestion* 1983; **26**: 24–32.
- 11 Loehry CA, Axon ATR, Hilton PJ, Hider RC, Creamer B. Permeability of the small intestine to substances of different molecular weight. *Gut* 1970; **11**: 466–70.

Peptic ulcer and non-steroidal anti-inflammatory agents

SIR,—I was most interested in the recent case control study by Duggan *et al* (*Gut* 1986; **27**: 929–33) confirming an association of non-steroidal anti-inflammatory agents (NSAIDs) with gastric ulcer in Australia. I note that no less than 42% of the cases presented with haemorrhage, and wonder whether comparison of this group with the remainder would add anything to the conclusions. British studies have shown a significant association of NSAIDs with haemorrhage from gastric ulcer in the elderly^{1–3} and also with ulcer perforations.⁴ Although it might seem logical that if an agent causes ulcers to bleed and perforate it could also cause an ulcer in the first place, this problem is far from simple – especially as

we do not know how gastric ulcers begin. If, for example, they can arise from gastric erosions, why are they usually single and why do they not follow mucosal biopsies?

If the authors found that NSAID ingestion was related to symptomatic non-bleeding ulcers, this would be valuable information. Unfortunately it still would not solve the question of cause, as so many ulcers are symptomless and/or undiagnosed,^{5,6} especially in the elderly⁷ who are the chief users of these agents.

R D MONTGOMERY

*Department of Medicine,
East Birmingham Hospital,
Birmingham B9 5ST.*

References

- 1 Thompson MR. The unnatural history of drug-associated gastric ulcers. [Abstract]. *Gut* 1981; **22**: A877.
- 2 Jolobe OMP, Montgomery RD. Changing clinical pattern of gastric ulcer: are anti-inflammatory drugs involved? *Digestion* 1984; **29**: 164–170.
- 3 Somerville K, Faulkner G, Langman M. Non-steroidal anti-inflammatory drugs and bleeding peptic ulcer. *Lancet* 1986; **i**: 462–64.
- 4 Collier D St J, Pain JA. Non-steroidal anti-inflammatory drugs and peptic ulcer perforation. *Gut* 1985; **26**: 359–63.
- 5 Jorde R, Bostad L, Burhol PG. Asymptomatic gastric ulcer: a follow-up study in patients with previous gastric ulcer disease. *Lancet* 1986; **i**: 119–20.
- 6 Bank J, Wright JP, Lucke W, Marks IN. Peptic ulcer: a follow-up study. *J Clin Gastroenterol* 1986; **8**: 381–84.
- 7 Clinch D, Banerjee AK, Ostick G. Absence of pain in elderly patients with peptic ulcer. *Age Ageing* 1984; **13**: 120–23.

Reply

SIR,—We thank Dr Montgomery for his comments about a possible difference between bleeding and non-bleeding ulcer. For bleeding ulcers there were nine pairs in which one member took NSAIDs regularly and the other none at all; seven were patients and two controls giving an odds ratio of 3.5. In the nine non-bleeding such pairs, eight were patients and one was a control (OR=8).

J M DUGGAN

*Department of Gastroenterology,
Royal Newcastle Hospital,
Newcastle NSW 2300,
Australia.*