Leading article

Pathogenesis of chronic pancreatitis

There have been few efforts to classify diseases of the pancreas by their specific pathological features and causes. In 1946 Comfort *et al*¹ observed that the changes of acute pancreatitis were often found in resected specimens of the pancreas from patients with chronic pancreatitis, suggesting that recurrent acute attacks were the basis for the development of chronic pancreatitis. It could be, however, that acute lesions complicated chronic lesions.

The hypothesis of Comfort *et al* was widely accepted and extended. Cattell and Warren² suggested that all possible transitions exist between the most acute haemorrhagic pancreatitis and the most chronic calcified pancreatitis, which represented two extremes of one disease: 'recurrent pancreatitis.'

Because pancreatic biopsy specimens are not readily available most clinicians attempted to classify their cases according to clinical or radiological criteria. The most recent attempt was the Cambridge classification³ which defined changes in the pancreatic ducts as seen with endoscopic retrograde pancreatography. The Marseille classifications⁴⁻⁶ represent an effort to isolate stable groups of pathological features and to study the statistical relationship between them and pathogenesis. The 1963 classification proposed that acute pancreatitis was not a common cause of chronic pancreatitis since the statistical studies showed that the average age of patients at first presentation with acute pancreatitis (the putative cause of recurrent pancreatitis) was 13 years more than the mean age at onset of chronic calcified pancreatitis. Further, the two diseases were statistically linked to different causes.^{7,8} Recent experiments have shown that acute haemorrhagic pancreatitis after injection of trypsin⁹⁻¹¹ or after hyperstimulation^{12,13} resolves after a phase of destruction and fibrosis of the exocrine parenchyma. These changes resemble the lesions of chronic pancreatitis but are reversible. Follow up of patients presenting with haemorrhagic necrosis by computed tomography has shown that even the most severe and extensive lesions will regress if the patient survives.¹⁴

The recent Marseille-Rome classification¹⁵⁻¹⁶ defined acute pancreatitis as 'a spectrum of inflammatory lesions in the pancreas and also in peripancreatic tissues: oedema, necrosis, haemorrhagic necrosis, fat necrosis.' Chronic pancreatitis was defined as 'the presence of chronic inflammatory lesions characterised by the destruction of exocrine parenchyma and fibrosis and at least in the later stages, the destruction of endocrine parenchyma.' It is frequently complicated in the early stages of its evolution by attacks of acute pancreatitis which are responsible for recurrent pain and which may represent the only clinical symptom. After some years there is insufficiency, both exocrine (steatorrhoea) and endocrine (diabetes), and acute attacks decrease and disappear.

Two large computer based studies of chronic pancreatitis^{17,18} distinguished two different types of chronic pancreatitis. A three dimensional reconstruction of the pancreatic ductal system using 7 µm sections drew the same conclusions.¹⁹ The two populations defined by their pathological features were statistically linked to different and specific causes.¹⁷

In the first and smaller group chronic pancreatitis was due to obstruction of the pancreatic ducts predating the

pancreatitis: tumours, papillary stenosis, cysts, or scars due to acute pancreatitis or trauma are the main causes. 'The lesions distal to the site of ductal obstruction are uniform, not patchy. The epithelium of the ducts in the obstructed part of the gland is usually preserved. Intraductal plugs may be observed but they are less frequent than in the following form and calcified plugs or pancreatic stones (calcifications) are not found.'

The usual form of chronic pancreatitis may be called chronic calcifying pancreatitis. Three studies have shown that after a variable time most chronic pancreatitis patients present with pancreatic calcifications visible on radiological films of the abdomen.²⁰⁻²² Calcification is intraductal pancreatic calculi.^{23,24} Pancreatic calculi are composed of two materials: calcium carbonate, generally in the form of calcite which constitutes approximately 95% of stone weight²⁵ and a curious protein called PSP.²⁶ Radiolucent pancreatic stones composed of different degraded forms of PSP occur additionally.²⁷ We have described one example of a patient presenting with large pancreatic stones which she had had since childhood. These stones consisted of pure calcium carbonate without PSP.²⁸

A morphometric ultrastructural study of exocrine pancreas of patients with chronic calcifying pancreatitis²⁹ was undertaken in those areas of the gland which were normal on light microscopy. The lobular distribution of the disease means that there are frequently such normal areas in a pathological pancreas. This study showed that in these areas acinar and ductal cells were intact but, compared to normal acinar cells, acinar cells from patients with chronic pancreatitis showed signs of increased protein secretion (larger diameter of the cell, nucleus, and nucleolus; increased length of endoplasmic reticulum; increased number of condensing vacuoles; decreased number of mature zymogen granules). Despite the normality of the cells, intra-acinar, or more frequently intraductal eosinophilic protein precipitates or plugs (which had been previously described at the optical microscope level) were found.^{15,21} Therefore, even in its earliest stages, before there is any apparent damage to the cell, chronic calcifying pancreatitis seems to be a lithiasis.

These precipitates are mostly composed of protein fibrils and many transitions between plugs and calculi have been described.^{30,31} Similar protein plugs are frequently observed in non-activated pure pancreatic juice collected from patients with chronic calcific pancreatitis. They comprise a degraded molecular form of PSP.³² During the early stage of the disease pure pancreatic juice in addition to plugs contains crystals of calcium carbonate which are larger and more numerous than in normal subjects.³³

Pancreatic lithogenesis seems to comprise precipitation of calcium salts and precipitation of different degraded forms of a special protein (PSP).

The original PSP molecules are found in pancreatic juice where they are termed PSP-S2-5. These molecules are glycosylated with a molecular mass of 16 000 to 19 000 daltons. They have the same 144 amino acid backbone, the different molecular masses being due to different glycosylation. The amino acid sequence of PSP and of its signal peptide and the nucleotide sequence of PSP mRNA are known.³⁴ This sequence does not resemble any other known

protein sequence and in particular it is completely different from trypsinogen and chymotrypsinogen.³⁵ PSP S2-5 is hydrolysed by trypsin but not by chymotrypsin to a long C-terminal 133 amino acid peptide which is not glycosylated, is insoluble, and polymerises to form fibrils. This has been called PSP-S1. A short N-terminal 11 amino acid peptide which is glycosylated and soluble is also formed.³⁶ PSP-S1 is easily formed in pure pancreatic juice when trypsin activation is not prevented by antiproteases. A similar fibrillar protein has been discovered in bovine³⁷ and human³⁸ pancreatic tissue by Gross *et al* with the name PTP. The partial amino acid sequence given by these authors strongly suggests that PTP is similar to PSP-S1, the degraded fragment of PSP-S2-5.

PSP-S is synthesised by the pancreatic acinar cell,³⁹ stored in the zymogen granules in the same way as enzymes,⁴⁰ and secreted in parallel with them.⁴¹ PSP can prevent the nucleation and crystal growth of calcium carbonate in a saturated solution,^{42,43} particularly PSP-S2-5 and the N-terminal glycosylated 11 amino acid peptide, but not in PSP-S1.

Pancreatic juice is normally saturated in calcium both in the dog⁴⁴ and in humans.⁴⁵ The observation that precipitation of calcium salts is one of the two phenomena observed in pancreatic lithogenesis and PSP prevents calcium precipitation in saturated solutions suggests that a deficiency of PSP in pancreatic secretion could play an important part in the disease. This is likely since the messenger RNA responsible for PSP biosynthesis is significantly decreased in the pancreas of patients with chronic calcifying pancreatitis (whether alcoholic or non-alcoholic),³⁹ PSP is severely decreased in the zymogen granules of acinar cells from these patients compared to normal subjects,⁴⁰ and PSP concentration related to total protein is significantly decreased in the pure pancreatic juice of these patients.^{41,46} PSP estimation in pancreatic juice requires precise experimental conditions. When the juice is activated or when it is stored hydrolysis of the molecule produces the insoluble residue PSP S-1 which polymerises and precipitates. This probably explains why there is no clear distinction between patients and controls when juice is frozen and stored or when PSP is estimated with a monoclonal antibody prepared in Marseille and raised against PSP-S1.^{46,47} However, with polyclonal antibodies, using immunodiffusion⁴¹ or enzyme-linked immunosorbent assay,⁴⁶ the concentration of PSP related to secretory protein is decreased with almost no overlap between chronic calcifying pancreatitis patients and normal non-alcoholic controls.

In apparently normal alcoholic controls (mean daily consumption >100 g per day) PSP is decreased by half. This could be due either to an alcohol induced decrease of PSP biosynthesis or to increased denaturation in the acinar or duct lumen, or both.

Low PSP concentrations in pancreatic juice probably have at least two causes: one is independent of alcohol consumption since it is found in hereditary and idiopathic pancreatitis, the other is related to alcoholism.

The explanation for the precipitation of residues of PSP is more difficult to explain. The fibrillar protein present in pancreatic plugs is the degraded polymerised form PSP-S1. This suggests that a hydrolysis of PSP-S takes place in pancreatic juice in the earliest stages of chronic calcifying pancreatitis and as high as the acinar lumen (plugs are observed in the acinar lumen but more frequently in the ducts).⁴⁸ It is just possible, and has not been completely excluded, that an active protease which has the same specificity as trypsin, either activated trypsin itself or cathepsin B, is present in the lumen of the acinus. A careful morphometric study did not show any modification of lysosomes.²⁹ Spontaneous degradation of PSP-S2-5 is another possibility. Protein plugs are observed in 2% of normal pancreatic juices and are significantly more frequent in the

samples collected from normal alcoholics and are found in more than 60% of patients with chronic calcifying pancreatitis, either alcoholic or abstinent³²; plug formation is therefore increased by alcohol consumption, but this is not the only factor.

Plug formation by itself is not a sufficient condition for pancreatic stone formation. It is observed in normal and alcoholic controls but it seems to be responsible for some duct lesions: in contact with plugs the basal membrane of ducts vanishes,⁴⁹ which is responsible for the observed transudation of serum protein⁵⁰ and calcium^{51,52} into pancreatic juice. This increased calcium concentration must facilitate calcium crystal formation and explains why calcifications visible on x ray films develop more rapidly after many years of clinical pancreatitis.²⁰⁻²² Spontaneous plug formation in normal juice could explain the frequent observation of tiny stones and duct lesions in localised areas of the pancreas at necropsy.^{53,54} In chronic calcifying pancreatitis duct lesions are more severe. Atrophy of the epithelium followed by stenosis, cyst formation, and parenchymal atrophy distal to obstructed ducts are apparently the final consequences of plugs and stones.¹⁷⁻¹⁹

Chronic calcifying pancreatitis is significantly associated with alcohol consumption, fat, a high protein diet, tobacco smoking, hypercalcaemia, and life in some tropical countries where malnutrition is common. There are also inherited cases.^{55,56}

Ethanol is metabolised by the pancreas⁵⁷ and it has been suggested that a direct toxic effect on the pancreas could explain alcoholic chronic calcifying pancreatitis.⁵⁸ But these lesions are an increased content of triglycerides and lesions of lysosomes in the acinar cell. They are observed in all organs of alcoholics and are reversible. They do not explain the formation of plugs and stones.⁵⁹

Two alcohol induced secretory modifications probably increase calcium crystal formation: a decreased secretion of citrate⁶⁰ and PSP.⁴⁶ This could be due to alcohol metabolism or to adaptation to alcohol.

The concentration and output of secretory protein is increased in the juice of alcoholic dogs and humans. This could play a part by increasing the viscosity of the juice. The same finding is observed in the juice of hypercalcaemic dogs, cats, and humans. As hypercalcaemia is also a cause of chronic calcifying pancreatitis, it could play a part, possibly by increasing the resistance to flow. The mechanism of this increased secretion is neural, cholinergic, and vagally mediated in alcoholics and neural and hormonal in hypercalcaemics (for review see Sarles⁵⁵ and Sarles *et al*⁵⁶).

The cause of tropical pancreatitis is not known. It is seen in non-alcoholic children and young adults of both sexes living in some tropical areas such as South India, Indonesia, Zaire, and Nigeria but not in others (for review see Sarles *et al*⁵⁶ and Mohan *et al*⁶¹). Kwashiorkor⁶² and cassava (manioc) consumption⁶³ have been implicated. Studies by Indian physicians⁶¹ and ourselves,⁶⁴ conducted in different areas and comparing the type of diet and the incidence of tropical pancreatitis, have shown that neither factor plays a part. The nutritional disorders linked to the disease are a low protein diet (50 g per day or less) and a very low fat intake (less than 30 g per day).^{56,64} It has previously been shown that a low as well as a high fat intake is a risk factor for chronic calcifying pancreatitis.⁶⁵ These disorders, if they have a role, could act directly on the pancreas of the child, but the disease begins early in life in Kerala, for instance, where restricted diets are common in adults as well as in children.⁶⁶ This is not the case in countries such as the Ivory Coast where malnutrition is mostly found during childhood.^{64,67} Malnutrition of the mother may play a part in the disease of the child. We have shown an increased enzyme and particularly an increased protease content in the pancreas of young rats born of animals

submitted to a moderately restricted protein diet (9% of calories). This modification persists several months after weaning.⁶⁸ In any case the pathological features of tropical pancreatitis are identical to those of alcoholic and idiopathic pancreatitis in Western countries. The biochemical composition of stones is similar.⁶⁹

Hereditary factors certainly play a part in the disease. There is a hereditary dominant form observed in children of both sexes.⁷⁰ The only known biochemical modification is a decrease of PSP in pancreatic juice.⁴¹ However, there is an increased risk that in the family of a patient with chronic calcifying pancreatitis there will be another case of chronic calcifying pancreatitis when compared with normal control subjects,⁷¹ which suggests that there is an underlying defect, probably different from the hereditary dominant pancreatitis, sometimes latent, sometimes shown by nutritional disorders, and sometimes sufficient to produce the disease by itself (idiopathic cases). The increased frequency of blood group O in chronic calcifying pancreatitis patients⁷² is consistent with this hypothesis. The many reports dealing with HLA markers are contradictory and do not provide consistent results.⁵⁵

Thus, although most cases of chronic calcifying pancreatitis in Western countries are related to consumption of alcohol, other rare causes exist (hypercalcaemia and heredity). In some tropical countries a form of chronic calcifying pancreatitis occurs which could be related to protein and fat malnutrition. All these forms are characterised by identical pathological changes in the gland and biochemical changes in the pancreatic secretion. When stones are present they have a similar composition independent of the cause of the chronic calcifying pancreatitis. We therefore have to conclude that chronic pancreatitis is due to stone formation consequent on changes in the composition of pancreatic juice.

The authors are grateful to Dr C D Johnson for correcting the style of this manuscript.

H SARLES, J P BERNARD, L GULLO

Clinique des maladies de l'Appareil Digestif et de la Nutrition –
Hôpital Sainte-Marguerite, Marseille, France
Unité de Recherches de Physiologie et Pathologie Digestives U 315, 46, Marseille, France

Correspondence to: Professeur Henri Sarles, Clinique des maladies de l'Appareil Digestif et de la Nutrition, Hôpital Sainte-Marguerite, BP29, 13277 Marseille Cedex 9, France.

- Comfort HW, Gambill EE, Baggenstoss AH. Chronic relapsing pancreatitis: a study of 29 cases without associated disease of the biliary or gastrointestinal tract. *Gastroenterology* 1946; 6: 239–85.
- Cattel RB, Warren KW. *Surgery of the pancreas*. London: WB Saunders, 1973.
- Sarner M, Cotton PB. Classification of pancreatitis. *Gut* 1984; 25: 756–9.
- Sarles H. *Pancreatitis. Symposium in Marseille* 1963. Basel: Karger, 1965.
- Singer MV, Gyr K, Sarles H. Revised classification of pancreatitis. Report of the second international symposium on the classification of pancreatitis in Marseille, France, March 28–30, 1984. *Gastroenterology* 1985; 89: 683–90.
- Sarles H. Classification et définition des pancréatites. Marseille-Rome 1988. *Gastroenterol Clin Biol* 1989; 13: 857–9.
- Sarles H, Sarles JC, Camatte R, et al. Observations on 205 confirmed cases of acute pancreatitis, recurring pancreatitis and chronic pancreatitis. *Gut* 1965; 6: 545–59.
- Howard JM, Jordan G. *Surgical diseases of the pancreas*. Philadelphia: Lippincott, 1960.
- Odaira C, Berger Z, Iovanna JL, Sarles H. Localized necrohemorrhagic pancreatitis in the rat after pancreatic interstitial trypsin injection. Regressive pseudochronic lesions. *Digestion* 1986; 34: 68–77.
- Lechene De La Porte P, Iovanna JL, Odaira C, Choux R, Sarles H, Berger Z. Tubular complexes: one of the sources of pancreatic exocrine parenchyma after acute necrohemorrhagic pancreatitis. *Pancreas* (in press).
- Iovanna JL, Odaira C, Berger Z, Sarles H. Temporary pseudochronic lesions during the recovery of acute necrohemorrhagic pancreatitis in rabbits. *Pancreas* 1983; 3: 433–8.
- Adler G, Hupp T, Kern HF. Course and spontaneous regression of acute pancreatitis in the rat. *Virchows Arch (A)* 1979; 382: 31–47.
- Elsasser HP, Adler G, Kern HF. Time course and cellular source of pancreatic regeneration following acute pancreatitis in the rat. *Pancreas* 1986; 1: 421–9.
- Delcenserie R, Tossou H, Jolly JP, Delamarre J, Deschepper B, Dupas JL. Interet pronostique de l'angiomodensitometrie pancreatique precoce dans les pancreatites aiguës. 24^e Forum de Recherches en Hepatogastroenterologie, Geneve, Avril 1989. *Gastroenterol Clin Biol* 1989; 13: A139.
- Sarles H, Adler G, Dani R, et al. The pancreatitis classification of Marcellles-Rome 1988. *Scand J Gastroenterol* 1989; 24: 641–2.
- Sarles H, Adler G, Dani R, et al. Classification of pancreatitis and definition of pancreatic diseases. *Digestion* 1989; 43: 234–6.
- Payan H, Sarles H, Demirdjian M, Gauthier AP, Cros RC, Durbec JP. Study of the histological features of chronic pancreatitis by correspondence analysis. Identification of chronic calcifying pancreatitis as an entity. *Biomedicine* 1972; 18: 663–70.
- Sahel J, Cros C, Durbec JP, et al. Multicenter pathological study of chronic pancreatitis. morphological regional variations and differences between chronic calcifying pancreatitis and obstructive pancreatitis. *Pancreas* 1986; 1: 471–7.
- Nakamura K, Sarles H, Payan H. Three-dimensional reconstruction of the pancreatic ducts in chronic pancreatitis. *Gastroenterology* 1972; 62: 942–9.
- Bernades P, Belghiti J, Athouel M, Mallardo N, Breil P, Fekete F. Histoire naturelle de la pancreatite chronique. Etude de 120 cas. *Gastroenterol Clin Biol* 1983; 7: 8–13.
- Ammann RW, Akovbiantz A, Largiader F, Schueler G. Course and outcome of chronic pancreatitis. Longitudinal study of a mixed medical-surgical series of 245 patients. *Gastroenterology* 1984; 86: 820–8.
- Ink O, Labayle D, Buffet C, Chaput JC, Etienne JP. Pancreatite chronique alcoolique: relations de la douleur avec le sevrage et la chirurgie pancreatique. *Gastroenterol Clin Biol* 1984; 8: 419–25.
- Gruber GB. Bauschspeicheldruse. In: Hencke F, Lubarsch D, eds. *Handbuch der Speziellen Pathologischen Anatomie und Histologie*. Berlin: Springer, 1929: 211–621.
- Edmondson HA, Bullock WK, Mehl JW. Chronic pancreatitis and lithiasis. Pathology and pathogenesis of pancreatic lithiasis. *Am J Pathol* 1950; 26: 37–55.
- De Caro A, Multigner L, Vernie H. Identification of two major proteins of bovine pancreatic stones as immunoreactive forms of trypsinogens. *Biochem J* 1982; 205: 543–9.
- Multigner L, De Caro A, Lombardo D, Campese D, Sarles H. Pancreatic stone protein. A phosphoprotein which inhibits calcium carbonate precipitation from human pancreatic juice. *Biochem Biophys Res Commun* 1983; 110: 69–74.
- Multigner L, Mariani A, Daudon M, Schmigel WH, Sarles H, De Caro A. Pancreatic stone protein and radiolucent pancreatic stones. Club Europ. Pancreas, Marseille, 1987. *Digestion* 1987; 38: 44.
- Sarles H, De Caro A, Multigner L, Martin E. Giant pancreatic stones in teetotal women due to absence of the 'stone protein'? *Lancet* 1982; ii: 714–5.
- Tasso F, Stemmelin N, Sarles H, Clop J. Comparative morphometric study of the human pancreas in its normal state and in primary chronic calcifying pancreatitis. *Biomedicine* 1973; 18: 134–44.
- Rumessen JJ, Marner B, Thorsgaard Pedersen N, Permin H. Autoantibodies in chronic pancreatitis. *Scand J Gastroenterol* 1985; 20: 966–70.
- Harada H, Takeda M, Tanaka J, Mini H, Ochi K, Kimura I. The fine structure of pancreatic stones as shown by scanning electron microscopy and x-ray probe microanalyser. *Gastroenterol Jpn* 1983; 18: 530–7.
- Guy O, Robles-Diaz G, Adrich Z, Sahel J, Sarles H. Protein content of precipitates present in pancreatic juice of alcoholic subjects and patients with chronic calcifying pancreatitis. *Gastroenterology* 1983; 84: 102–7.
- Multigner L, Mariani A, Sahel J, De Caro A, Sarles H. Occurrence of calcium carbonate (calcite) crystals in normal and pathological pancreatic juice. Club Europ. Pancreas, Marseille 1987. *Digestion* 1987; 38: 43.
- Giorgi D, Bernard JP, Rouquier S, Iovanna J, Sarles H, Dagorn JC. Secretory pancreatic stone protein messenger RNA. Nucleotide sequence and expression in chronic calcifying pancreatitis. *J Clin Invest* 1989; 84: 100–6.
- Montalto G, Bonicelli J, Multigner L, Rovey M, Sarles H, De Caro A. Partial amino-acid sequence of human pancreatic stone protein. A novel pancreatic secretory protein. *Biochem J* 1986; 238: 227–32.
- De Caro JD, Rouimi P, Rovey M. Hydrolysis of p-nitrophenyl acetate by the peptide chain fragment (336–449) of porcine pancreatic lipase. *Eur J Biochem* 1986; 158: 601–7.
- Gross J, Brauer AW, Bringham RF, Corbett C, Margolies MN. An unusual bovine pancreatic protein exhibiting pH-dependent globule-fibril transformation and unique amino acid sequence. *Proc Natl Acad Sci USA* 1985; 82: 5627–31.
- Gross J, Carison RI, Brauer AW, Margolies MN, Warshaw AL, Wands JR. Isolation, characterization and distribution of an unusual pancreatic human secretory protein. *J Clin Invest* 1985; 76: 2215–26.
- Giorgi D, Bernard JP, De Caro A, et al. Pancreatic stone protein. I. Evidence that it is encoded by a pancreatic messenger ribonucleic acid. *Gastroenterology* 1985; 89: 381–6.
- Lechene De La Porte P, De Caro A, Lafont H, Sarles H. Immunocytochemical localization of pancreatic stone protein in the human digestive tract. *Pancreas* 1986; 1: 301–8.
- Multigner L, Sarles H, Lombardo D, De Caro A. Pancreatic stone protein. II. Implication in stone formation during the course of chronic calcifying pancreatitis. *Gastroenterology* 1985; 89: 387–91.
- Multigner L, De Caro A, Lombardo D, Campese D, Sarles H. Pancreatic stone protein. A phosphoprotein which inhibits calcium carbonate precipitation from human pancreatic juice. *Biochem Biophys Res Commun* 1983; 110: 69–74.
- Multigner L, De Caro A. Kinetic studies on calcium carbonate crystal growth inhibition by human pancreatic stone protein. Club Europ. Pancreas, Marseille, 1987. *Digestion* 1987; 38: 43.
- Moore EW, Verine HJ. Pancreatic calcification and stone formation: a thermodynamic model of calcium in pancreatic juice. *Am J Physiol* 1987; 252: 707–18.
- Gerolami A, Marteau C, Matteo A, et al. Calcium carbonate saturation in human pancreatic juice: possible role of ductal H⁺ secretion. *Gastroenterology* 1989; 96: 881–4.
- Provansal Cheylan M, Bernard JP, Mariani A, Sarles H, Dupuy P. Pancreatic stone protein: quantification in pancreatic juice by enzyme linked immunosorbent assay and comparison with other methods. *Pancreas* 1990; 4: 680–9.
- Schmigel WH, Burchert M, Kalthoff H, et al. Pancreatic stone protein in serum of patients with pancreatitis. *Lancet* 1986; ii: 686–7.
- Sarles H, Muratore R, Sarles JC. Etude anatomique des pancreatites chroniques de l'adulte. *Semin Hop Paris* 1961; 25: 1507–22.
- Uscanga L, Kennedy RH, Choux R, Druguet M, Grimaud JA, Sarles H. Sequential connective matrix changes in experimental acute pancreatitis. An immunohistochemical and biochemical assessment in the rat. *Int J Pancreatol* 1987; 2: 33–45.
- Clemente F, Ribeiro T, Figarella C, Sarles H. Albumine IgA et IgG dans le suc pancreatique humain normal chez l'adulte. *Clin Chim Acta* 1971; 33: 317–24.
- Gullo L, Sarles H, Mott CB, Tiscornia OM, Pauli AM, Pastor J. Pancreatic secretion of calcium in healthy subjects and various diseases of the pancreas. *Rendiconti Gastroenterol* 1974; 6: 35–44.

- 52 Regan PT, Malagelada JR, Dimagno EP. Duodenal calcium outputs in health and pancreatic disease. *Gut* 1980; 21: 614-8.
- 53 Birnstingl M. A study of pancreatography. *Br J Surg* 1959; 47: 128-39.
- 54 Martin E. Different pathomorphological aspects of pancreatic fibrosis correlated with etiology: anatomical study of 300 cases. In: Gyr K, Singer MV, Sarles H, eds. *Pancreatitis. Concepts and classification*. Amsterdam: Elsevier Science, 1984: 77-80.
- 55 Sarles H. Etiopathogenesis and definition of chronic pancreatitis. *Dig Dis Sci* 1986; 31: 91S-107S.
- 56 Sarles H, Bernard JP, Johnson CD. Pathogenesis and epidemiology of chronic pancreatitis. *Ann Rev Med* 1989; 40: 453-68.
- 57 Clemente F, Estival A, Durand S, Ribet A. Biochemical events in rat pancreatic cells in acute and chronic alcohol intoxications. In: Gyr KE, Singer MV, Sarles H, eds. *Pancreatitis. Concepts and classification*. Amsterdam: Excerpta Medica, 1984: 111-6.
- 58 Norohna M, Bordalo O, Dreiling DA. Alcohol and the pancreas. II. Pancreatic morphology of advanced alcoholic pancreatitis. *Am J Gastroenterol* 1981; 76: 120-4.
- 59 Tasso F. Etude ultrastructurale du pancreas dans 8 cas de pancreatites chroniques. *Ann Anat Pathol* 1967; 12: 343-72.
- 60 Boustiere C, Sarles H, Lohse J, Durbec JP, Sahel J. Citrate and calcium secretion in the pure human pancreatic juice of alcoholic and nonalcoholic men and of chronic pancreatitis patients. *Digestion* 1985; 32: 1-9.
- 61 Mohan V, Mohan R, Sushella L, et al. Tropical pancreatic diabetes in South India: heterogeneity in clinical and biochemical profile. *Diabetologia* 1985; 28: 229-32.
- 62 Clain JE, Barbezat GO, Marks IN. Exocrine pancreatic enzyme and calcium secretion in health and pancreatitis. *Gut* 1981; 22: 355-8.
- 63 Pitchumoni CS, Thomas E. Chronic cassava toxicity. Possible relationship to chronic pancreatic disease in malnourished populations. *Lancet* 1973; i: 1397-8.
- 64 Saunier JF, Sarles H. Exocrine pancreatic function and protein-calorie malnutrition in Dakar and Abidjan (West Africa). Silent pancreatic insufficiency. *Am J Clin Nutr* 1988; 48: 1233-8.
- 65 Durbec JP, Sarles H. Multicenter survey of the etiology of pancreatic diseases. Relationship between the relative risk of developing chronic pancreatitis and alcohol protein and lipid consumption. *Digestion* 1978; 18: 337-50.
- 66 Balakrishnan V, Saunier JF, Hariharan M, Sarles H. Diet, pancreatic function and chronic pancreatitis in South India and in France. *Pancreas* 1988; 3: 30-5.
- 67 Saunier JF, Sarles H, Attia Y, et al. Exocrine pancreatic function of children from the Ivory Coast compared to French children. Effect of kwashiorkor. *Dig Dis Sci* 1986; 31: 481-6.
- 68 Sarles H, Lahaie R, Dollet JM, Beck B, Michel R, Debry G. Effect of parental malnutrition on enzyme content of rat pancreas. *Dig Dis Sci* 1987; 32: 520-8.
- 69 Montalto G, Multigner L, Sarles H, De Caro A. Organic matrix of pancreatic stones associated with nutritional pancreatitis. *Pancreas* 1988; 3: 262-8.
- 70 Stafford RJ, Grand RJ. Hereditary disease of the exocrine pancreas. *Clin Gastroenterol* 1982; 11: 141-70.
- 71 Sarles H, Camatte R. Etiopathogenesis of pancreas. In: *Recent advances in gastroenterology*, Vol 4. Tokyo: Third Congress of Gastroenterology, 1966.
- 72 Marks IN, Bank S, Louw JH. Chronic pancreatitis in the Western Cape. *Digestion* 1973; 9: 447-53.