LETTERS TO THE EDITOR

Postprandial mesenteric blood flow

SIR,—We read with interest the study by Sieber *et al*¹ in which they used Doppler ultrasound to measure superior mesenteric arterial blood flow in response to liquid test meals given orally and intraduodenally (with and without atropine), and the additional effects of the infusion of combinations of cholecystokinin octapeptide 8, secretin, gastrin 17, and glucagon. We would like to raise several points.

(1) The authors used a 3 MHz Doppler flowmeter. Wavelength considerations with this frequency probe dictate that the best AP spatial resolution of the diameter measuring cursor is ± 1 mm (a best 'guess' of 0.5 mm can sometimes be made). The authors, however, have quoted superior mesenteric artery diameter readings to two decimal points and claim to be able to show increases in the diameter of between 5 and 26% after food intake. Furthermore, it is not clear whether an AP and lateral cross sectional diameter reading or a single longitudinal AP diameter reading was taken (in which case the superior mesenteric artery was assumed to be circular). It would have been better if the authors had quoted the mean and the range of the diameter reading.

(2) The quoted increase in superior mesenteric arterial blood flow in response to the oral test meal (180–360% above basal) is far greater than previous reports.^{2–5} It is not clear from the data whether the same subjects were studied on all occasions and compared – a total of nine subjects was enrolled in the study but data are only shown for six. If the same six subjects were not examined on each occasion, direct comparisons between experiments are less reliable. This is further compounded by the clear differences in basal mean (SEM) flow rates between the experiments (data not all shown) of 443 (38) ml/minute before the oral test meal and 520 (56) ml/minute before the intraduodenal test meal. What is the day to day coefficient of variation for the technique in their hands? Were all observations performed by one operator?

(3) The authors state that they infused glucagon, cholecystokinin octapeptide 8, secretin, and gastrin in doses designed to simulate postprandial circulating concentrations. No data, however, are presented to support this, and the authors cannot therefore be sure that physiological concentrations of these hormones were achieved. Furthermore, the dose of glucagon used (500 ng/kg/hour or 8.33 ng/kg/minute) is a supraphysiological dose^{6–10} and would result in plasma concentrations much greater than those normally circulating postprandially. A dose of less than 3 ng/kg/minute would have been more appropriate. We cannot therefore agree with the authors' contention that the hormones 'are unlikely to be involved as blood borne hormones' in mediating splanchnic vasodilation on the data presented but agree with the hypothesis.

We have recently shown (unpublished data) that fasting superior mesenteric arterial blood flow in six subjects measured by Doppler ultrasound decreases during physiological

infusions of glucagon. A 1 ng/kg/minute infusion produced no detectable rise in basal glucagon concentrations (mean (SEM) basal glucagon values 144 (15) ng/l; after 30 minutes 1 ng/kg/minute glucagon infusion resulted in a value of 129 (80) ng/l) and a 3 ng/kg/minute glucagon infusion raised values to 214 (37) ng/l. Mean (SEM) fasting blood flow fell from 678 (97) ml/minute to 549 (88) ml/minute after 30 minutes of 1 ng/kg/minute glucagon infusion and to 453 (63) ml/minute after 20 minutes of a 3 ng/kg/minute infusion. There were no associated changes in cardiac output, stroke volume, blood pressure, pulse, or peripheral vascular resistance. This suggests that glucagon is a selective splanchnic vasoconstrictor at physiological concentrations and, indeed, is not involved in mediating the postprandial hyperaemia observed in previous studies. It is noteworthy, however, that Lee *et al*¹¹ using indocyanine dye clearance methods showed that supraphysiological infusions of 10 and 20 ng/kg/minute glucagon caused no systemic haemodynamic or total hepatic blood flow changes in a group of cirrhotic subjects. Azygous blood flow, however (and hence superior mesenteric blood flow), rose significantly in those with well compensated cirrhosis, suggesting that at supraphysiological doses (similar to those used by Sieber *et al*¹) glucagon may be a splanchnic vasodilator.

(4) The authors state that blood pressure and pulse were monitored throughout the experiments but apart from incomplete data for the intraduodenal test meal with and without atropine, they present no evidence of blood pressure or pulse rate changes in response to any of their meals or infusion experiments. The pulse and blood pressure response to meals varies with both the age of the subject¹² and meal composition,¹³ and these data should have been included or discussed in the current study.

(5) Atropine was shown to attenuate the postprandial hyperaemic response to the meal, suggesting that the cholinergic nervous system has a role in this change. The mechanism for the postprandial response is, however, likely to be multifactorial and may also involve β adrenergic¹⁴ and peptidergic mechanisms. Neurotensin,¹⁵ vasoactive intestinal polypeptide,¹⁶ and calcitonin gene related peptide¹⁷ are known to be powerful splanchnic vasodilators and may also be involved in postprandial splanchnic vasodilation.

G D BRAATVEDT

A E READ

R J M CORRALL

Department of Medicine

M HALLIWELL

P N T WELLS

Department of Medical Physics,

Bristol Royal Infirmary,

Bristol BS2 8HW

- 1 Sieber C, Beglinger C, Jaeger K, Hildebrand P, Stalder GA. Regulation of postprandial mesenteric blood flow in humans: evidence for a cholinergic nervous reflex. *Gut* 1991; 32: 361–6.
- 2 Norrby C, Dencker H, Lunderquist A, Olin T, Tylen U. Superior mesenteric blood flow during digestion in man. *Acta Chir Scand* 1975; 141: 197–202.
- 3 Qamar MI, Read AE, Skidmore R, Evans JM, Wells PNT. Transcutaneous Doppler ultrasound measurement of superior mesenteric artery blood flow in man. *Gut* 1986; 27: 100–5.
- 4 Cooper A, Braatvedt GD, Qamar M, *et al*. Fasting and postprandial splanchnic blood flow is reduced by a somatostatin analogue (octreotide) in man. *Clin Sci* 1991; 81: 169–75.
- 5 Kooner JS, Peart WS, Mathias CJ. The peptide release inhibitor, octreotide (SMS 201-995) prevents the haemodynamic changes following food ingestion in normal human subjects. *Q J Exp Physiol* 1989; 74: 569–72.
- 6 Rizza RA, Cryer PE, Gerich JE. Role of glucagon, catecholamines and growth hormone in human

glucose counterregulation. *J Clin Invest* 1979; 64: 62–71.

- 7 Bajorunas DR, Fortner JG, Jaspan J, Sherwin RS. Total pancreatectomy increases the metabolic response to glucagon in humans. *J Clin Endocrinol Metab* 1986; 63: 439–46.
- 8 Alford FP, Bloom SR, Nabarro JDN. Glucagon metabolism in man. Studies on the metabolic clearance rate and plasma acute disappearance time of glucagon in normal and diabetic subjects. *J Clin Endocrinol Metab* 1976; 42: 830–8.
- 9 Liddle RA, Gertz BJ, Kanayama S, *et al*. Regulation of pancreatic endocrine function by cholecystokinin: studies with MK-329 a nonpeptide cholecystokinin receptor antagonist. *J Clin Endocrinol Metab* 1990; 70: 1312–8.
- 10 Fioretto P, Trevisan R, Valerio A, *et al*. Impaired renal response to a meat meal in insulin-dependent diabetes – role of glucagon and prostaglandins. *Am J Physiol* 1990; 258: F675–83.
- 11 Lee SS, Moreau R, Hadengue A, Cerini R, Koshy A, Lebre C. Glucagon selectively increases splanchnic blood flow in patients with well-compensated cirrhosis. *Hepatology* 1988; 8: 1501–5.
- 12 Westenberg M, Lenders JW, Thien T. The course of blood pressure after a meal: a difference between young and elderly subjects. *J Hypertens* 1985; 3 (suppl 3): S417–9.
- 13 Heseltine D, Potter JF, Hartley G, Macdonald IA, James OFW. Blood pressure, heart rate and neuroendocrine responses to a high-carbohydrate and a high-fat meal in healthy young subjects. *Clin Sci* 1990; 79: 517–22.
- 14 Braatvedt GD, Flynn MD, Stanners A, Halliwell M, Goldie B, Corral RJM. Splanchnic vasodilation during hypoglycaemia in man – evidence for a beta adrenergic mechanism? [Abstract]. *Diabetic Med* 1991; 8: 11A.
- 15 Rossell S, Rokaeus A. Actions and hormonal functions of circulating neurotensin. *Clin Physiol* 1981; 1: 3–20.
- 16 Said SI, Mutt V. Potent peripheral and splanchnic vasodilator peptide from normal gut. *Nature (London)* 1970; 225: 863–4.
- 17 O'Halloran DJ, Bloom SR. Calcitonin gene related peptide. *BMJ* 1991; 302: 739–40.

Reply

SIR,—Dr Wells raises important points about our recent publication,¹ especially about the methodology used to assess superior mesenteric artery blood flow (SMABF). Several comments need clarification, however, and we would like to reply in the same order as given in the letter:

(1) We used a 3.5 MHz sector scanner for diameter measurements (and not a 3 MHz Doppler flowmeter as stated in the letter). This sector probe was combined with a pulsed Doppler flow meter (3.0 MHz). Diameters were presented as mean (SEM) cm units. The statement that we were reporting two decimal points is therefore rather misleading.

In our experience, the superior mesenteric artery has a circular anatomy which is an advantage compared with the portal vein where the determination of the cross sectional area of the vessel is complicated by an ellipsoid vessel shape.

(2) We are puzzled by the statement that the increases in postprandial SMABF reported in our paper were far higher than in the published reports: similar increases have been reported by others^{2–4} with comparable meals and caloric loads. On the other hand, the reports cited by Dr Wells are hardly comparable.

In one study,⁵ a different technique was employed to quantify SMABF (dye dilution). In a second, a lower caloric load was tested⁶ and in the study of Qamar⁷ only fasting SMABF was measured.

Nine subjects participated in this study. For all oral food experiments as well as the hormone studies, the same six volunteers were studied. For the experiments involving intraduodenal food application, three additional subjects were