PROGRESS REPORTS

Biology of pancreatic cancer

G J Poston, J Gillespie, P J Guillou

Abstract

Pancreatic cancer is the fifth leading cause of death from malignant disease in Western society. Apart from the fortunate few patients who present with a resectable small pancreatic adenocarcinoma, conventional treatment offers no hope of cure and has little palliative value. Over the past two decades major steps have been made in our understanding of the biology of pancreatic growth and neoplasia. This review sets out to explore these advances, firstly in the regulation of normal pancreatic growth, and secondly the mechanism which may be involved in malignant change of the exocrine pancreas. From an understanding of this new biology, new treatment strategies may be possible for patients with pancreatic cancer.

Carcinoma of the exocrine pancreas is now the fifth leading cause of death from malignant disease in Western society¹ yet there has been only marginal improvement in the outcome of treatment for pancreatic cancer since the beginning of this century.² This improvement is largely due to the use of radical surgery with low mortality in specialist centres of excellence.³ The diagnosis of carcinoma of the pancreas still remains a virtual death sentence for the patient. Unfortunately, most studies show an increasing incidence of the disease, particularly among the elderly, in whom it seems to have trebled over the past 50 years.¹ It is also apparent that despite advances in surgical technique, anaesthetic support, intensive care, anticancer chemotherapy, and radiotherapy, the five year survival rate for all patients diagnosed as having carcinoma of the pancreas is less than 1%.² The only cure at present for pancreatic cancer is surgical resection, which in the very few patients who present with tumours smaller than 2 cm in diameter may achieve five year survival rates as high as 37%.⁴ It seems clear therefore that if any improvement is to be made in the treatment of this awful condition then a radical rethink is necessary in approaches to treatment and this must begin with a thorough understanding of the biology of the disease from which potential new remedies might be designed. The purpose of this review is to set out our current knowledge of the biology of pancreatic cancer and explore some of the potential avenues of treatment which may arise.

Cancer is an abnormality of cell division and tissue growth. In necropsy studies ductal hyperplasia, which may be premalignant, and separate synchronous carcinoma in situ can be found in as

many as 20% of patients who die from pancreatic cancer.⁵ Allen-Merish⁶ carefully dissected 102 pancreases taken at necropsy from 102 patients dying from non-pancreatic disease and found evidence of ductal mucinous hyperplasia in 63 separate pancreases. In both the established carcinogen induced rodent models of pancreatic cancer, azaserine treated rats which develop acinar tumours⁷ and *N*-nitrosobis(2-oxopropyl) amine treated hamsters which causes ductal disease^{8,9} similar to most human disease,¹⁰ malignant tumours are preceded by hyperplastic and nodular changes and also an alteration of pancreatic endocrine activity.¹¹ It seems rational therefore to start with an overview of the regulation of normal pancreatic growth, discussing the possible abnormalities in this process which may lead to or promote tumour growth.

Normal pancreatic growth

Pancreatic development in humans first becomes detectable as two separate evaginations of the primitive foregut during the fifth week in utero. Over the subsequent two weeks these dorsal and ventral buds fuse to form a single glandular structure from which develop the ductular and acinar components of the exocrine pancreas and the separate cellular components of the endocrine pancreas.¹² Specific cell types arise either by mitotic division of existing cells, differentiation of specialised cells, such as ductal cells, from uncommitted precursor or stem cells, or by transformation of one type of differentiated cell to another type – for example, acinar to ductal.¹³ Despite the evidence that a major attack of acute pancreatitis in humans which results in loss of acinar tissue, if survived, will be followed by full recovery of both structure and function of the exocrine gland,¹⁴ no other objective data are available on the regulation of exocrine pancreatic morphology in humans. Most of our data are derived from animal studies, usually in rats.¹⁵

During fetal life cell replication occurs before acinar cell differentiation¹⁶; however, there is a high frequency of mitoses in fully differentiated acinar and ductal cells which exists even late in the gestational period¹⁷ and values as high as 14% have been reported for acinar cells immediately before birth.¹⁸ Over the period of suckling, however, this labelling index rapidly decreases to levels of 1%, falling one month postnatally to 0.1–0.2%, where it remains thereafter.¹⁸ Indeed, with advancing age there is even a loss of the normal pancreatic trophic response to hormones (cholecystokinin, secretin, bombesin, penta-

Academic Surgical Unit,
St Mary's Hospital
Medical School, London
W2 1NY
G J Poston
J Gillespie
P J Guillou

Correspondence to:
Mr G J Poston.

Accepted for publication
24 August 1990

gastrin) which normally promote pancreatic growth.^{19,20}

The mature pancreas will undergo hyperplastic growth in response to three situations: dietary manipulation, surgical procedures, and hormone stimulation. Obviously the result of each of these events does not occur in isolation from the other two but the evidence for each is best described individually.

Diet

Starvation in the rat will cause an appreciable reduction in total pancreatic DNA content after six days.²¹ These changes can be prevented by administering pentagastrin²² and during starvation are associated with a commensurate fall in intestinal mucosal cholecystokinin content and a compensatory increase in intestinal cholecystokinin content during refeeding which coincides with the recovery of pancreatic DNA content.²² In a subsequent study the same researchers showed that chronic feeding with cholestyramine, which binds luminal bile salts, produced pancreatic hyperplasia.²³ Similarly, inhibition of the other major components of pancreatobiliary secretion, the pancreatic enzymes, and in particular trypsin, by soya bean trypsin inhibitor, promotes pancreatic growth.^{24,25} Diet also modulates the response of the pancreas to epidermal growth factor. A diet in which two thirds of the calories are administered as fat causes increased amino acid incorporation into rat acinar cells in response to epidermal growth factor stimulation.²⁶

Surgical procedures

In 1885 Di Mattei reported that pancreatic resection in the dog resulted in pronounced mitotic activity in the residual gland.²⁷ More recent studies have shown that as little as 50% pancreatectomy will promote pancreatic regeneration in the remaining tissue in the rat with maximal incorporation of tritium labelled thymidine into replicating acinar cells seen as early as 36 hours after resection.²⁸ Surgical pancreatobiliary diversion of bile and pancreatic juice away from the proximal small bowel promotes pancreatic growth,²⁹ and vagal deafferentation after jejunectomy causes pancreatic atrophy in the pig,³⁰ suggesting a feedback role of pancreatic secretion in maintaining pancreatic size.

Ileocaecal resection³¹ and colectomy³² both promote pancreatic growth in the rat and several studies have shown that this response is not due to the subsequent hypergastrinaemia associated with these procedures.³³⁻³⁵

Hormones and other growth factors

It became apparent in the early 1970s that gastrointestinal hormones were of fundamental importance in the maintenance of gut integrity and promotion of intestinal growth. By the mid 1970s Enochs and Johnson were proposing that cholecystokinin, which is a prime mover of pancreatic acinar secretion, was also a major trophic hormone for the pancreas.³⁶ Chronic

treatment with this or its analogue caerulein causes pancreatic growth in rats³⁷⁻³⁹ and mice.^{40,41} Similarly gastrin, which shares the same four C-terminal amino acids as cholecystokinin, when administered as its analogue pentagastrin promotes pancreatic growth.^{37,40,42} The physiological role of cholecystokinin in maintaining pancreatic size in health and disease is important.⁴³ It has recently been shown that promotion of endogenous cholecystokinin is the major agent responsible for pancreatic growth after the addition of soya bean trypsin inhibitor⁴³⁻⁴⁵ and cholestyramine⁴⁶ to the diet and also after chronic pancreatobiliary diversion⁴⁶ and jejunal bypass.⁴⁷ Plasma concentrations of cholecystokinin are raised in pancreatic atrophy,⁴⁸ and caerulein will induce pancreatic hyperplasia in rat models of chronic pancreatic insufficiency⁴⁹ and after DL-ethionine diet induced pancreatic degeneration.⁵⁰

Other peptide hormones that promote pancreatic growth include secretin,³⁷⁻³⁹ neurotensin,^{51,52} bombesin,^{53,54} and epidermal growth factor.⁵⁵ On the other hand, chronic treatment with somatostatin results in a decrease in resting DNA synthesis of the normal exocrine pancreas within 24 hours of the onset of treatment, with appreciable reduction of total pancreatic DNA content after five days.⁵⁶ Caerulein stimulated growth of the normal pancreas is also inhibited by somatostatin.⁵⁶

Glucocorticoids promote pancreatic hyperplasia during suckling and acinar cell hypertrophy at all ages,⁵⁷ and this hypertrophic response is both promoted by and promotes the hypertrophic response of acinar cells to the cholecystokinin analogue caerulein.⁵⁸ Prostaglandins of the E and F series stimulate pancreatic growth and this response is independent of their effect on serum gastrin release.⁵⁹

Until recently the major problem in understanding normal pancreatic growth has been that all available models have been in whole animal preparations. The only tissue culture in vitro models used cell lines derived from spontaneous or carcinogen induced pancreatic carcinomas. In 1986 Logsdon and Williams reported the methodology for establishing cultures of adult mouse pancreatic acinar cells as monolayers on collagen gels.⁶⁰ They were able to show that the cholecystokinin analogue caerulein would directly stimulate growth in vitro,⁶⁰ and Logsdon has subsequently shown that cholecystokinin-8 and gastrin, by stimulating the cholecystokinin receptor, promote cell division.⁶¹ In contrast, bombesin, substance P, and carbachol, factors which interact with separate receptors but stimulate pancreatic secretion in the same way as cholecystokinin by mobilising intracellular Ca²⁺, did not have any effect on the growth of pancreatic acinar cells in vitro. Most recently, Logsdon has shown that epidermal growth factor insulin, and insulin-like growth factor 1, which work through separate intracellular tyrosine kinase pathways, will also induce acinar cell division.⁶² Pancreatic ductal cells, dissected from acinar deficient pancreas produced by DL-ethionine deficient diets, have been maintained with secretory integrity in vitro^{63,64} for periods of up to six months in tissue culture.⁶⁵

Intracellular mechanisms

The function of regulatory peptides and steroids is to transmit information; however, a specific receptor must exist for each to initiate its own specific response. Molecules of similar structure – for example, cholecystokinin and gastrin, epidermal growth factor and transforming growth factor alpha, insulin and insulin-like growth factor – may in turn act upon unique receptors, related receptors, or the same receptor particularly when they may involve two or more separate responses from the same cell. For example, cholecystokinin will both stimulate acinar secretion of enzymes and promote acinar cell division. The receptor proteins for peptide hormones exist on the cell membrane, whereas those for steroid hormones exist either in the cytosol or within the nucleus.⁶⁶ In the case of cholecystokinin the initial site of peptide receptor binding in acinar cells is the basolateral membrane domain and at physiological temperatures (30°–37°C) the bound hormone is subsequently internalised to be localised within specific intracellular compartments where it may have direct actions.⁶⁷ Similarly insulin, insulin-like growth factor II, epidermal growth factor, and somatostatin are all internalised in the acinar cells after receptor binding, although to varying degrees and extents.^{68,69} The main problem in interpretation of these data is that whereas Scatchard analysis of binding of radiolabelled agonist to receptor ligand will give the number and binding affinity of membrane receptors in each sample, this technique produces a static picture which does not clarify the events occurring beyond the membrane once internalisation has occurred. On the other hand, autoradiographic studies of radiolabelled agonist bound to receptors either on microscopy or after gel electrophoresis of lysed cells will give information about receptor size and site but not about the kinetics of the initial reaction. Thus most of the data derived so far about number of functional receptors per cell at any given moment are probably inaccurate.

Cell receptor populations are not static, constantly being increased ('upregulation') or decreased ('downregulation') by alterations in the rate of receptor synthesis and degradation in response to stimulus drive and target cell response.⁶⁶ Downregulation of acinar receptors has been shown *in vitro* for insulin and acetylcholine.^{70,71} Other factors that affect the binding characteristics of specific receptors include occupancy of totally separate hormone-receptor species. For instance, somatostatin enhances binding of oestradiol to its cytosolic protein in the rat pancreas⁷² and treatment with cholecystokinin decreases the binding of epidermal growth factor, insulin-like growth factor II, and somatostatin to their respective receptors.^{68,69,73}

Chronic treatment with cholecystokinin *in vivo*, sufficient to cause pancreatic hypertrophy and hyperplasia, does not increase the number of cholecystokinin receptors per cell but the total number of cholecystokinin receptors per pancreas increases.⁷⁴ Finally, the number and concentration of cholecystokinin receptors in the pancreas changes with age. During postnatal maturation and weaning in the rat, pancreatic

cholecystokinin receptors increase in number.⁷⁵ In the guinea pig they reach a peak by 1 year of age and then fall threefold by 3 years of age (median age of survival and equivalent to a 70 year old man).⁷⁶

In general those hormones which stimulate the release of pancreatic enzymes are mediated intracellularly by Ca^{2+} and diacylglycerol while those hormones which regulate ductal secretion are mediated via cyclic AMP,⁷⁷ although this division is not absolute.⁶⁶ A third class of hormone regulators which are primarily concerned with metabolism and induction of protein synthesis activate tyrosine kinase.⁷⁸ Unlike the stimulatory peptides, somatostatin regulates growth by stimulation of tyrosine phosphatase resulting in activated tyrosine phosphatase which in turn activates serine kinase and subsequently inhibits cell division and growth.⁷⁹

Pancreatic cytosolic receptors have been described for glucocorticoids, oestrogens, and androgens.^{66,72,80–86} Although oestrogen binding proteins exist in pancreatic cytosol with high binding affinity^{80,82} they show different reaction kinetics and are of different molecular size to the high binding affinity oestrogen receptors found in uterine cytosol.^{83,86} A separate atypical low affinity oestrogen binding site requiring a peptide cofactor has also been described in the pancreas.⁸⁷ Pancreatic oestradiol binding proteins are found in both ductal and acinar cells^{83,88} and in these cells act within the region of the endoplasmic reticulum.

Second messengers

As discussed earlier hormone and growth factor signals generally require the stimulation of a second intracellular chemical messenger (transduction) to begin the initiation of a response by the cell. These transduction pathways show autoregulation and directly interact at a number of important modulatory steps.⁸⁹ The initial generation of the calcium signal with release of neutral lipids is mediated by the breakdown of phosphatidylinositol and changes in cyclic nucleotide levels involve the activation of adenylate cyclase.

The breakdown of phosphatidylinositol is initiated through the action of a series of phospholipase C's and kinases to produce neutral lipids (which activate protein kinase C), inositol-triphosphate, and arachidonic acid.^{90,92} Subsequent studies have shown the action of some transforming oncogenes on phosphatidylinositol turnover,⁹³ the effect of phorbol esters on cell transformation,⁹⁴ and the influence of C-kinase on the epidermal growth factor receptor⁹⁵ and transferin receptor.⁹⁶ The phosphatidylinositol pathway promotes intracellular Ca^{2+} through inositol 1,4,5-triphosphate and this soluble product is seen immediately after a hormone-receptor interaction which is subsequently mediated via Ca^{2+} .^{90,91} Raised intracellular Ca^{2+} activates further enzyme pathways by binding to calmodulin, a calcium binding protein.⁹⁹ A rise in the intracellular concentrations of calmodulin have been found in cells undergoing malignant transformation.^{97,98} Lastly phosphatidylinositol breakdown generates inositol phosphate

mediators from arachidonate, probably through the action of phospholipase A2 on phosphatidyl choline or phosphatidyl ethanolamine.⁹⁴

The adenylate cyclase system responds to ligand receptor interaction with the generation of cyclic nucleotides, either promoting or inhibiting cyclase activity.⁹⁰ The G proteins of the cyclase system influence the affinities of receptors for ligand and so may play a part in receptor sensitisation and desensitisation.⁹⁹ Adenylate cyclase has three primary components: G_s , a stimulating subunit; G_i , an inhibiting subunit; and a third, catalytic, subunit.¹⁰⁰

Nuclear events

Little is known of what occurs after intracellular second messenger promotion during events which stimulate pancreatic growth. More is known, however, about the regulation of pancreatic enzyme production during the same dietary changes described earlier which promote pancreatic growth.^{101, 102} It seems reasonable therefore to review these pathways since some inference may be drawn from stimulatory events which promote both pancreatic secretion and growth. In the pancreas the synthesis of functional groups of enzymes is regulated by specific hormones, whether these hormones are administered exogenously or produced endogenously.¹⁰³

Production of proteins by the cell, whether for use within the cell or for export, begins with transcription of the DNA code of a particular gene to messenger RNA (mRNA) by RNA polymerase II. Within the nucleus this primary mRNA transcript is processed by either the removal of intron sequences by splicing mechanisms or the addition of a poly (A) tail to the 3' terminus.¹⁰³ The processed mRNA is now transported out to the cytoplasm where, after the functional binding of the 40S and 60S ribosome subunits, the coding sequence of the mRNA is translated into protein.¹⁰⁴ The protein product is then either used or packaged for export. Regulation of gene expression as protein product can therefore occur: (a) at the original transcription of DNA; (b) post-transcriptional processing; (c) rate of mRNA transport; (d) mRNA translation; (e) mRNA breakdown; (f) post-translational processing and cleavage of the protein product; (g) storage of protein product; (h) release or use of protein product. The events which regulate gene expression involve the interaction of regulatory proteins with specific nucleotide sequences on DNA.¹⁰⁵ RNA polymerase II initiates transcription after binding to the promoter (TATA) sequence of the gene which usually lies 30 nucleotides upstream from the DNA nucleotide at which transcription begins.¹⁰³ The RNA polymerase then moves downstream from the promoter site, unwinding the DNA and polymerising a single strand of mRNA, regulated further by enhanced gene sequences which lie further upstream in the 5' flanking region of genes.¹⁰³ Enhanced elements either regulate gene expression in a tissue specific fashion or in response to individual hormones.¹⁰⁶ Our knowledge of hormone regulation is, however, largely confined to steroid hormones which have receptor proteins that bind directly to DNA¹⁰⁷ and so

regulate gene expression at the level of transcription. Little is understood of the mechanisms of gene expression regulation by peptide hormones and growth factors. Scheele and Kern with their groups have carefully analysed this response by the exocrine pancreas to cholecystokinin and secretin.¹⁰⁸ They have shown that pronounced changes in individual enzyme biosynthesis were not accompanied by changes in mRNA levels, at least during the first six hours of stimulation, suggesting that these initial changes in gene expression are due to alterations in post-translational processing. This time related difference in the mechanism of gene expression is probably because exocrine pancreatic product mRNAs have relatively long half lives of up to six hours and therefore sudden increases in demand for protein synthesis can be met both at and beyond the translational level,¹⁰³ which is the rate limiting step in protein synthesis. Changes at this level may involve nucleotide signals in the 5'-non-translated region of mRNAs and the interaction of these signals or sequences with either regulatory proteins or regulatory RNA molecules.¹⁰³

In general, growth factors such as insulin, insulin-like growth factor-1, epidermal growth factor, and platelet derived growth factor all stimulate phosphorylation of the ribosomal protein S6, but the subsequent effect on promoting or inhibiting cell growth and mitosis is cell type specific.¹⁰⁹⁻¹¹³ In 3T3 fibroblasts, treatment with platelet derived growth factor stimulates expression of the *c-myc* and *c-fos* oncogenes,¹¹⁰ and similar changes in expression of these oncogenes are seen in many cancer cells.

Polyamines

Polyamines (putrescine, spermidine, and spermine) are ubiquitous highly charged cations that are required for growth and differentiation in all eukaryotic cells^{114, 115} and are particularly important in the regulation of DNA, RNA, and protein synthesis.¹¹⁶ Membrane stability and cyclic AMP independent protein kinases are apparently also influenced by polyamines.¹¹⁶ The mechanisms by which polyamines exert their effects are poorly understood, though their polybasic nature has been proposed as being an important factor.¹¹⁷ High levels of polyamines are found in rapidly dividing cells and a depletion of intracellular polyamines causes a slowing and eventual stopping of cell growth.¹¹⁸

The biosynthesis of polyamines in mammalian systems begins with the decarboxylation of ornithine by the enzyme ornithine decarboxylase to form the diamine putrescine. This is the first, and probable rate limiting step, in the polyamine biosynthetic pathway, and ornithine decarboxylase activity is greatly raised during cell division.¹¹⁹ Subsequently putrescine is converted by spermidine synthase to spermidine, which in turn is converted to spermine by spermine synthase.¹²⁰ These last two steps are potentially reversible by polyamine oxidase.¹²⁰ During caerulein stimulated pancreatic growth there are increases in content and concentration of all three polyamines which occur within 12-96 hours after the onset of caerulein administration.^{121, 122} Total

spermidine and spermine content also correlates with the rates of increase in both pancreatic weight and DNA content.¹²¹ Because of their possible antimitogenic potential, many pharmacological inhibitors of polyamine biosynthesis have been developed,¹¹⁶ most importantly against ornithine decarboxylase because of its key regulatory role.¹²³ One of these is alpha-difluoromethyl-ornithine (DFMO). DFMO treatment suppresses the increase in pancreatic weight induced by caerulein¹²⁴ and this effect can be reversed by administration of putrescine, the biosynthetic product of ornithine decarboxylase.¹²⁵ Thus the stimulation of pancreatic DNA synthesis by caerulein seems to be a polyamine dependent process.

The potent phosphodiesterase inhibitor 3-isobutylmethylxanthine causes increases in ornithine carboxylase, putrescine, and *N*'-acetyltransferase levels in the rat pancreas and liver. DFMO decreases the accumulation of putrescine in the liver but not the pancreas.¹²⁶ These findings suggest that the accumulation of putrescine in the pancreas occurs primarily by the action of the polyamine interconversion enzymes, *N*'-acetyltransferase and polyamine oxidase.

Having reviewed the current opinions on the regulation of normal pancreatic growth we now consider the present views on the abnormalities and regulation of growth in pancreatic cancer.

Aetiological factors of pancreatic cancer in humans

Exocrine pancreatic cancer is a disease of Western society with a tenfold difference in prevalence between countries where the disease is common (UK, USA) and countries where it is less frequent (India, Nigeria).¹²⁷ There are also racial predilections with blacks in the Bay Area of California having 1.5 times the risk of developing the disease as whites in the same area; however, diet and environment also play a part as American blacks have a greater chance of developing the disease than African blacks.¹²⁷ Within the United States the incidence of pancreatic cancer in blacks is five times greater than that of Japanese Americans or Hispanics from Puerto Rico.¹²⁷

Other established risk factors include age and sex. All studies show the disease to be commoner in men than in women, with a ratio ranging from 1.5:1 to 2:1 depending on the community.¹²⁸ This may be hormonal, but men are also exposed on the whole to other risk factors for the disease. There is an established link between pancreatic cancer and cigarette smoking; it seems likely that nitrosamines may have a causal role.¹²⁷ Whether these chemicals reach the pancreas directly or, after hepatic metabolism, are excreted in the bile and so affect the pancreas is unknown. Smoking also raises serum lipids which may predispose to pancreatic cancer, since high fat diets are also related to the increased incidence of the disease.¹ Other factors associated with an increased incidence of pancreatic cancer include working in chemical industries (particularly beta-naphthylamine and benzidine), petroleum plants, coke and coal gas plants, dry cleaning industries,

and plants handling radioactive materials for nuclear fuels and warheads.¹²⁷ An unsubstantiated risk factor is urban living. Earlier studies between 1935 and 1974 suggested this association but intensive county-by-county studies performed in Connecticut and Iowa in 1974 did not bear this out.¹²⁷

Coffee and caffeine have been linked to pancreatic cancer but this link is tenuous since such drinking habits are often linked to other risk factors like cigarette smoking. Also, the incidence of coffee drinking tends to increase during the early stages of the disease when patients become anxious about insidious symptoms or develop glucose intolerance.¹ Similarly, excessive alcohol consumption does not predispose to pancreatic cancer. Gall stone disease and chronic pancreatitis have been linked to the development of pancreatic cancer but close analysis of the data, combined with better diagnostic criteria, have challenged this connection.^{127 128} A causal relation with diabetes (which at one time was thought to increase the chance of developing pancreatic cancer threefold) has been confused by the fact that up to 15% of patients with pancreatic cancer will have developed diabetes because of their cancer in the immediate period before presentation. When this factor is taken into account, there is no difference in the prevalence of pancreatic cancer between diabetics and non-diabetics.¹²⁹

Aetiological factors of pancreatic cancer in animal models

The adaptive changes of hypertrophy and hyperplasia in the pancreas are diffuse and reversible; on the other hand, neoplastic growth is clonal and involves multiple steps through preneoplastic changes to produce a tumour.¹³⁰ Completion of the sequence of changes to irreversible malignant growth usually requires a major portion of the lifespan of the host, and the rate of progression of preneoplastic lesions to cancer may be regulated by the same factors that control adaptive growth. The major difference between neoplastic growth and adaptive growth is that the former is the result of altered division of single cells (clonal growth) and the latter represents the regulated growth of the whole organ. Certainly in the two rodent models of pancreatic cancer (azaserine induced acinar cancer in the rat and *N*-nitrosobis(2-oxopropyl)amine induced ductal cancer in the hamster) the sequence of progression is: initiated cell → focus → nodule → neoplasm (carcinoma).¹³⁰ The sensitivity of cells to carcinogenesis is promoted if the cells are already undergoing mitosis – rats are highly sensitive to the induction of preneoplastic acinar lesions when carcinogens are given during the first three weeks of life¹³¹; during pancreatic regeneration¹³²; after pancreaticobiliary diversion^{133 134}; and after administration of exogenous or a rise in endogenous cholecystokinin.^{130 135-137}

As in humans, dietary lipids promote the growth of carcinogen induced pancreatic cancer in rodents. Diets that include 4–8% unsaturated linoleic fatty acid enhance the development of azaserine induced acinar cancer in Lewis rats,¹³⁸ and lithocholic acid when combined with

cholecystectomy produced greater numbers of *N*-nitrosobis(2-hydroxypropyl)amine ductal tumours in hamsters.¹³⁹

Endocrine regulation of pancreatic cancer

The idea that cancers which arise from hormone target organs possibly maintain their hormone responsiveness has evolved into the classification of cancers as hormone dependent or independent and has led to the development of treatment regimens aimed at arresting tumour growth by either hormone deprivation or administration of antagonistic agents of these hormones. The latter would include treatment using other hormones whose action would oppose those hormones which promote tumour growth. The concept of growth factor and hormone regulation of cancer has been extensively and recently reviewed by Goustin and colleagues¹⁴⁰ and Townsend and coworkers have specifically reviewed the role of growth factors in intestinal neoplasms.¹⁴¹

Direct effects of hormones on pancreatic carcinogenesis

Importantly, a chronic rise in plasma cholecystokinin by longterm feeding with raw soya flour (for more than four months) will cause hyperplastic foci of pancreatic acinar cells.^{142 143} If this diet is discontinued after 24 weeks then pancreatic morphology will subsequently revert to normal; however, if the diet is continued for up to 36 weeks some of these hyperplastic foci will progress to pancreatic cancer, even without the addition of carcinogens.¹⁴³ The link between cholecystokinin and carcinogenesis, however, is not straightforward since Pour and colleagues have shown in the hamster model of pancreatic ductal carcinogenesis that exogenous cholecystokinin, given simultaneously with or shortly before the carcinogen *N*-nitrosobis(2-oxopropyl)amine, inhibited cancer induction.¹⁴⁴⁻¹⁴⁶ On the other hand, Satake and colleagues found that weekly administration of caerulein enhanced the carcinogenic effect of *N*-nitrosobis(hydroxypropyl)amine.¹⁴⁷ Similarly, Howatson and Carter found that both cholecystokinin¹⁴⁸ and secretin¹⁴⁹ enhanced *N*-nitrosobis(2-oxopropyl)amine pancreatic carcinogenesis in hamsters. It may be that the studies which failed to show any effect used insufficient dosages of caerulein to achieve the hyperplastic response necessary for carcinogenesis, but the inhibitory studies may also relate to the phenomenon that caerulein also inhibits the growth of a human cholangiocarcinoma, known to possess cholecystokinin receptors, when growing in nude mice.¹⁵⁰ Epidermal growth factor promotes *N*-nitrosobis(2-oxopropyl)amine pancreatic carcinogenesis in hamsters¹⁵¹ and both the somatostatin analogue RC-160 and luteinising hormone releasing hormone reduce *N*-nitrosobis(2-oxopropyl)amine pancreatic carcinogenesis and prolong survival in hamsters.¹⁵²⁻¹⁵⁴

Hormone regulation of established tumours

CHOLECYSTOKININ AND SECRETIN

There are good data to show that cholecystokinin

will promote the growth of established pancreatic cancers in the same way that cholecystokinin stimulates normal pancreatic growth. Townsend and coworkers reported that the combination of caerulein and secretin (but neither agent alone) produced significant stimulation of growth in a transplantable hamster pancreatic ductal cancer (H2T).¹⁵⁵ Receptors for both hormones have been found in a human pancreatic cancer cell line.¹⁵⁶ Asperlicin, a competitive non-peptide cholecystokinin antagonist, significantly inhibits the growth of a human pancreatic adenocarcinoma (SKI), established in nude mice in Dr Courtney Townsend's laboratory in Galveston, that possesses cholecystokinin receptors,¹⁵⁷ probably by inhibiting the effect of endogenous cholecystokinin.¹⁵⁷ A similar effect is seen with the cholecystokinin antagonist CR-1409 on exogenous cholecystokinin promoted tumour growth in azaserine induced rat pancreatic cancer.¹⁵⁸ Upp and colleagues reported that the presence of cholecystokinin receptors on a human pancreatic cancer will predict the response of the cancer to caerulein treatment.¹⁵⁹ Caerulein produced a pronounced stimulation of growth in the tumour possessing cholecystokinin receptors and this effect was inhibited by proglumide, the gastrin-cholecystokinin antagonist. Growth of the tumour without cholecystokinin receptors was not affected by treatment. Furthermore, caerulein treatment enhanced the presence of high affinity binding receptors for cholecystokinin in the SKI tumour.

BOMBESIN

Bombesin, a member of the gastrin releasing peptide family of hormones, normally promotes pancreatic growth¹⁶⁰ and indeed will promote the growth of azaserine induced rat acinar tumours.¹⁵⁸ Chronic treatment with bombesin, however, inhibits the growth of human pancreatic cancer in nude mice while stimulating the growth of the normal nude mouse pancreas.¹⁶¹ This demonstration of simultaneous trophic and inhibitory responses by the same hormone on normal and malignant tissue suggests that the action of bombesin, directly or indirectly through the hormones that it stimulates, may be site specific, interacting differently in normal and neoplastic tissues. Possible explanations for this phenomenon may be that different dosages of the same hormone may act differently: multiple forms of hormone receptor may exist to serve different functions (note the effects on growth and secretion discussed earlier); hormones may act directly on target cells through hormone-receptor regulation (homologous or heterologous); or hormones may act indirectly by releasing other hormones or growth factors (autocrine, paracrine or endocrine) to produce their response from the target cell.

VASOACTIVE INTESTINAL PEPTIDE

Vasoactive intestinal peptide receptors exist on both normal¹⁶² and neoplastic¹⁶³ pancreatic tissue. Binding of the peptide to these receptors promotes intracellular cyclic AMP accumulation,¹⁶⁴ which may be a regulatory factor in cell division

and growth.¹⁶⁵⁻¹⁶⁷ Poston and colleagues have shown that chronic treatment with vasoactive intestinal peptide inhibits the growth of hamster pancreatic cancer *in vivo* but not human pancreatic cancer.¹⁶⁸ This effect may be related to the presence of two vasoactive intestinal peptide receptor types (one with molecular weight 66 kDa, and the other of 90 kDa) on the hamster pancreatic cancer cells whereas only one type of peptide receptor (MW 66 kDa) was detectable on the human cancer cell lines.¹⁶³

SOMATOSTATIN

Somatostatin inhibits the growth of several malignant tumours and tumour cell lines¹⁶⁹⁻¹⁷³ and its effect on pancreatic carcinogenesis has been described.¹⁵²⁻¹⁵⁴ The oncological application of somatostatin and its analogues have recently been extensively reviewed by Schally and colleagues.¹⁷⁴⁻¹⁷⁶

In 1984 Redding and Schally described the inhibition of both rat and hamster pancreatic cancer by administration of somatostatin.¹⁷⁷ Subsequently Upp and colleagues have reported that the somatostatin analogue SMS 201-995 inhibits the growth of human pancreatic cancer in nude mice.¹⁷⁸ Recent reports suggest that this response may be potentiated by the oestrogen antagonist tamoxifen.¹⁷⁹ *In vitro*, somatostatin reverses the growth potentiating effect of epidermal growth factor on MIA PaCa-2 human pancreatic cancer cells⁷⁹ and H2T hamster ductal pancreatic cancer cells (unpublished data from our laboratory), which may be through promotion of tyrosine phosphatase activity acting on the epidermal growth factor receptor.⁷⁹

Not only does somatostatin treatment prolong tumour doubling time¹⁷⁸ but it may result in programmed cell death (apoptosis) in carcinogen induced pancreatic cancer in hamsters, in some cases completely preventing the development of carcinogen induced tumours.¹⁸⁰ Although evidence exists from *in vitro* studies for a directly mediated response,^{181 182} somatostatin may also affect tumour growth *in vivo* by inhibiting other hormones such as cholecystokinin which may promote tumour growth.¹⁸³ We have recently shown somatostatin analogue inhibition of epidermal growth factor and transforming growth factor alpha stimulated growth of pancreatic cancer in a hamster model (unpublished data from our laboratory). Somatostatin receptors have been shown on a well differentiated cell line derived from rat carcinogen induced acinar cancer AR42J.¹⁸⁴ Under the auspices of the Cancer Research Campaign, we have now started clinical trials on the use of long acting somatostatin analogues for treatment of pancreatic cancer but care will be necessary because of the potential side effects on other gastrointestinal pancreatic hormones, and in particular on the management of serum glucose.¹⁸⁵

Epidermal growth factor and transforming growth factors

As described earlier, epidermal growth factor promotes pancreatic carcinogenesis in hamsters^{151 186} and promotes the growth of human

pancreatic cancer cells *in vitro*.⁷⁹ Retrovirus insertion of epidermal growth factor receptor gene into 3T3 fibroblasts is associated with malignant transformation, the degree of which is proportional to the extent of epidermal growth factor receptor expression.¹⁸⁷ Carcinoma cells which overexpress the epidermal growth factor receptor produce a 270 kDa epidermal growth factor receptor gene transcription factor¹⁸⁸ and that form of this factor can result in tenfold gene expressions of epidermal growth factor receptor gene.¹⁸⁹ DNAase 1 footprinting studies have identified four binding sites for these transcription factors on the epidermal growth factor receptor gene promoter.¹⁹⁰

Normally, stimulation by epidermal growth factor downregulates its receptor expression, so regulating the number of available epidermal growth factor receptors¹⁹¹⁻¹⁹³ with phosphorylation of membrane proteins.¹⁹⁴ The pancreas is one of the richest sources in the body of mRNA for prepro-epidermal growth factor,¹⁹⁵ so raising the question of autocrine growth stimulation of pancreatic cancer by epidermal growth factor production.¹⁹⁶ The epidermal growth factor receptor is the cellular homologue of the avian erythroblastosis virus *erb-B* proto-oncogene.^{197 198} The gene for pancreatic neuropeptide Y and *erb-B* overlap on human chromosome 7¹⁹⁹ and Korc *et al* have recently shown that human pancreatic cancer cells overexpress the gene for epidermal growth factor receptor associated with concomitant changes on chromosome 7.²⁰⁰ These increases in expression of epidermal growth factor receptor may be as great as 100-fold.^{201 202}

Normal fibroblasts must attach to a surface to grow *in vitro*. If exposed to substances which reversibly transform them, however, these same fibroblasts can grow in suspension; such factors have been termed transforming growth factors (TGF).^{203 204} Two groups of these growth factors have been identified. TGF α , 5-6 kDa polypeptide structurally related to epidermal growth factor with which it shows a 35% homology by possession of six cysteine residues in the same relative positions, and can thus bind to and stimulate the epidermal growth factor receptor.²⁰⁵ The TGF β s are a separate distinct polypeptide family of about 25 kDa in size and have no sequence homology with TGF α .^{203 206 207} TGF β s do not bind to the epidermal growth factor receptor²⁰⁸ but stimulate a separate unique receptor to induce *c-sis* oncogene mRNA expression which in turn induces a mitogenic response through production of platelet derived growth factor.²⁰⁹ But unlike their effect on mesenchymal cells, TGF β s have a different effect on epithelial cells, being the most potent growth inhibitory peptide known for a number of epithelial cells.^{210 211} Interestingly, transfection of mouse fibroblast cells with a *c-myc* oncogene results in acquisition of responsiveness to TGF β .²¹² Both TGF α and β are produced by breast^{213 214} and colon cancers.²¹⁵

Several cultured human pancreatic cancer cell lines produce TGF α in addition to epidermal growth factor^{201 205} and TGF α is 10-100 times more potent than epidermal growth factor in stimulating anchorage independent growth of

these same cell lines.²⁰⁵ TGF α binds to pancreatic epidermal growth factor receptors where it is degraded after receptor stimulation.²⁰⁵ The binding of TGF α to the epidermal growth factor receptor does not however downregulate the expression of the epidermal growth factor receptor as occurs normally with epidermal growth factor stimulation^{191-193 205} and so may result in an uncontrolled autocrine growth effect. This effect can be reversed by culturing the cells on soft agar containing anti-TGF α monoclonal antibody.²⁰⁵ On the other hand, some pancreatic cancer cells have been shown to produce TGF β , which in turn inhibits their own growth.²⁰⁵ These findings of Korc's group suggest that TGF α participates in the regulation of pancreatic cancer cell proliferation and point to the possible existence of a TGF α epidermal growth factor receptor autocrine cycle whose function may be modulated by TGF β .²¹⁶ We have just shown that exogenous administration of TGF α stimulates the growth of hamster pancreatic cancer in vivo (unpublished data from our laboratory).

STEROIDS AND LUTEINISING HORMONE-RELEASING HORMONE

Recent interest has focused on the use of steroid hormones and their antagonists in the treatment of pancreatic cancer.^{217 218} Certainly, oestrogen receptors can be shown on normal^{219 220} and neoplastic²²⁰⁻²²⁵ pancreatic tissue in both human and carcinogen induced rat acinar tumours. Androgen receptors are found on both normal pancreas²²⁶ and pancreatic cancers²²⁷ and both glucocorticoids^{228 229} and sex steroid hormones have a regulatory role in normal exocrine pancreatic secretion and growth.²³⁰ When given to spayed female mice relatively high amounts of the active metabolite of tamoxifen, 4-hydroxy-tamoxifen, have been found in the normal pancreas as long as 24 hours after administration,²³¹ and these findings have formed the basis of early clinical trials in patients with pancreatic cancer which have claimed some limited success for antioestrogen²³² and luteinising hormone-releasing hormone antagonists to suppress endogenous oestrogen.^{233 234}

In the laboratory it is much more difficult to induce pancreatic cancers using azaserine in female rats than it is in male rats,²³⁵ but this can be overcome with prior oophorectomy and tamoxifen treatment.²³⁵ Similarly, oestrogen treatment and castration inhibit the early stages of acinar pancreatic carcinogenesis after azaserine treatment in male rats.²³⁶ In human pancreatic cancer cells there is good evidence that 4-hydroxy-tamoxifen produces its growth inhibitory effect through specific antioestrogen binding sites,^{237 238} although these effects bear little relation to oestrogen receptor content.²³⁸

Plasma concentrations of circulating androgens are decreased in patients, particularly men, with pancreatic cancer,²³⁹⁻²⁴³ and this has led some to claim that low concentrations of testosterone may be an endocrine marker for the disease.^{241 243} Certainly studies on rat pancreatic cancer²⁴⁴ and human pancreatic cancer in nude mice^{245 246} and tissue culture^{238 246} show a trophic effect of androgens on the growth of the tumour cells.

This may be a factor in the predilection of the disease for males in general.¹ In male hamsters, however, castration promotes the growth of transplantable ductal cancers but this may be mediated through the tumour promoting effect of castration by increasing the release of endogenous gastrin.²⁴⁷ Testosterone may act either directly on the cancer cells or be converted by aromatase to oestrogen or by 5- α -reductase to the more potent 5- α -dihydrotestosterone, after internalisation into the pancreatic cancer cells.²⁴⁸

Glucocorticoids stimulate the growth of human pancreatic cancer cells and rat AR42J acinar cancer cells in vitro²³⁸ and also promote the synthesis of amylase mRNA levels, secretory organelles, and enzyme secretion in AR42J cells in tissue culture.²⁴⁹ Again, there is variance in transplantable hamster ductal pancreatic cancer where dexamethasone treatment inhibits tumour growth (unpublished results).²⁵⁰

Intracellular events

Little is known about oncogene activity and the role of oncogenes in human pancreatic cancer. Cooper and colleagues have characterised expression of activated *ras*^K gene in human pancreatic cancer cell lines²⁵⁰ and activation of this gene appears to be due to an amino acid change on position 12 of the *ras*^{K-2} sequence.²⁵¹

Palmiter and Brinster have created a transgenic model of acinar pancreatic cancer in mice by fusing the SV-40 tumour virus to the elastase 1 gene, which is only expressed in the pancreas.²⁵² These mice develop pancreatic cancer by three months of age and will prove a powerful tool in the dissection of molecular events leading to the induction of pancreatic cancer.

Ornithine decarboxylase activity is greatly raised in hamster pancreatic cancer cell lines²⁵³ and inhibition of this with DFMO inhibits the growth of both human and hamster pancreatic cancers.^{254 255} This effect is reversed by the administration of exogenous putrescine²⁵⁴ and potentiated by the addition of cyclosporine both in vitro^{256 257} and in vivo.²⁵⁸

Tumour cells characteristically exhibit an increased rate of glycolysis,²⁵⁹ the so called 'Warburg effect,' but little is known of the regulation of this mechanism in pancreatic cancer cells. Schek and colleagues have shown by using complementary DNA (cDNA) probes that there is an overexpression of mRNA for glyceraldehyde-3-phosphate dehydrogenase, enolase, and glucose transporter protein which may play a part in the Warburg effect.²⁶⁰

The ability of malignant tumours to invade normal tissue is dependent on the production of proteases such as collagenase, plasminogen activators, and lysosomal proteases. The human pancreatic carcinoma cell line HPC-YT secretes excessive amounts of acid proteases which may play a part in this process.²⁶¹

Finally, much clinical interest has focused recently on the production of antigen markers by pancreatic cancer. In 1979 Kaprowski and colleagues reported that raised levels of CA19-9, a carbohydrate antigenic determinant were present in sera of patients with gastrointestinal cancer²⁶² and subsequent workers have shown

that it is raised in the serum of as many as 80% of patients with pancreatic cancer.^{263 264} Pancreatic cancer can also produce carcinoembryonic antigen,²⁶⁵ tumour associated glycoprotein (TAG 72), and DU-PAN-2.²⁶⁶

Conclusion

Much is known about the physiological, pharmacological, and surgical events which will promote pancreatic growth and carcinogenesis. We are beginning to understand the intracellular events which regulate normal pancreatic hypertrophy and hyperplasia. Although we now know that pancreatic cancers will respond in many similar ways to these events, there are vast gaps in our knowledge of the intracellular events which occur after pancreatic carcinogenesis. Certainly it is well established that there is a wide variability of biology and response to treatment within pancreatic exocrine cancers as has been carefully shown by Klapdor's group.^{267 268} The greatest challenge to pancreatic oncology during the next decade will be in dissecting out these intracellular mechanisms and differences in biology and establishing where the disorders of regulation exist.

This work is supported by the British Digestive Foundation of which GJP is the Amelie Waring scholar, and by the Cancer Research Campaign (Project Grant SP 2088).

- Gordis L, Gold LB. Epidemiology of pancreatic cancer. *World J Surg* 1984; 8: 808-21.
- Williamson RCN. Pancreatic cancer: the greatest oncological challenge. *BMJ* 1988; 296: 445-6.
- Trede M, Schwall G. The complications of pancreatotomy. *Ann Surg* 1988; 207: 39-47.
- Tsuchiya R, Noda T, Harada N, et al. Collective review of small carcinomas of the pancreas. *Ann Surg* 1986; 203: 77-81.
- Cubilla AL, Fitzgerald PT. Morphological lesions associated with human primary invasive non endocrine pancreatic cancer. *Cancer Res* 1976; 36: 2690-5.
- Allen-Merish TG. Pancreatic ductal mucinous hyperplasia: distribution within the pancreas, and effect of variation in ampullary and pancreatic duct anatomy. *Gut* 1988; 29: 1392-6.
- Longnecker DS, Crawford BG. Hyperplastic nodules and adenomas of exocrine pancreas in azaserine treated rats. *J Natl Cancer Inst* 1974; 53: 573-7.
- Pour P. Islet cells as a component of pancreatic ductal neoplasia. I. Experimental study: ductular cells, including islet cell precursors, as primary progenitor cells of tumors. *Am J Pathol* 1978; 90: 295-307.
- Pour P. Experimental pancreatic ductal (ductular) tumours. In: Moosa AR, eds. *Tumours of the pancreas*. Baltimore: Williams & Wilkins, 1980: 111.
- Ewing J. *Neoplastic diseases*. Philadelphia: Saunders, 1919.
- Pour PM, Bell RH. Alterations of pancreatic endocrine cell patterns and their secretions during pancreatic carcinogenesis in the hamster model. *Cancer Res* 1989; 49: 6396-400.
- Lebenthal E, Lev R, Lee PC. Prenatal and postnatal development of the human exocrine pancreas. In: Go VWL, ed. *The exocrine pancreas: biology, pathobiology and diseases*. New York: Raven Press, 1986: 33-43.
- Logothetopoulos J. Islet cell regeneration and neogenesis. In: Freinkel N, Renold AE, eds. *Handbook of physiology*. Section 7. Vol 2. Washington DC: American Physiological Society, 1972: 67-76.
- Gyr KE, Singer MV, Sarles H. Revised classification of pancreatitis, Marseille 1984. In: Gyr KE, Singer MV, Sarles H, eds. *Pancreatitis, concepts and classification*. Amsterdam: Excerpta Medica, 1984: xxiii-v.
- Solomon TE. Control of exocrine pancreatic secretion. In: Johnson LR, ed. *Physiology of the gastrointestinal tract*. 2nd ed. New York: Raven Press, 1987: 1173-207.
- Wessels NK. DNA synthesis, mitosis and differentiation in pancreatic acinar cells *in vitro*. *J Cell Biol* 1964; 20: 415-33.
- Pictet RL, Clark WR, Williams RM, Rutter WJ. An ultrastructural analysis of the developing embryonic pancreas. *Dev Biol* 1972; 29: 430-67.
- Wenzel G, Stocker E, Heine WD. Zur Zellproliferation in exokrinen Pankreas der Ratte. Autoradiographische Untersuchungen mit ³H-Thymidine. *Virchows Arch [B] Z* 1972; 10: 118-26.
- Greenberg RD, Dominguez A, Washington A, Holt PR. Impaired response of aging rat to caerulein-secretin stimulation [Abstract]. *Gastroenterology* 1986; 90: 1438.
- Poston GJ, Saydjari R, Lawrence JP, et al. Trophic effect of CCK, bombesin and pentagastrin on the aged rat pancreas [Abstract]. *Pancreas* 1989; 4: 637.
- Mainz DL, Parks NM, Webster PD. Effect of fasting and refeeding on pancreatic DNA synthesis and content. *Proc Soc Exp Biol Med* 1977; 156: 340-4.
- Brand SJ, Morgan RGH. The influence of starvation of intestinal cholecystokinin-like activity and pancreatic growth. *J Physiol* 1981; 321: 469-82.
- Brand SJ, Morgan RGH. Stimulation of pancreatic secretion and growth in the rat after feeding cholestyramine. *Gastroenterology* 1982; 83: 851-9.
- Goke B, Siegel EG, Stockmann F, et al. Effect of a specific serine protease inhibitor on the rat pancreas. *Digestion* 1985; 31: 128-35.
- McGuinness EE, Morgan RGH, Wormsley KG. Trophic effects on the pancreas of trypsin and bile salt deficiency in the small-intestinal lumen. *Scand J Gastroenterol* 1985; 20 (suppl 112): 64-7.
- Brannon PM, Demarest AS, Sabb JE, Korc M. Dietary modulation of epidermal growth factor action in cultured pancreatic acinar cells of the rat. *J Nutr* 1986; 116: 1306-15.
- Di Mattei E. Degli effetti della imitazione rugli elementi glandulari del pancreas. *Giornale dell reale. Accademia Medica Torino Seriale* 1885; 3.33: 476-7.
- Lehv MD, Fitzgerald PJ. Pancreatic acinar cell regeneration IV: regeneration after surgical resection. *Am J Pathol* 1968; 53: 513-35.
- Newman BM, Lee PC, Brooks S, et al. Enteral feedback control of pancreatic hypertrophy: the role of pancreatic biliary secretions. *Proc Soc Exp Biol Med* 1986; 181: 226-32.
- Laplace JP, Simoes Nunes C. Pancreatic size and enzyme content after vagal deafferentation in jejunectomized pigs under free or restricted feeding. *Gut* 1987; 28 (suppl): 169-73.
- Baba N, Chowdhury P, Inoue K, et al. Ileocaecal resection induced pancreatic growth in rats. *Peptides* 1985; 6: 211-5.
- Malfertheiner P, Buckler M, Ditschuneit H. Sequential changes in pancreatic growth following colectomy in rat [Abstract]. *Dig Dis Sci* 1985; 30: 983.
- Stock-Damge C, Aprahamian M, Lhoste E, et al. Pancreatic hyperplasia after small bowel resection in the rat: dissociation from endogenous gastric levels. *Digestion* 1984; 29: 223-30.
- Lawrence JP, Poston GJ, James E, et al. Hyperplasia of colon mucosa and pancreas after ileocaecal resection is not caused by gastrin. *Surgical Forum* 1988; 39: 177-9.
- Hakanson R, Blom H, Carlsson E, et al. Hypergastrinaemia produces trophic effects in stomach but not in pancreas and intestines. *Reg Pept* 1986; 13: 225-33.
- Enochs MR, Johnson LR. Trophic effects of gastrointestinal hormones: physiological implications. *Federation Proceedings* 1977; 36: 1942-7.
- Peterson H, Solomon T, Grossman MI. Effect of chronic pentagastrin, cholecystokinin and secretin on pancreas of rats. *Am J Physiol* 1978; 234: E286-93.
- Solomon T, Vanier M, Morisset J. Cell site and time course of DNA synthesis in pancreas after cerulein and secretin. *Am J Physiol* 1983; 245: G99-105.
- Haarstad H, Steffensud S, Winnberg A, Petersen H. The trophic effect on the pancreas of long term continuous intravenous infusion of secretin and a cholecystokinin-like peptide in rats. *Scand J Gastroenterol* 1986; 21: 589-97.
- Balas D, Senegas-Balas F, Pradayrol L, et al. Long-term comparative effect of cholecystokinin and gastrin on mouse stomach, antrum, intestine and exocrine pancreas. *Am J Anat* 1985; 174: 27-43.
- Hudd C, La Regina MC, Devine JE, Johnson FE. Response of nude mouse pancreas and gastrointestinal tract to gradual doses of exogenous cholecystokinin. *Curr Surg* 1986; 43: 105-9.
- Solomon TE. Trophic effects of pentagastrin on gastrointestinal tract in fed and fasted rats. *Gastroenterology* 1986; 91: 108-16.
- Niederer C, Liddle RA, Williams JA, Grendell JH. Pancreatic growth: interaction of exogenous cholecystokinin, a protease inhibitor, and a cholecystokinin receptor antagonist in mice. *Gut* 1987; 28 (suppl): 63-9.
- Goke B, Printz H, Koop I, et al. Endogenous CCK release and pancreatic growth in rats after feeding a proteinase inhibitor (Camostat). *Pancreas* 1986; 1: 509-15.
- Wisner JR, McLaughlin RE, Rich KA. Effects of L-364, 718, a new cholecystokinin receptor antagonist, on Camostat-induced growth of the rat pancreas. *Gastroenterology* 1988; 94: 109-13.
- Lee PC, Newman BM, Praissman M, et al. Cholecystokinin: a factor responsible for the enteral feedback control of pancreatic hypertrophy. *Pancreas* 1986; 1: 335-40.
- Levan VM, Liddle RA, Green GM. Jejunal bypass stimulation of pancreatic growth and cholecystokinin secretion in rats: importance of luminal nutrition. *Gut* 1987; 28 (suppl): 25-9.
- Folsch UR, Schafmayer A, Ebur R, et al. Elevated plasma cholecystokinin concentrations in exocrine pancreatic atrophy in the rat. *Digestion* 1984; 29: 60-4.
- Steinberg WM, Burns MK, Henry JP, et al. Cerulein induces hyperplasia of the pancreas in a rat model of chronic pancreatic insufficiency. *Pancreas* 1987; 2: 176-80.
- Majumdar APN, Vesenka GD, Dubick MA. Acceleration of pancreatic regeneration by cholecystokinin in rats. *Pancreas* 1987; 2: 199-204.
- Feurle GE, Muller B, Ohnheiser G, Baca I. Action of neurotensin on size, composition and growth of pancreas

- and stomach in the rat. *Regul Pept* 1985; **13**: 53–62.
- 52 Feurle GE, Muller B, Rix E. Neurotensin induces hyperplasia of the pancreas and growth of the gastric antrum in rats. *Gut* 1987; **28** (suppl): 19–23.
 - 53 Lehy T, Puccio F, Chariot J, Labeille D. Stimulatory effect of bombesin on the growth of gastrointestinal tract and pancreas in suckling rats. *Gastroenterology* 1986; **90**: 1942–9.
 - 54 Upp JR, Poston GJ, MacLellan DG, *et al.* Mechanism of the trophic action of bombesin on the pancreas. *Pancreas* 1988; **3**: 193–8.
 - 55 Dembinski A, Gregory H, Konturek SJ, Pulanski M. Trophic action of epidermal growth factor on the pancreas and gastroduodenal mucosa in rats. *J Physiol* 1982; **325**: 35–40.
 - 56 Morisset J, Geilik P, Lord A, Solomon TE. Effect of chronic administration of somatostatin on rat exocrine pancreas. *Regul Pept* 1982; **4**: 49–58.
 - 57 Morisset J, Jolicœur L. Effect of hydrocortisone on pancreatic growth in rats. *Am J Physiol* 1980; **239**: G95–8.
 - 58 Morisset J, Jolicœur L, Genik P, Lord A. Interaction of hydrocortisone and caerulein on pancreatic size and composition in rats. *Am J Physiol* 1981; **241**: G37–42.
 - 59 Dembinski A, Konturek SJ. Effects of E, F and I series prostaglandins and analogues on growth of gastroduodenal mucosa and pancreas. *Am J Physiol* 1985; **248**: G170–5.
 - 60 Logsdon CD, Williams JA. Pancreatic acinar cells in monolayer culture: direct trophic effects of caerulein in vitro. *Am J Physiol* 1986; **250**: G440–7.
 - 61 Logsdon CD. Effects of calcium mediated secretagogues on the growth of pancreatic acinar cells in vitro. *Gut* 1987; **28** (suppl): 117–20.
 - 62 Logsdon CD. Growth and differentiation of pancreatic acinar cells in primary culture [Abstract]. *Pancreas* 1989; **4**: 259.
 - 63 Githens S, Holmgeist DRG, Whelan JF, Ruby JR. Characterization of ducts isolated from the pancreas of the rat. *J Cell Biol* 1980; **85**: 122–35.
 - 64 Githens S, Finley JJ, Patke CL, *et al.* Biochemical and histochemical characterization of cultured rat and hamster pancreatic ducts. *Pancreas* 1987; **4**: 427–38.
 - 65 Hubchak S, Mangino MM, Reddy K, Scarpelli DG. Characterization of syrian golden hamster pancreatic duct cells maintained in extended monolayer culture [Abstract]. *Pancreas* 1989; **4**: 620.
 - 66 Williams JA. Role of receptors in mediating trophic stimuli in the pancreas. *Gut* 1987; **28** (suppl): 45–9.
 - 67 Goldfine ID, Williams JA. Receptors for insulin and CCK in the acinar pancreas: relationship to hormone action. *Int Rev Cytol* 1983; **85**: 1–38.
 - 68 Logsdon CD, Williams JA. Intracellular Ca^{2+} and phorbol esters synergistically inhibit EGF intermediaries in pancreatic acini. *Biochem J* 1984; **223**: 893–900.
 - 69 Mossner J, Logsdon CD, Petau N, *et al.* Effect of intracellular Ca^{2+} in insulin like growth factor II internalization into pancreatic acini: role of insulin and cholecystokinin. *J Biol Chem* 1984; **259**: 12350–6.
 - 70 Mossner J, Logsdon CD, Goldfine ID, Williams JA. Regulation of pancreatic acinar cell insulin receptors by insulin. *Am J Physiol* 1984; **247**: G155–60.
 - 71 Hootman SR, Brown ME, Williams JA, Logsdon CD. Regulation of muscarinic acetyl choline receptors in cultured guinea pig pancreatic acini. *Am J Physiol* 1986; **251**: G75–83.
 - 72 Band P, Richardson SB, Boctor AM, Grossman A. Somatostatin enhances binding of [3H]estradiol to a cytosolic protein in rat pancreas. *J Biol Chem* 1983; **258**: 7284–7.
 - 73 Matozaki T, Saramoto C, Nagao M, Baba S. Phorbol ester or diacylglycerol modulates somatostatin binding to its receptors on rat pancreatic acinar cell membranes. *J Biol Chem* 1986; **261**: 1414–20.
 - 74 Otsuki M, Williams JA. Altered amylase secretion by isolated pancreatic acini following in-vivo administration of cholecystokinin. *Am J Physiol* 1983; **244**: G683–8.
 - 75 Leung YK, Lee PC, Lebenthal E. Maturation of cholecystokinin receptors in pancreatic acini of rats. *Am J Physiol* 1986; **250**: G594–7.
 - 76 Poston GJ, Singh P, Dravian ED, *et al.* Development and age related changes in pancreatic cholecystokinin receptors and duodenal cholecystokinin in guinea pigs. *Mech Ageing Develop* 1988; **46**: 59–66.
 - 77 Williams JA, Hootman SR. Stimulus-secretion coupling in pancreatic acinar cells. In: Go VLW, ed. *The exocrine pancreas: biology, pathophysiology and disease*. New York: Raven Press, 1986: 123–39.
 - 78 Williams JA, Goldfine ID. The insulin pancreatic acinar axis. *Diabetes* 1985; **34**: 980–6.
 - 79 Liebow C, Reilly C, Serrano M, Schally AV. Somatostatin analogues inhibit growth of pancreatic cancer by stimulating tyrosine phosphatase. *Proc Natl Acad Sci USA* 1989; **86**: 2003–7.
 - 80 Sandberg AA, Rosenthal HE. Estrogen receptors in the pancreas. *J Steroid Biochem* 1974; **5**: 969–77.
 - 81 Sandberg AA, Rosenthal HE. Steroid receptors in exocrine glands: the pancreas and prostate. *J Steroid Biochem* 1979; **11**: 293–9.
 - 82 Pousette A, Carlstrom K, Skoldefors H, *et al.* Purification and partial characterization of a 17-beta estradiol-binding macromolecule in the human pancreas. *Cancer Res* 1982; **42**: 633–7.
 - 83 Boctor AM, Band P, Grossman A. Analysis of binding of [3H] estradiol to the cytosol fraction of rat pancreas: comparison with sites in the cytosol of uterus. *Endocrinology* 1983; **113**: 453–62.
 - 84 Grossman A, Richardson SB, Altszuler N, Lane B. Evidence that the [3H] estradiol binding protein in pancreas is localized in exocrine cells. *Endocrinology* 1985; **116**: 2221–6.
 - 85 Pousette A, Fernstad R, Theve N-O, *et al.* Further characterization of the estrogen binding macromolecule in human pancreas. *J Steroid Biochem* 1985; **23**: 115–9.
 - 86 Petterson A, Lea O, Bakkerold K, Arnesjo B. Progress in pancreatic oestrogen receptors. *Eur J Surg Oncol* 1986; **12**: 321–4.
 - 87 Boctor AM, Band P, Grossman A. Requirement for an accessory factor for binding of [3H] estradiol to protein in the cytosol fraction of rat pancreas. *Proc Natl Acad Sci USA* 1981; **78**: 5648–51.
 - 88 Grossman A. Steroid receptors in normal pancreas [Abstract]. *Pancreas* 1989; **4**: 265.
 - 89 Gorelick FS. Second messenger systems and adaptation. *Gut* 1987; **28** (suppl): 79–84.
 - 90 Rasmussen HR. The calcium messenger system: part 1. *New Engl J Med* 1986; **314**: 1095–101.
 - 91 Rasmussen HR. The calcium messenger system: part 2. *New Engl J Med* 1986; **314**: 1164–70.
 - 92 Hannun YA, Carson RL, Bell RM. Protein kinase C activation in mixed micelles. *J Biol Chem* 1986; **261**: 7184–90.
 - 93 Nishizuka Y. The role of protein kinase C in cell surface signal transduction and tumour promotion. *Nature* 1984; **308**: 693–8.
 - 94 Nishizuka Y. Studies and perspectives of protein kinase C. *Science* 1986; **233**: 305–12.
 - 95 Roos W, Fabbro D, Kung W, *et al.* Correlation between hormone dependency and the regulation of epidermal growth factor receptor by tumor promoters in human mammary carcinoma lines. *Proc Natl Acad Sci USA* 1986; **83**: 991–5.
 - 96 May WS, Sayhoun N, Jacobs S, *et al.* Mechanism of phorbol diester-induced regulation of surface transferrin receptor involves the action of activated protein kinase C and intact cytoskeleton. *J Biol Chem* 1986; **260**: 9419–26.
 - 97 Means AR, Dedman JR. Calmodulin – an intracellular calcium receptor. *Nature* 1980; **285**: 73–7.
 - 98 Chafouleas JG, Pardue RL, Brinkley BR, *et al.* Regulation of intracellular levels of calmodulin and tubulin in normal and transformed cells. *Proc Natl Acad Sci USA* 1981; **78**: 996–1000.
 - 99 Sibley DR, Lefkowitz RJ. Molecular mechanisms of receptor desensitisation using the beta-adrenergic receptor-coupled adenylate cyclase system as a model. *Nature* 1985; **317**: 124–9.
 - 100 Gilman AG. G proteins and dual control of adenylate cyclase. *Cell* 1984; **36**: 577–9.
 - 101 Grossman MJ, Greengard H, Ivy AC. The effect of dietary composition on pancreatic enzymes. *Am J Physiol* 1943; **138**: 676–82.
 - 102 Schick J, Verspohl R, Kern HF, Scheele GA. Two distinct adaptive responses in the synthesis of exocrine pancreatic enzymes to inverse changes in proteins and carbohydrates in the diet. *Am J Physiol* 1984; **247**: G611–6.
 - 103 Kern HF, Rausch U, Scheele GA. Regulation of gene expression in pancreatic adaptation to nutritional substrate or hormones. *Gut* 1987; **28** (suppl): 89–94.
 - 104 Darnell IE. Variety in the level of gene control in eukaryotic cells. *Nature* 1982; **297**: 365–71.
 - 105 Dynan WS, Tjian R. Control of eukaryotic messenger RNA synthesis by sequence specific DNA-binding proteins. *Nature* 1985; **316**: 774–8.
 - 106 Scheele GA. Regulation of gene expression in the exocrine pancreas. In: Go VLW, ed. *The exocrine pancreas: biology, pathology and disease*. New York: Raven Press, 1986: 55–67.
 - 107 Scheidereit C, Greisse S, Westphal HW, Beato M. The glucocorticoid receptor binds to defined nucleotide sequences near the promoter of mouse mammary tumour tissue. *Nature* 1983; **304**: 749–52.
 - 108 Wicker C, Puigserver A, Rausch U, *et al.* Multiple level caerulein control of the gene expression of secretory proteins in the rat pancreas. *Eur J Biochem* 1985; **151**: 461–6.
 - 109 Cochran AB. The molecular action of platelet derived growth factor. *Adv Cancer Res* 1985; **45**: 183–216.
 - 110 Armelin HA, Armelin MCS, Kelly K, *et al.* Functional role for C-myc in mitogenic response to platelet derived growth factor. *Nature* 1984; **310**: 655–60.
 - 111 Cochran BH, Refell AL, Stiles CD. Molecular cloning of gene sequences regulated by platelet derived growth factor. *Cell* 1983; **33**: 939–47.
 - 112 Kelly K, Cochran BH, Stiles CD, Leder P. Specific regulation of the C-myc gene by lymphocyte mitogens and platelet derived growth factor. *Cell* 1983; **35**: 603–10.
 - 113 Chambard JC, Franchi A, Le Cam A, Pouyssegur J. Growth factor stimulated protein phosphorylation in G0-G1-arrested fibroblasts. *J Biol Chem* 1983; **258**: 1706–13.
 - 114 Pegg AE, McCann PP. Polyamine metabolism and function. *Am J Physiol* 1982; **243**: C212–21.
 - 115 Pegg AE. Recent advances in the biochemistry of polyamines in eukaryotes. *Biochem J* 1986; **234**: 249–62.
 - 116 Tabor CW, Tabor H. Polyamines. *Ann Rev Biochem* 1984; **53**: 749–90.
 - 117 Scalabrino G, Ferioli ME. Polyamines in mammalian tumors. Part I. *Adv Cancer Res* 1981; **35**: 152–268.
 - 118 Janne J, Poso H, Raina A. Polyamines in rapid growth and cancer. *Biochim Biophys Acta* 1978; **473**: 241–93.
 - 119 McCann PP. Regulation of ornithine decarboxylase in eukaryotic cells. In: Gangas JM, ed. *Polyamines in biomedical research*. New York: Wiley, 1980: 109–23.
 - 120 Dowling RH, Hosomi M, Stace NM, *et al.* Hormones and polyamines in intestinal and pancreatic adaptation. *Scand J Gastroenterol* 1985; **20** (suppl 112): 84–95.

- 121 Baylin SB, Stevens SA, Shakir KMM. Association of diamine oxydase and ornithine decarboxylase with maturing cells in rapidly proliferating epithelium. *Biochim Biophys Acta* 1978; 541: 415-9.
- 122 Morisset J, Benrezzak O. Polyamines and pancreatic growth induced by caerulein. *Life Sci* 1984; 35: 2471-80.
- 123 Scalabrino G, Ferioli ME. Polyamines in mammalian tumors. Part II. *Adv Cancer Res* 1982; 36: 1-102.
- 124 Benrezzak O, Morisset J. Effects of alpha-difluoromethyl-ornithine on pancreatic growth induced by caerulein. *Regul Pept* 1984; 9: 143-53.
- 125 Morisset J, Benrezzak O. Reversal of alpha dimethyl fluoro-ornithine inhibition of caerulein induced pancreatic growth by putrescine. *Regul Pept* 1985; 11: 201-8.
- 126 Danzin C, Claverie N, Wagner J, et al. Comparison of 3-isobutyl-1-methylxanthine-induced accumulation of putrescine in rat pancreas and liver. *Life Sci* 1983; 33: 2173-8.
- 127 Fontham E, Corresa P, Colin I. Epidemiology of cancer of the pancreas. In: Howard IM, Jordan GL, Reber HA, eds. *Surgical diseases of the pancreas*. Philadelphia: Lea & Febiger, 1984: 613-26.
- 128 Gudjonsson B, Linstone EM, Spiro HM. Cancer of the pancreas: diagnostic accuracy and survival statistics. *Cancer* 1978; 42: 2494-506.
- 129 Poston GJ, Williamson RCN. Causes, diagnosis and management of exocrine pancreatic cancer. *Compr Ther* 1990; 16: 36-42.
- 130 Longnecker DS. Interface between adaptive and neoplastic growth in the pancreas. *Gut* 1987; 28 (suppl): 253-8.
- 131 Longnecker DS, French J, Hyde E, et al. Effects of age on nodule induction by azaserine. *J Natl Cancer Inst* 1977; 58: 1769-75.
- 132 Rao M, Subbarao M, Leuteke N, Scarpelli D. Further characterization of carcinogen-induced hepatocyte like cells in hamster pancreas. *Am J Pathol* 1983; 110: 89-94.
- 133 Stace NH, Palmer TJ, Vaja S, Dowling RH. Long term pancreaticobiliary diversion stimulates hyperplastic and adenomatous nodules in the rat pancreas: a new model for spontaneous tumour formation. *Gut* 1987; 28 (suppl): 265-8.
- 134 Stewart I, Flaks B, Cooper MJ, et al. Precancerous lesions in the pancreas after short-term pancreaticobiliary diversion [Abstract]. *Br J Surg* 1988; 75: 1257.
- 135 Longnecker DS, Morgan R. Diet and cancer of the pancreas: epidemiological and experimental evidence. In: Reddy B, et al. *Diet, nutrition and cancer: a critical evaluation*. New York: CRC Press, 1986: 12-22.
- 136 Miazza BM, Widgren S, Chayvialle JA, et al. Exocrine pancreatic nodules after long term pancreaticobiliary diversion in rats. An effect of raised CCK plasma concentrations. *Gut* 1987; 28 (suppl): 269-74.
- 137 Roebuck BD. Effect of CCK in the rat model of pancreatic carcinogenesis. *Pancreas* 1989; 4: 259-60.
- 138 Roebuck BD, Longnecker DS, Baumgartner KJ, Throw CD. Carcinogen induced lesions in the rat pancreas: effects of varying levels of essential fatty acids. *Cancer Res* 1985; 45: 5252-6.
- 139 Ura H, Makino T, Ito S, et al. Combined effects of cholecystectomy and lithocholic acid on pancreatic carcinogenesis of N-nitrosobis (2-hydroxypropyl) amine in Syrian golden hamsters. *Cancer Res* 1986; 46: 4782-6.
- 140 Goustin AS, Leof EB, Shipley GD, Moses HL. Growth factors and cancer. *Cancer Res* 1986; 46: 1015-29.
- 141 Townsend CM Jr, Beauchamp RD, Singh P, Thompson JC. Growth factors and intestinal neoplasms. *Am J Surg* 1988; 155: 526-36.
- 142 McGuinness EE, Morgan RGH, Levison DA, et al. The effect of long term feeding of soya flour on the rat pancreas. *Scand J Gastroenterol* 1980; 15: 497-502.
- 143 McGuinness EE, Morgan RGH, Wormsley KG. Fate of pancreatic nodules induced by raw soya flour in rats. *Gut* 1987; 28 (suppl): 207-12.
- 144 Pour PM, Hegelson S, Lawson T, et al. Effect of cholecystokinin on pancreatic carcinogenesis in the hamster model. *Carcinogenesis* 1988; 9: 597-601.
- 145 Pour PM. Effect of CCK in the hamster model [Abstract]. *Pancreas* 1989; 4: 260.
- 146 Johnson FE, La Regina MC, Martin SA, Bajniti HM. Cholecystokinin inhibits pancreatic and hepatic carcinogenesis. *Cancer Detect Prev* 1983; 6: 389-402.
- 147 Satake K, Mukai R, Kato Y, Umeyama K. Effect of caerulein on the normal pancreas and an experimental pancreatic carcinoma in the Syrian golden hamster. *Pancreas* 1986; 1: 246-53.
- 148 Howatson AG, Carter DC. Pancreatic carcinogenesis enhancement by cholecystokinin in the hamster-nitrosamine model. *Br J Cancer* 1985; 51: 107-14.
- 149 Howatson AG, Carter DC. Pancreatic carcinogenesis: effect of secretin in the hamster-nitrosamine model. *J Natl Cancer Inst* 1987; 78: 101-5.
- 150 Hugh T, Alexander RW, Upp JR, et al. Caerulein inhibits growth of a human cholangiocarcinoma with cholecystokinin receptors [Abstract]. *Gastroenterology* 1987; 92: 1801.
- 151 Chester JF, Gaissert MA, Ross JA, Malt RA. Pancreatic cancer in the Syrian hamster induced by N-nitroso bis (2-oxopropyl)-amine: carcinogenic effect with epidermal growth factor. *Cancer Res* 1986; 46: 2954-70.
- 152 Paz-Bouza JI, Redding TW, Schally AV. Treatment of nitrosamine-induced pancreatic tumors in hamsters with analogues of somatostatin and luteinizing hormone-releasing hormone. *Proc Natl Acad Sci USA* 1987; 84: 1112-6.
- 153 Zalutnai A, Schally AV. Treatment of the N-nitrosobis(2-oxopropyl) amine-induced pancreatic cancer in Syrian golden hamsters with D-Trp-6-LH-RH and somatostatin analogue RC-160 microcapsules. *Cancer Res* 1989; 49: 1810-15.
- 154 Zalutnai A, Schally AV. Responsiveness of the hamster pancreatic cancer to treatment with microcapsules of d-Trp-6-LH-RH and somatostatin analogue RC-160. *Int J Pancreatol* 1989; 4: 149-60.
- 155 Townsend CM Jr, Franklin RB, Watson LC, et al. Stimulation of pancreatic cancer growth by caerulein and secretin. *Surg Forum* 1981; 32: 228-30.
- 156 Estival A, Clemente F, Ribet J. Adenocarcinoma of the human exocrine pancreas: presence of secretin and caerulein receptors. *Biochem Biophys Res Commun* 1981; 102: 1336-41.
- 157 Alexander RW, Upp JR Jr, Singh P, et al. Asperlicin inhibits the growth of a xenografted human pancreatic carcinoma [Abstract]. *Gastroenterology* 1987; 92: 1293.
- 158 Douglas BR, Worters RA, Jansen JBM, et al. Influence of cholecystokinin antagonist on the effects of cholecystokinin and bombesin on azaserine induced lesions on rat pancreas. *Gastroenterology* 1989; 96: 462-9.
- 159 Upp JR Jr, Singh P, Townsend CM Jr, Thompson JL. Predicting response to endocrine therapy in human pancreatic cancer with cholecystokinin receptors [Abstract]. *Gastroenterology* 1987; 92: 1677.
- 160 Stock-Damge C, Lhoste E, Aprahamian M, Pousse A. Effect of chronic bombesin on pancreatic size, composition and secretory function in the rat. *Gut* 1987; 28 (suppl): 1-7.
- 161 Alexander RW, Upp JR, Jr, Poston GJ, et al. Bombesin inhibits growth of human pancreatic adenocarcinoma in nude mice. *Pancreas* 1988; 3: 297-302.
- 162 McArthur KE, Wood CL, O'Dorisio MS, et al. Characterization of receptors for VIP on pancreatic acinar cell plasma membrane using covalent cross linking. *Am J Physiol* 1987; 252: G404-12.
- 163 Yao CZ, Poston GJ, Townsend CM Jr, Thompson JC. Covalent cross linking identification of vasoactive intestinal receptors in human colon and pancreatic cancers and hamster pancreatic cancer [Abstract]. *Gastroenterology* 1987; 92: 1702.
- 164 Estival A, Mounielou P, Trocheris V, et al. Presence of VIP reception in a human pancreatic adenocarcinoma cell line. Modulations of the cAMP response during cell proliferation. *Biochem Biophys Res Commun* 1983; 111: 958-63.
- 165 Fernandez-Moreno MD, Diaz-Juarez JL, Arilla E, Prieto JC. Effect of resection of small intestine in the interaction of vasoactive intestinal peptide with rat colonic epithelial cells. *Horm Metab Res* 1985; 17: 289-92.
- 166 De Rubertis FR, Craven PA. Early alteration in rat colonic mucosal cyclic nucleotide metabolism and protein kinase activity induced by 1,2-dimethylhydrazine. *Cancer Res* 1980; 40: 4589-98.
- 167 Craven PA, De Rubertis FR. Cyclic nucleotide metabolism in rat colonic epithelial cells with different proliferative activities. *Biochim Biophys Acta* 1981; 676: 155-69.
- 168 Poston GJ, Yao CZ, Upp JR Jr, et al. Vasoactive intestinal peptide inhibits the growth of hamster pancreatic cancer but not human pancreatic cancer in vivo. *Pancreas* 1988; 3: 439-43.
- 169 Kraenzlin ME, Ch'ng JLL, Wood SM, et al. Long-term treatment of a vipoma with somatostatin analogue resulting in remission of symptoms and possible shrinkage of metastases. *Gastroenterology* 1985; 88: 185-7.
- 170 Clements D, Elias E. Regression of metastatic vipoma with somatostatin analogue SMS 201-995. *Lancet* 1985; i: 874-5.
- 171 Reubi JC. A somatostatin analogue inhibits chondrosarcoma and insulinoma tumour growth. *Acta Endocrinol* 1985; 109: 108-14.
- 172 Upp JR Jr, Alexander RW, Poston GJ, et al. Inhibition of a human malignant carcinoid growing in nude mice [Abstract]. *Gastroenterology* 1987; 92: 1676.
- 173 Smith JP, Solomon TE. Effects of gastrin, proglumide and somatostatin on growth of human colon cancer. *Gastroenterology* 1988; 95: 1541-8.
- 174 Schally AV, Comaro-Schally M, Redding T. Antitumor effects of analogues of hypothalamic hormones in endocrine dependent cancers. *Proc Soc Exp Biol Med* 1984; 175: 259-81.
- 175 Schally AV, Cai R-Z, Torres-Aleman I, et al. Endocrine, gastrointestinal and antitumor activity of somatostatin analogues. In: Moody TW, ed. *Neural and endocrine peptides and receptors*. New York: Plenum, 1986: 73-88.
- 176 Schally AV. Oncological applications of somatostatin analogues. *Cancer Res* 1988; 48: 6977-85.
- 177 Redding TW, Schally AV. Inhibition of growth of pancreatic carcinomas in animal models by analogues of hypothalamic hormones. *Proc Natl Acad Sci USA* 1984; 81: 248-52.
- 178 Upp JR Jr, Olson D, Poston GJ, et al. Inhibition of growth of two human pancreatic adenocarcinomas in vivo by somatostatin analogue SMS 201-995. *Am J Surg* 1988; 155: 29-35.
- 179 Poston GJ, Townsend CM Jr, Rajaraman S, et al. Effect of somatostatin and tamoxifen on the growth of human pancreatic cancers in nude mice. *Pancreas* 1990; 5: 151-7.
- 180 Szende B, Zalutnai A, Schally AV. Programmed cell death (apoptosis) in pancreatic cancer of hamsters after administration of analogs of luteinizing hormone releasing hormone and somatostatin. *Proc Natl Acad Sci USA* 1989; 86: 1643-7.
- 181 Liebow C, Hierowski M, Du Sapin K. Hormonal control of pancreatic cancer growth. *Pancreas* 1986; 1: 44-8.
- 182 Fekete M, Zalutnai A, Comaru-Schally A-M, Schally AV.

- Membrane receptors for peptides in experimental and human pancreatic cancers. *Pancreas* 1989; 4: 521-8.
- 183 Schlegel N, Hawes RF, Raptis S, et al. Inhibition of cholecystokinin-pancreozymin release by somatostatin. *Lancet* 1977; i: 166-8.
 - 184 Viguerie N, Esteve JP, Tahiri-Jouti N, et al. Functional somatostatin receptors in a rat pancreatic acinar cell line [Abstract]. *Gastroenterology* 1987; 92: 1660.
 - 185 Osei K, O'Dorisio TM. Malignant insulinoma: effects of a somatostatin analogue (compound 201-996) on serum glucose, growth and gastro-entero-pancreatic hormones. *Ann Intern Med* 1985; 103: 223-5.
 - 186 Malt RA, Chester JF, Gaissert HA, Ross JS. Augmentation of chemically induced pancreatic and bronchial cancers by epidermal growth factor. *Gut* 1987; 28 (suppl): 249-52.
 - 187 Velu TJ, Beguinot L, Vass WC, et al. Epidermal growth factor-dependent transformation by a human EGF receptor proto-oncogene. *Science* 1987; 238: 1408-10.
 - 188 Kagayama R, Merlino GT, Pastan I. Epidermal growth factor (EGF) receptor gene transcription. *J Biol Chem* 1988; 263: 6329-36.
 - 189 Kagayama R, Merlino GT, Pastan I. A transcription factor active on the epidermal growth factor receptor gene. *Proc Natl Acad Sci USA* 1988; 85: 5016-20.
 - 190 Johnson AC, Ishii S, Jinno Y, Pastan I, Merlino GT. Epidermal growth factor receptor gene promoter. *J Biol Chem* 1988; 263: 5693-9.
 - 191 Korc M, Magun BE. Recycling of epidermal growth factor in a human pancreatic carcinoma cell line. *Proc Natl Acad Sci USA* 1985; 82: 6172-5.
 - 192 Clark AJL, Ishii S, Richert N, et al. Epidermal growth factor regulates the expression of its own receptor. *Proc Natl Acad Sci USA* 1985; 82: 8374-8.
 - 193 Earp HS, Austin KS, Blaisdell J, et al. Epidermal growth factor (EGF) stimulates EGF receptor synthesis. *J Biol Chem* 1986; 261: 4777-80.
 - 194 Hierowski MT, Liebow C, du Sabin K, Schally AV. Stimulation by somatostatin of dephosphorylation of membrane proteins in pancreatic cancer MIA PaCa-2 cell lines. *FEBS Lett* 1985; 179: 252-6.
 - 195 Rall LB, Scott J, Bell GJ, et al. Mouse prepro-epidermal growth factor synthesis by kidney and other tissues. *Nature* 1985; 313: 228-31.
 - 196 Sporn MB, Roberts AB. Autocrine growth factors and cancer. *Nature* 1985; 313: 745-7.
 - 197 Weinberg RA. The action of oncogenes in the cytoplasm and nucleus. *Science* 1985; 230: 770-6.
 - 198 Downward J, Yarden Y, Mayes E, et al. Close similarity of epidermal growth factor receptor and V-erb-B oncogene protein sequences. *Nature* 1984; 307: 521-7.
 - 199 Takeuchi T, Gumucio DL, Yamada T, et al. Genes encoding pancreatic polypeptide and neuropeptide Y are on human chromosomes 17 and 7. *J Clin Invest* 1986; 77: 1038-41.
 - 200 Korc M, Meltzer P, Trent J. Enhanced expression of epidermal growth factor receptor correlates with alterations of chromosome 7 in human pancreatic cancer. *Proc Natl Acad Sci USA* 1986; 83: 5141-4.
 - 201 Korc M, Shadle KM, Meltzer PS. Analysis of EGF receptor gene structure in human pancreatic cancer cells [Abstract]. *Dig Dis Sci* 1987; 32: 1173.
 - 202 Korc M. EGF and pancreatic cancer. In: Thompson JL, ed. *Proceedings second Galveston international symposium on gastrointestinal hormones*. New York: McGraw-Hill (in press).
 - 203 De Larco JE, Todaro GJ. Growth factors from urine sarcoma virus transformed cells. *Proc Natl Acad Sci USA* 1978; 75: 4001-5.
 - 204 Roberts AB, Frolik C, Anzano MA, Sporn MB. Transforming growth factors from neoplastic and non neoplastic tissue. *Federation proceedings* 1983; 42: 2621-6.
 - 205 Korc M. Regulation of pancreatic cancer cell proliferation by an autocrine cycle coupled to the epidermal growth factor receptor [Abstract]. *Pancreas* 1989; 4: 262-3.
 - 206 Roberts AB, Anzano MA, Meyers CA, et al. Purification and properties of a type B transforming growth factor from bovine kidney. *Biochemistry* 1983; 22: 5692-8.
 - 207 Frolik CA, Dart LL, Meyers CA, et al. Purification and initial characterization of a type B transforming growth factor from human placenta. *Proc Natl Acad Sci USA* 1983; 80: 3676-80.
 - 208 Moses HL, Childs CB, Halper J, et al. Role of transforming growth factors in neoplastic transformations. In: Veneziale CM, ed. *Control of cell growth and proliferation*. New York: Van Nostrand Reinhold, 1984: 147-67.
 - 209 Leof EB, Proper JA, Goustin AS, et al. Induction of C-sis and m RNA and activity similar to platelet derived growth factor beta: a proposed model for indirect mitogenesis involving autocrine activity. *Proc Natl Acad Sci USA* 1986; 83: 2453-7.
 - 210 Moses HL, Tucker RF, Leof EB, Derynck R. Type beta transforming growth factor is a growth stimulant and growth inhibitor. In: Feremisco J, et al, eds. *Growth factors and transformation*. Cold Spring Harbor NY: Cold Spring Harbor Laboratory, 1985: 65-71.
 - 211 Moses HL. Positive and negative control by growth factors [Abstract]. *Pancreas* 1989; 4: 261.
 - 212 Leof EB, Proper JA, Moses HL. Modulation of transforming growth factor type beta action by activated ras and c-myc. *Mol Cell Biol* 1987; 7: 2649-52.
 - 213 Perroteau I, Salomon D, De Bortoli M, et al. Immunological detection and quantification of alpha transforming growth factors in human breast carcinoma cells. *Breast Cancer Research and Treatment* 1986; 7: 201-10.
 - 214 Lippman ME, Dickson RB, Bates S, et al. Autocrine and paracrine growth regulation of human breast cancer. *Breast Cancer Research and Treatment* 1986; 7: 59-70.
 - 215 Coffey RJ Jr, Shipley GD, Moses HL. Production of transforming growth factors by human colon cancer lines. *Cancer Res* 1986; 46: 1164-9.
 - 216 Smith JJ, Derynck R, Korc M. Production of transforming growth factor alpha in human pancreatic cancer cells: evidence of a superagonist autocrine cycle. *Proc Natl Acad Sci USA* 1987; 84: 7567-70.
 - 217 Sandberg AA. Estrogens and pancreatic cancer: some recent aspects. *Scand J Gastroenterol* 1986; 21: 129-33.
 - 218 Greenway BA. Carcinoma of the exocrine pancreas: a sex hormone responsive tumour? *Br J Surg* 1987; 74: 441-2.
 - 219 Tesone M, Chazenbalk D, Ballesos G, Charreau EH. Estrogen receptors in rat pancreatic islets. *J Steroid Biochem* 1979; 11: 1309-14.
 - 220 Greenway B, Iqbal MJ, Johnson PJ, Williams R. Oestrogen receptor proteins in malignant and fetal pancreas. *BMJ* 1981; 283: 751-3.
 - 221 Kiang DT, Kennedy BJ. Estrogen receptor assay in differential diagnosis of adenocarcinomas. *JAMA* 1977; 238: 32-4.
 - 222 Molteni A, Rao MS, Reddy JK, Fors EM. Estradiol receptors in the transplantable pancreatic carcinoma of the rat [Abstract]. *Federation Proceedings* 1978; 37: 897.
 - 223 Andren-Sendberg A, Borg A, Dawiskiba S, et al. Estrogen receptors and estrogen binding proteins in pancreatic cancer [Abstract]. *Digestion* 1982; 25: 12.
 - 224 Stedman KE, Moore GE, Morgan RT. Estrogen receptor protein in diverse human tumors. *Arch Surg* 1980; 115: 244-8.
 - 225 Satake K, Yoshimori T, Mukai R, Umeyama K. Estrogen receptors in 7,12-dimethyl benz(a)anthracene (DMBA) induced pancreatic carcinoma in rats and human pancreatic carcinoma. *Clin Oncol* 1982; 8: 49-54.
 - 226 Pousette A. Demonstration of an androgen receptor in rat pancreas. *Biochem J* 1976; 157: 229-32.
 - 227 Corbishley TP, Iqbal MJ, Wilkinson ML, Williams R. Androgen receptors in human normal and malignant pancreatic tissue and cell lines. *Cancer* 1986; 57: 1992-5.
 - 228 Gullo L, Priori P, Laro G. Influence of adrenal cortex on exocrine pancreatic function. *Gastroenterology* 1982; 83: 92-6.
 - 229 Kohen A, Kulka RG. Induction of chymotrypsinogen by hydrocortisone in embryonic chick pancreas in vitro. *J Biol Chem* 1974; 249: 4522-7.
 - 230 Andriulli A, Tappero GF, Gaia E, et al. Sex related differences in exocrine pancreatic secretion [Abstract]. *Digestion* 1982; 25: 13.
 - 231 Wilking N, Appelgren L-E, Carlstrom K, et al. The distribution and metabolism of ¹⁴C labelled tamoxifen in spayed female mice. *Acta Pharmacol Toxicol* 1982; 50: 161-8.
 - 232 Theve NO, Pousette A, Carlstrom K. Adenocarcinoma of the pancreas - a hormone sensitive tumour. A preliminary report on nolvadex treatment. *Clin Oncol* 1983; 9: 193-7.
 - 233 Gonzalez-Barcena D, Rangel-Garcia NE, Perez-Sanchez PL, et al. Response to D-TRP-6-LH-RH in advanced adenocarcinoma of the pancreas. *Lancet* 1986; ii: 154.
 - 234 Gonzalez-Barcena D, Ibarra-Olmas MA, Garcia-Carrasco F, et al. Influence of d-Trp-6-LH-RH on the survival time in patients with advanced pancreatic cancer. *Biomed Pharmacother* 1989; 43: 313-7.
 - 235 Lhoste EF, Roebuck BD, Longnecker DS. Effect of steroids on the early stage of azaserine-induced pancreatic carcinogenesis in the rat [Abstract]. *Dig Dis Sci* 1986; 31: 1139.
 - 236 Sumi C, Longnecker D, Roebuck BD, Brinck-Johnson T. Inhibitory effect of estrogen and castration on the early stages of pancreatic carcinogenesis in Fischer rats treated with azaserine. *Cancer Res* 1989; 49: 2332-6.
 - 237 Lointier P, Raboudi SH, Jones LA, et al. Antiestrogen binding sites (AEBS) may mediate 4-OH-tamoxifen (4-OH-TAM) growth inhibitory effects in human pancreatic adenocarcinoma [Abstract]. *Dig Dis Sci* 1986; 31: 1140.
 - 238 Benz C, Hollander C, Miller B. Endocrine-responsive pancreatic carcinoma: steroid binding and cytotoxicity studies in human tumor cell lines. *Cancer Res* 1986; 46: 2276-81.
 - 239 Greenway B, Iqbal MJ, Johnson PJ, Williams R. Low serum testosterone concentration in patients with carcinoma of the pancreas. *BMJ* 1983; 286: 93-5.
 - 240 Militello C, Sperti C, Foresta C, et al. Plasma levels of testosterone and hypophyseal gonadotrophins in men affected by pancreatic cancer [Abstract]. *Digestion* 1984; 30: 122.
 - 241 Shearer MG, Taggart D, Gray C, Imrie CW. Useful differentiation between pancreatic carcinoma and chronic pancreatitis by testosterone assay [Abstract]. *Digestion* 1984; 30: 106.
 - 242 Corbishley TP, Iqbal MJ, Wilkinson ML, Williams R. Correlating sex steroids and sex hormone binding globulin in pancreatic adenocarcinoma. *Anticancer Res* 1986; 5: 219-22.
 - 243 Tulassay Z, Sandoz Z, Bodrogi L, et al. An endocrine model for pancreatic cancer. *BMJ* 1988; 297: 1447-8.
 - 244 Hayashi Y, Katayama H. Promoting effect of testosterone propionate on experimental exocrine pancreatic tumours by 4-hydroxy amino quinoline 1-oxide in rats. *Toxicol Lett* 1981; 9: 349-54.
 - 245 Greenway B, Duke D, Pym B, et al. The control of human pancreatic adenocarcinoma xenografts in nude mice by hormone therapy. *Br J Surg* 1982; 69: 595-7.
 - 246 Kloppel G, Lohr M, Moesta V, Bulow M. The effects of sex steroid hormones on pancreatic carcinoma grown in nude mice and tissue culture [Abstract]. *Dig Dis Sci* 1986; 31: 1137.
 - 247 Lawrence JP, Poston GJ, Townsend CM, et al. Stimulatory effect of castration on the growth of hamster pancreatic

- cancer in vivo: mediating role of gastrin. *Gastroenterology* 1988; **94**: A253.
- 248 Iqbal MJ, Greenway B, Wilkinson ML, *et al.* Sex steroid enzyme, aromatase and 5-alpha-reductase in the pancreas: a comparison of normal adult, foetal and malignant tissue. *Clin Sci* 1983; **65**: 71-5.
 - 249 Logsdon CD, Moessner J, Williams JA, Goldfine ID. Glucocorticoids increase amylase mRNA levels, secreting organelles, and secretion in pancreatic acinar AR42J cells. *J Cell Biol* 1985; **100**: 1200-8.
 - 250 Cooper CS, Blair DG, Oskarsson MK, *et al.* Characterization of human transforming genes for chemically transformed, teratocarcinoma, and pancreatic carcinoma cell lines. *Cancer Res* 1984; **44**: 1-10.
 - 251 O'Hara BM, Oskarsson M, Tainsky MA, Blair DG. Mechanism of activation of human ras gene cloned from a gastric adenocarcinoma and a pancreatic carcinoma cell line. *Cancer Res* 1986; **46**: 4695-700.
 - 252 Ornitz DM, Hammer RE, Messing A, *et al.* Pancreatic regulation induced by SV40 T-antigen expression in acinar cells of transgenic mice. *Science* 1987; **238**: 188-93.
 - 253 Black O Jr, Chang BK. Ornithine decarboxylase enzyme activity in human and hamster pancreatic tumour cell lines. *Cancer Lett* 1982; **17**: 87-93.
 - 254 Chang BKR, Black O Jr, Gutman R. Inhibition of growth of human or hamster pancreatic cancer cell lines by alpha-difluoromethylornithine alone and combined with cis-diaminedichloroplatinum (II). *Cancer Res* 1984; **44**: 5100-4.
 - 255 Marx M, Townsend CM Jr, Barranco SC, *et al.* Treatment of pancreatic cancer by alpha-difluoromethylornithine (DFMO), an inhibitor of polyamine synthesis [Abstract]. *Dig Dis Sci* 1983; **28**: 941.
 - 256 Saydjari R, Townsend CM Jr, Barranco SC, *et al.* Effects of cyclosporine A and alpha-difluoromethylornithine on the growth of pancreatic cancer in vitro. *J Natl Cancer Inst* 1986; **77**: 1087-92.
 - 257 Saydjari R, Townsend CM Jr, Barranco SC, Thompson JC. Cyclosporine and alpha-difluoromethylornithine exhibit differential effects on colon and pancreatic cancer in vitro. *Invest New Drugs* 1987; **5**: 251-8.
 - 258 Saydjari R, Townsend CM Jr, Barranco SC, Thompson JC. Differential sensitivity of pancreatic and colon cancer to cyclosporine and alpha-difluoromethylornithine in vivo. *Life Sci* (in press).
 - 259 Warburg O. On the origin of cancer cells. *Science* 1956; **123**: 309-14.
 - 260 Schek N, Hall BL, Finn OJ. Increased glyceraldehyde-3-phosphate dehydrogenase gene expression in human pancreatic adenocarcinoma. *Cancer Res* 1988; **48**: 6354-9.
 - 261 Yamaguchi N, Kawai K. Acid protease secreted from human pancreatic carcinoma cell line HPC-YT into serum free, chemically defined, medium. *Cancer Res* 1986; **46**: 5353-9.
 - 262 Kaprowski H, Steplewski Z, Mitchell K, *et al.* Colorectal carcinoma antigen detected by hybridoma antibodies. *Somatic Cell Mol Genet* 1979; **5**: 957-72.
 - 263 Malesci A, Tommasini M, Bocchia P, *et al.* Differential diagnosis of pancreatic cancer and chronic pancreatitis by a monoclonal antibody detecting a new cancer associated antigen (CA19-9). *La Ricerca Clin Lab* 1984; **14**: 303-6.
 - 264 Malesci A, Tommasini MA, Bonato C, *et al.* Determination of CA19-9 antigen in serum and pancreatic juice for differential diagnosis of pancreatic adenocarcinoma from chronic pancreatitis. *Gastroenterology* 1987; **92**: 60-7.
 - 265 Kyriazis AA, Kyriazis AP, Sternberg CN, *et al.* Morphological, biological, biochemical and karyotypic characterization of human pancreatic ductal adenocarcinoma CAPAN-2 in tissue culture and nude mouse. *Cancer Res* 1986; **46**: 5810-5.
 - 266 Takasaki H, Tempero MA, Uchida E, *et al.* Comparative studies on the expression of tumour-associated glycoprotein (TAG-72) CA19-9 and DU-Pan-2 in normal, benign and malignant pancreatic tissue. *Int J Cancer* 1988; **42**: 681-6.
 - 267 Klapdor U, Klapdor R, Bahld M, *et al.* Effects of cytotoxic drugs on tumor marker secretion of gastrointestinal/pancreatic carcinomas - comparison of studies in nude mice and in patients. *Strahlenther Oncol* 1989; **165**: 550-2.
 - 268 Dietel M, Arps H, Bals U, *et al.* Individualization of chemotherapy through in vitro predictive determination of the cytostatic sensitivity of malignant tumors. *Dtsch Med Wochenschr* 1989; **114**: 1645-52.