Polymyositis associated with ulcerative colitis

S Chugh, J B Dilawari, I M S Sawhney, N Dang, B D Radotra, Y K Chawla

Abstract

An elderly woman with chronic ulcerative colitis who developed proximal muscle weakness, increased serum creatine phosphokinase activity, and histological and electromyographic abnormalities characteristic of polymyositis is described. Treatment with corticosteroids and 5-acetylsalicylic acid was followed by a remission in bowel symptoms, improvement in muscle power, and reversal of electromyographic changes. An autoimmune link between the two disorders seems likely. (Gut 1993; 34: 567-569)

Extraintestinal manifestations of inflammatory bowel disease are known to affect several organ systems.¹ Although 'granulomatous' myositis is mentioned occasionally as a rare complication of inflammatory bowel disease,² most reports on ulcerative colitis do not mention any type of myositis as a complication or association of the disorder.^{1,3} We have been able to find only five cases of myositis⁴⁻⁸ of different types complicating ulcerative colitis, of which only two are published in English.^{4,5} Because of its rarity, we document the case of an elderly woman who developed polymyositis as a complication of long standing ulcerative colitis.

Case report

A 78 year old woman was admitted to hospital for investigation and management of diarrhoea and muscle weakness. She had been having recurrent episodes of diarrhoea associated with the passage of blood and mucus, low grade fever, and a feeling of exhaustion for the past 15 years. Each episode lasted for 7 to 10 days, followed by a symptom free period of 2-3 months. She had never visited hospital and was treated symptomatically by general practitioners with anti-diarrhoeal drugs. During the past 15 days, she had started having arthralgias that affected multiple joints and appreciable proximal muscle weakness of both lower and upper limbs. The patient was prescribed indomethacin (25 mg thrice daily) for arthralgia by her treating physician. She had lost about 4 kg of weight over the previous 2 weeks.

Physical examination at the time of hospital admission showed moderate pallor and mild dehydration. The patient's pulse rate was 96 bpm and the volume was slightly decreased. Her blood pressure was 100/60 mm Hg. The small joints of the hands were tender to pressure and movement. Abdominal examination showed mild tenderness in the hypogastrium, and the patient's spleen was palpable 2 cm below the left costal margin and was firm in consistency. Neurological examination showed weakness of the proximal muscles of the lower and upper

limbs and muscles of the trunk and flexors of the neck (grade 3/5). The superficial and deep reflexes were normal and her plantars were flexor. Ocular fundi were normal.

Laboratory investigations showed a haemoglobin concentration of 78 g/l and a haematocrit of 23%. The patient's total leukocyte count was 8600/mm³ ($8.6 \times 10^9/l$), with neutrophils 85%, lymphocytes 13%, and eosinophils 2%. The erythrocyte sedimentation rate was 67 mm in the first hour (Westergren) and reticulocyte count 5%. A coagulogram was normal. Platelets were 110 000/mm³ ($110 \times 10^9/l$). Serum Na was 135 mmol/l and K 3.3 mmol/l. Blood urea was 28 mg/dl (9.9 mmol/l) and serum creatinine 1.0 mg/dl (88.4 mmol/l). Serum bilirubin was 0.8 mg/dl (13.6 mmol/l); SGOT was 13 U/l, SGPT was 7 U/l and alkaline phosphatase was 10 KAU. The total serum protein was 74 g/l with an albumin concentration of 36 g/l, globulin 38 g/l, and an albumin/globulin ratio of 0.95. The patient's urine was negative for protein and sugar, and microscopy showed 10-12 pus cells/high power field. Urine culture showed growth of *Pseudomonas aeruginosa*. Stools showed the presence of red cells (+++++) but no ova or cysts, and culture grew *Candida albicans*. Upper gastrointestinal endoscopy showed features of oesophagitis and erosive gastritis. A barium enema showed extensive ulceration in the entire large bowel (Figs 1 and 2) and pseudopolyp formation (Fig 3). Colonoscopy confirmed the presence of multiple ulcerations and pseudopolyps. Rectal biopsy specimen showed infiltration of the lamina propria by inflammatory cells severe mucodepletion, and cryptitis (Fig 4). No granuloma formation or vasculitis was noted. A biopsy specimen from the left quadriceps muscle showed focal intermyseal infiltration by lymphocytes and occasional phagocytosis of degenera-

Figure 1: Barium enema showing extensive mucosal ulceration (arrows) in the ascending, transverse, and descending colon.

Departments of
Medicine, Hepatology,
Neurology and
Pathology, Postgraduate
Institute of Medical
Education and Research,
Chandigarh, India
S Chugh
J B Dilawari
I M S Sawhney
N Dang
B D Radotra
Y K Chawla

Correspondence to:
Professor J B Dilawari,
Department of Hepatology,
Postgraduate Institute of
Medical Education and
Research, Chandigarh,
160012, India

Accepted for publication
31 August 1992

Figure 2: Barium enema showing ulceration (arrows) in the descending and sigmoid colon.

tive muscle fibres (Fig 5). No evidence of regenerative activity, vasculitis, or granuloma was seen. Electromyography of the right deltoid and right quadriceps muscles showed normal insertional activity. No spontaneous activity was observed. Motor unit potentials were small in amplitude and duration. The interference pattern was full, and 30–40% polyphasia was noted (Fig 6). Myopathic motor unit potentials and significant polyphasia were consistent with the diagnosis of polymyositis.

Nerve conduction studies were normal. The patient's creatine phosphokinase activity was increased five fold (1010 U/l). Rheumatoid factor was positive, and antinuclear antibodies were negative.

COURSE AND MANAGEMENT

The patient was put on parenteral nutrition, oral prednisolone (40 mg once daily), and 5-acetylsalicylic acid (400 mg thrice daily). She also received 20 mg of prednisolone daily by enema and 3 units of blood. In addition, she was given ciprofloxacin (500 mg twice daily) for 7 days for urinary tract infection and ketoconazole (200 mg twice daily) for candidiasis. Her ulcerative colitis

Figure 3: Barium enema showing pseudopolyps in the transverse colon (arrows) and ulceration in the ascending colon.

Figure 4: Photomicrograph of the rectum showing intense inflammatory cell infiltration of the lamina propria, mucodepletion and cryptitis (arrow). (Original magnification $\times 240$)

went into remission after 4 weeks and a significant improvement in her muscle power (grade 4/5) became noticeable after 12 weeks of therapy. Repeat study of the electromyographic pattern 3 months later showed a distinct improvement and this was accompanied by a significant decrease in the serum creatinine phosphokinase activity. The enlarged spleen regressed in size and became impalpable.

Discussion

Skeletal muscle involvement associated with inflammatory bowel disease was first reported in 1970 by Spiro,⁹ who illustrated granuloma formation in striated muscle in a patient with Crohn's disease without giving any further details. In a subsequent study, Tydd and Dyer¹⁰ performed muscle biopsies in 15 patients with Crohn's disease in a search for granulomata. None of these patients, however, had any clinical evidence of myopathy, and the muscle histology was normal in all of them. In 1976, Menard *et al*¹¹ reported granulomatous myositis and myopathy in a middle aged man with Crohn's disease. Two more cases were reported subsequently, one by Gilliam *et al*,¹² who described an unusual vasculitic myositis in a young man with Crohn's disease, and another by Hall *et al*,¹³ who reported a non-specific localised myositis without histological evidence of muscle granulomata or vasculitis in a 32 year old woman with long standing Crohn's disease.

The earliest documentation of myositis com-

Figure 5: Photomicrograph of the left quadriceps muscle showing focal intermyseal lymphocytic infiltration and phagocytosis of degenerative muscle fibres. (Original magnification $\times 550$)

Figure 6: Electromyograph of the right deltoid muscle showing myopathic motor unit potentials, polyphasia (above) and normal interference pattern (below).

Comparison of three reported cases of myositis in ulcerative colitis

Clinical and laboratory parameters	Kaneoka et al (1990)	Bhigjee et al (1987)	Present case
Age at onset of myositis (y)	57	42	78
Sex (M/F)	F	M	F
Duration of colitis prior to onset of muscle weakness (y)	11	12	15
Drugs before onset of myositis	Steroids	Steroids and sulphasalazine	No steroids or sulphasalazine
Splenomegaly	Massive, regressed after treatment of myositis	Absent	Mild, regressed after treatment of myositis
Creatine phosphokinase activity	301 U/l	262 U/l	1010 U/l
Antinuclear factor	+	—	—
Rheumatoid factor	+	Not mentioned	+

plicating ulcerative colitis came from Oshitani *et al*,⁸ who described dermatomyositis and mononeuritis multiplex in ulcerative colitis and Basedow's disease. Four more cases, excluding ours, have subsequently been reported to date,^{4,7} of which only two are reported in English.^{4,5}

Bhigjee *et al*⁴ recorded the case of a 44 year old man who presented with ulcerative colitis of 12 years' duration and muscle weakness of 3 years. He underwent colectomy for his bowel disease but despite surgery, the leg weakness progressed resulting in wasting of the quadriceps muscles.

Kaneoka *et al*⁵ described polymyositis in a 63 year old woman who suffered from ulcerative colitis and developed liver cirrhosis secondary to post-transfusional hepatitis. After a relatively symptom free course of colitis for several years, she developed severe proximal muscle weakness and generalised atrophy. She was observed to have an enlarged spleen, which regressed after administration of steroids for polymyositis.

In both cases mentioned above and the one observed by us, myositis occurred after

long standing ulcerative colitis (Table). Serum creatinine phosphokinase activity was raised and there was no evidence of vasculitis or granulomata on histology. In the absence of a history of sulphasalazine ingestion before the onset of myositis in two cases, one reported by Kaneoka *et al*⁵ and the second observed by us, the role of this drug in the pathogenesis of muscle disease is doubtful. In addition, it is evident from the case described by Bhigjee *et al*⁴ that presence of active colitis is not essential for the progression of the muscle disease. The disease entity seems to affect both sexes and all ages since the condition has also been observed in a child.⁷ Regression of splenomegaly after steroid therapy in two patients (Table) and the presence of autoantibodies in the present patient as well as that described by Kaneoka *et al*⁵ suggest a possible autoimmune link between ulcerative colitis and polymyositis.

In the somewhat unrelated case of a 19 year old man with ulcerative colitis and necrotising cerebral vasculitis observed by Nelson *et al*,¹⁴ a biceps muscle biopsy specimen showed 'a few necrotic myofibres and mild perivascular inflammation'. Though not commented on by the authors, the presence of myositis with or without vasculitis remains a distinct possibility in their patient. Recognition of myositis as an extraintestinal manifestation of ulcerative colitis seems to be important for future documentation of this association.

1 Greenstein AJ, Janowitz HD, Sachar D. The extra intestinal complications of Crohn's disease and ulcerative colitis. A study of 700 patients. *Medicine (Baltimore)* 1976; 55: 401-12.

2 Glickman RM. Inflammatory bowel disease. In: Wilson JD, Braunwald E, Isselbacher KJ, Petersdorf RG, Martin JB, Fauci AS, Root RK, eds. *Harrison's principles of internal medicine*. Vol 2, 12th ed. New York: McGraw Hill, 1991: 1268-81.

3 Cello JP, Schneiderman DJ. Ulcerative colitis. In: Sleisenger MH, Fordtran JS, eds. *Gastrointestinal disease. Pathophysiology, diagnosis and management*. 4th ed. Philadelphia: W B Saunders, 1989: 1435-76.

4 Bhigjee AI, Bill PLA, Cosnett JE. Ulcerative colitis and interstitial myositis. *Clin Neurol Neurosurg* 1987; 89: 261-3.

5 Kaneoka H, Iyadomi I, Hiida M, Yamamoto K, Kisu T, Tokunaga O, Yamaguchi M. An overlapping case of ulcerative colitis and polymyositis. *J Rheum* 1990; 17: 274-6.

6 Evrard P, Lefebvre C, Brucher JM, Coche E. Myositis associated with ulcero-hemorrhagic rectocolitis. Apropos of a case. *Acta Gastroenterol Belg* 1987; 50: 675-9.

7 Hernandez MA, Diez-Tejedor E, Barbado J, Morales C. Polymyositis and ulcerative colitis. A new form of association? *Neurologia* 1986; 1: 134-5.

8 Oshitani H, Wakabayashi Y, Sawaguchi Y, Koike H, Yoshino Y. A case of dermatomyositis followed by multiple mononeuritis, with the history of ulcerative colitis and Basedow's disease. *J Kyojin Med Soc* 1981; 12: 47-53.

9 Spiro HM. *Clinical Gastroenterology*. Toronto: MacMillan, 1970: 631.

10 Tydd TF, Dyer NH. Muscle biopsy in Crohn's disease. *Lancet* 1974; ii: 1574-5.

11 Menard DB, Haddad H, Blain JG, Beaudry R, Devroede G, Masse S. Granulomatous myositis and myopathy associated with Crohn's colitis. *New Engl J Med* 1976; 295: 818-9.

12 Gilliam JH, Challa VR, Agudelo CA, Albertson DA, Huntley CC. Vasculitis involving muscle associated with Crohn's colitis. *Gastroenterology* 1981; 81: 787-90.

13 Hall MJ, Thomas WEG, Cooper BT. Gastrocnemius myositis in a patient with inflammatory bowel disease. *Digestion* 1985; 32: 296-300.

14 Nelson J, Barron MM, Riggs JE, Gutmann L, Schochet SS Jr. Cerebral vasculitis and ulcerative colitis. *Neurology* 1986; 36: 719-21.