One week's anti-*Helicobacter pylori* treatment for duodenal ulcer

R P H Logan, P A Gummett, J J Misiewicz, Q N Karim, M M Walker, J H Baron

Abstract

This open study tested whether eradication of *Helicobacter pylori* (*H. pylori*) heals duodenal ulcers as well as decreasing recurrence. *H. pylori* was detected in patients with endoscopic duodenal ulcers by histology, CLO-test, culture, and ^{13}C -urea breath test (^{13}C -UBT). Tripotassium dicitrate bismuthate (120 mg) and amoxycillin (500 mg) each four times daily, were given for seven days, with 400 mg metronidazole five times a day on days 5-7. The ^{13}C -UBT was repeated immediately after treatment and endoscopy repeated within 21 days. After treatment unhealed ulcers were reinspected one month later and healed ulcers followed up by ^{13}C -UBT alone for 12 months. Of 45 patients, 44 were available for follow up. Mean pretreatment excess $\delta^{13}\text{CO}_2$ excretion was 25.6 per mil, which fell to 2.4 per mil immediately after finishing treatment, indicating clearance of *H. pylori* in every patient. At the second endoscopy (median interval 20 days from start of treatment) 33 of 44 (75%) duodenal ulcers had healed. Ten of the remaining 11 duodenal ulcers were smaller and those 10 healed in the next two weeks with no further treatment. Two patients' ulcers that initially healed with clearance of *H. pylori* recurred three weeks later (both had metronidazole resistant *H. pylori*). *H. pylori* was eradicated in 28 of 44 (64%) patients (^{13}C -UBT negative for median follow up 10.2 months). Overall 41 of 43 (93%, 95% confidence intervals 81%-99%) duodenal ulcers were healed at one month. This study suggests that one week of anti-*H. pylori* triple treatment is effective in healing duodenal ulcers.

(Gut 1994; 35: 15-18)

Helicobacter pylori (*H. pylori*) causes non-autoimmune gastritis¹ and is one important factor in the aetiology of recurrent duodenal ulcer.² Thus recurrence of *H. pylori* precedes ulcer recurrence,³ the density of antral *H. pylori* increases the rate of ulcer recurrence⁴ and successful eradication of *H. pylori* after duodenal ulcer healing substantially prevents recurrence.⁵⁻⁷ All of these studies, however, focused on the role of *H. pylori* in recurrence of duodenal ulcers rather than on primary ulcer healing. We have previously developed a novel one week triple treatment that eradicated 93% of metronidazole sensitive *H. pylori*.⁸ We have now for the first time investigated healing of duodenal ulcers with this regimen as the sole primary treatment.

Patients and methods

After routine diagnostic endoscopy of the upper gastrointestinal tract patients with a duodenal

ulcer greater than 5 mm in diameter were invited to enter and give written consent to the study, which was approved by the Parkside ethics committee. Patients with previous gastric surgery, known bleeding diathesis, taking oral anticoagulants, or who had been treated in the previous two months with bismuth compounds or antibiotics known to be active against *H. pylori* were excluded. Patients taking H_2 antagonists, proton pump inhibitors, or whose ulcers had recently bled were not excluded.

Endoscopes were disinfected with an automatic washing machine (Olympus EW20)⁹ after each examination and the biopsy forceps were sterilised by autoclaving.

ASSESSMENT OF *H. PYLORI* STATE

The presence of *H. pylori* was assessed by the ^{13}C -urea breath test (^{13}C -UBT), histological examination of two antral biopsy specimens, CLO-test (Delta West Ltd, Western Australia), and culture (two antral biopsies, selective and non-selective media, microaerophilic conditions for up to 10 days). Tests for metronidazole sensitivity (5 μg) were completed by an in vitro disc method (mast sensitivity discs, Mast Laboratories, Liverpool). Metronidazole resistant or metronidazole sensitive *H. pylori* are subsequently referred to as MR *H. pylori* and MS *H. pylori*, respectively.

Patients were classified as *H. pylori* positive by a positive ^{13}C -UBT and CLO-test together with positive histology or culture. Clearance of *H. pylori* was defined as a negative ^{13}C -UBT at the end of treatment. Eradication was defined as a negative ^{13}C -UBT one month after the end of treatment (negative ^{13}C -UBT = excess $\delta^{13}\text{CO}_2$ excretion <5 per mil).¹⁰

^{13}C -UREA BREATH TEST

The ^{13}C -UBT (European standard protocol)¹⁰ was completed within 24 hours of the initial endoscopy in all patients. Briefly, a baseline sample of expired breath was obtained before ingesting a fatty liquid test meal to delay gastric emptying. After 10 minutes, 100 mg ^{13}C -urea (99% pure, Cambridge Isotopes, Boston, USA) in 50 ml of tap water was swallowed and distributed within the stomach by turning the patient to the left and right decubitus position. Two litre serial breath samples were collected every five minutes into a large reservoir collecting bag, from which a single 20 ml sample was taken at the end of the test and analysed by mass spectrometry (BSIA, Brentford, London). A positive result was defined as excess $\delta^{13}\text{CO}_2$ excretion >5 per mil, as determined from previous studies.¹⁰

Patients with a positive ^{13}C -UBT and positive

Parkside *Helicobacter*
Study Group, Central
Middlesex and St Mary's
Hospitals, London
R P H Logan
P A Gummett
J J Misiewicz
Q N Karim
M M Walker
J H Baron

Correspondence to:
Dr R P H Logan, Department
of Gastroenterology and
Nutrition, Central Middlesex
Hospital, Acton Lane, Park
Royal, London NW10 7NS.

Accepted for publication
19 May 1993

Outline of study design with the number of patients with healed ulcers at first or second follow up. OGD=endoscopy.

CLO-test were given 120 mg tripotassium dicitrato bismuthate and 500 mg amoxycillin, each four times a day, for seven days (days 1–7). Metronidazole (400 mg) was given five times a day for the last three days (days 5–7). This high dose was used to decrease the risk of promoting the development of MR *H pylori*.

FOLLOW UP
Immediately after the end of eradication treatment, a second ¹³C-UBT was performed to assess clearance (to determine the effectiveness of treatment), followed as soon as possible by a second endoscopy to assess ulcer healing. In those patients with healed ulcers further ¹³C-UBTs were done at one, three, six, and 12 months: they were re-endoscoped only if ulcer symptoms recurred. Patients whose ulcer had not healed were again endoscoped and had a ¹³C-UBT one month after the end of eradication treatment. If the ulcer was healed, they had further ¹³C-UBTs at three, six, and 12 months (Figure).

Ulcer healing and relapse, metronidazole sensitivity, and eradication

No of patients	H pylori cleared	Ulcer healed at first follow up	H pylori eradicated	Ulcer healed only at one month	Ulcer recurred or unhealed
MS <i>H pylori</i> (23)	23	15	20	8	0
MR <i>H pylori</i> (12)	12	12	2	0	2
M sensitivity not known (9)	9	6	6	2	2
Total (44)	44	33	28	10	4

Patients who failed to clear *H pylori* after treatment, or whose ulcers failed to heal, or who had recurrent *H pylori* at one month were not studied further.
Confidence intervals on proportions are based on the binomial distribution.

Results
The Figure shows the study design. The Table summarises ulcer healing and relapse, metronidazole sensitivity, and eradication data.

PATIENTS
Forty five patients (27 men, median age 37, range 17–76 years) with duodenal ulcers (>5 mm) and *H pylori* were entered into the study; 44 are available for follow up. The remaining patient declined further endoscopy and follow up because he had no symptoms. Twenty two patients were smokers, five presented with either haematemesis or melaena, of whom three had been started on H₂ antagonists within 24 hours of the initial endoscopy, and a further three patients had duodenal ulcers resistant to standard treatment: 300 mg ranitidine at night (n=1), 400 mg cimetidine twice a day (n=1), or 40 mg omeprazole in the morning (n=1), each for one month. Any treatment with H₂ antagonists or proton pump inhibitor was stopped within 24 hours of the initial endoscopy in all patients.

ENDOSCOPY
At the second endoscopy 33 of 44 (75%, 95% confidence intervals (95% CI) 60%–87%) ulcers were completely healed, 10 of the remaining 11 were smaller (median diameter 3 mm), and one was larger. The median interval between the end of treatment and the second endoscopy was 13 days (range 1–31 days, variation due to missed appointments and holidays).
One month after the end of treatment 10 of the initially unhealed 11 ulcers had healed completely without any further treatment in the intervening three weeks. The remaining ulcer had increased in size (MR *H pylori*). Two patients whose ulcers initially healed (both infected with MR *H pylori*) had an early symptomatic ulcer recurrence, which was confirmed at endoscopy three weeks after the end of treatment. Thus overall 43 of 44 (98%) duodenal ulcers healed after the one week *H pylori* eradication regimen, with 41 of 44 (93%, 95% CI 81%–99%) still healed at one month.

H PYLORI AND METRONIDAZOLE SENSITIVITY
All 44 patients were colonised by *H pylori* as detected by a positive ¹³C-UBT and positive antral histology. The ¹³C-UBT became negative in all 44 patients immediately after finishing treatment indicating that the treatment regimen had cleared *H pylori* in every patient. *H pylori* was grown from 38 of 44 patients and metronidazole sensitivity established in 35: 12 (34%) isolates were resistant and 23 (66%) were sensitive to metronidazole. In these 23 patients the

ulcers were healed at one month or earlier in 20 of 23 (87%, 95% CI 60%–95%) (Table), with neither recurrent symptoms nor recurrence of *H pylori* during subsequent follow up (median = 10, range 6–14 months). In the 12 patients with pretreatment MR *H pylori* duodenal ulcer healing was achieved in all 12, but *H pylori* was eradicated in only two of 12 (17%).

FOLLOW UP

Overall *H pylori* was eradicated in 28 of 44 patients (64%, 95% CI 48%–76%). In these patients there was no evidence by ¹³C-UBT of recurrent *H pylori* at three and six months and no recurrence of ulcer symptoms during this period of follow up. In the other 16 patients the ¹³C-UBT became positive indicating failed eradication: in these 16 with persistent *H pylori*, one duodenal ulcer had not healed at one month follow up, two had recurrent ulcers within three weeks of finishing treatment, and five had recurrent dyspepsia but no recurrent ulcers on repeat endoscopy.

Discussion

For many years the treatment of duodenal ulcer has been based on Schwartz's dictum 'no acid – no ulcer', and whereas this is remarkably successful with 90%–100% duodenal ulcer healing within two months of antisecretory treatment, there is a disappointingly high rate of ulcer recurrence on stopping treatment – 50% can recur within six months.¹¹ It is now clear that acid inhibitors do not change the natural history of the underlying ulcer diathesis. The natural history can be changed, however, by eradication of *H pylori*, which decreases the incidence of ulcer recurrence.^{5–7}

Previous studies of duodenal ulcer and *H pylori* (with comparatively lengthy treatment regimens of antibiotics in conjunction with either bismuth salts, or antisecretory agents) have focused on recurrence of *H pylori* and relapse to duodenal ulcers,^{5–7} whereas all the studies of duodenal ulcer healing with bismuth monotherapy were done before the discovery of *H pylori*.

ULCER HEALING

This study is the first to examine the alternative strategy of using an anti-*H pylori* regimen as the primary (and only) treatment for healing duodenal ulcers. On an intention to treat basis 75% of duodenal ulcers healed on the seven day *H pylori* eradication regimen. Ulcer healing was not complete in 11 patients at the first follow up endoscopy (median interval from the end of treatment nine days). Further treatment was not needed and 10 of 11 duodenal ulcers healed completely within 28 days, or earlier, after finishing eradication treatment, achieving a 93% healing rate at one month after the end of the regimen.

This open pilot study was uncontrolled, but the results are comparable with a recent two week trial of 20 mg omeprazole in the morning *v* 150 mg ranitidine twice daily with two

week healing rates of 86% and 63% respectively.¹²

BISMUTH

There are no studies of the rate of duodenal ulcer healing with bismuth salts after less than four weeks of treatment^{13–15} and it is possible that the 75% healing rate could have been due to the bismuth in our triple treatment regimen. In studies with bismuth salts as the sole agent, healing and prevention of recurrence may have been promoted by the anti-*H pylori*, mucosal protective, or other properties of accumulated bismuth. Bismuth monotherapy can heal duodenal ulcers without inhibition of acid secretion,^{13,15} but it is not clear whether the benefit is due to suppression of *H pylori* or other mechanisms – for example, decreasing pepsin.¹⁶ Eradication rates with bismuth salts alone, however, are very low.¹⁷

H PYLORI

Although many workers are at present attempting to identify pathogenic factors characterising strains of *H pylori* associated with duodenal ulcers,^{18–21} our study suggests that regardless of pathogenicity, the overall load of the organism may be important in determining the clinical outcome. Thus in two patients duodenal ulcers healed after suppressing *H pylori* to a low level of infection (excess $\delta^{13}\text{CO}_2$ excretion 1.3 and 2.1 per mil for each patient at the end of treatment). Within days of finishing treatment the level of infection, as measured by the ¹³C-UBT, had returned to pretreatment levels: this was followed by ulcer recurrence 10 days later.

METRONIDAZOLE SENSITIVITY

Duodenal ulcers healed in all the 23 patients infected with MS *H pylori*. In 20 of these 23 patients *H pylori* was eradicated and neither ulcer nor *H pylori* had recurred within the nine months of follow up. The ulcers also healed in the 12 patients with MR *H pylori*, but *H pylori* was eradicated in only two of 12. Although those patients with persistent *H pylori* are at risk of duodenal ulcers, symptomatic ulcer recurrence has so far not occurred in any patient. Recurrence without symptoms is possible but could be delayed, as seen after bismuth monotherapy,^{14,15} which generally only suppresses *H pylori*.¹⁷

We and others have shown that the most important factor in preventing recurrent duodenal ulcer is the success of the treatment regimen in eradicating *H pylori*. Previous studies suggest that any regimen that includes metronidazole will not work if the stomach is infected with MR *H pylori*.^{8,21,22} The design of this study prevented the exclusion of patients with MR *H pylori* and this is the most likely reason for the low overall eradication rate. For optimal management strategy, routine *in vitro* antibiotic sensitivity tests are desirable before attempting to eradicate *H pylori*, in order to treat patients with drugs that are likely to be successful against their bacteria and to prevent the emergence of resistant strains.

Although no formal symptom assessments were made in this study, no patient needed additional treatment after their initial one week eradication regimen, including the patients with unhealed ulcers at the first follow up endoscopy.

The main conclusion of this study is that one week's triple treatment (which had been shown to be highly effective in eradicating *H pylori*) is highly effective in healing duodenal ulcers. Ulcer healing is probably, but not certainly, due to clearance of *H pylori* rather than to other effects of the bismuth, amoxycillin, or metronidazole. The ulcers that healed after clearance of *H pylori* all remained healed during the period of follow up if the *H pylori* had been eradicated, but two of those whose *H pylori* had been cleared but not eradicated recurred during follow up.

This approach to duodenal ulcer healing poses new problems, particularly that of MR *H pylori*. Anti-*H pylori* regimens containing metronidazole will not be successful in areas where the incidence of MR *H pylori* is high.²² Further studies, including symptom assessments and follow up endoscopy in all patients are needed, before the proper place of this strategy in the management of duodenal ulcer can be established.

We thank the endoscopy staff at the Central Middlesex and St Mary's Hospitals and Luke Whitaker for their help with this study. Dr R Logan was supported by a grant from Glaxo Group Research.

An abstract of this study was presented to British Society of Gastroenterology in September 1991.

- 1 Rauws EA, Langenberg W, Houthoff HJ, Zanen HC, Tytgat GNJ. Campylobacter pyloridis-associated chronic active antral gastritis. A prospective study of its prevalence and the effects of antibacterial and antiulcer treatment. *Gastroenterology* 1988; **94**: 33-40.
- 2 Axon AR. Duodenal ulcer: the villain unmasked? *BMJ* 1991; **302**: 919-21.
- 3 Logan RPH, Gummert PA, Polson RJP, Walker MM, Baron JHB, Misiewicz JJ. The recurrence of Helicobacter pylori in relation to duodenal ulcer. *Rev Esp Enferm Dig* 1990; **78** (suppl 1): 72.
- 4 Hui WM, Ho J, Lam SK. Pathogenic role of Helicobacter pylori in duodenal ulcer disease. *Dig Dis Sci* 1991; **36**: 424-30.
- 5 Marshall BJ, Goodwin CS, Warren JR, et al. A prospective double-blind trial of duodenal ulcer relapse after eradication of Campylobacter pylori. *Lancet* 1988; **2**: 1437-42.
- 6 Coghlan JG, Humphries H, Dooley C, et al. Campylobacter pylori and recurrence of duodenal ulcers - a 12-month follow-up study. *Lancet* 1987; **ii**: 1109-11.
- 7 Rauws EA, Tytgat GNJ. Cure of duodenal ulcer with eradication of Helicobacter pylori. *Lancet* 1990; **335**: 1233-5.
- 8 Logan RPH, Gummert PA, Misiewicz JJ, et al. A one week eradication regime for Helicobacter pylori. *Lancet* 1991; **338**: 1249-52.
- 9 Weller IVD, Williams CB, Jeffries DJ, et al. Cleaning and disinfection of equipment for gastrointestinal flexible endoscopy: interim recommendations of a Working Party of the British Society of Gastroenterology. *Gut* 1988; **29**: 1134-51.
- 10 Logan RPH, Dill S, Bauer FE, et al. The European ¹³C-urea breath test for the detection of Helicobacter pylori. *European Journal of Gastroenterology and Hepatology* 1991; **3**: 905-11.
- 11 Thomas JM, Misiewicz JJ. Histamine H₂ antagonists in the short and longterm treatment of duodenal ulcer. *Clinics in Gastroenterology* 1984; **13**: 501-4.
- 12 Hui WM, Lam SK, Ho J, et al. Effect of omeprazole on duodenal ulcer-associated antral gastritis and Helicobacter pylori. *Dig Dis Sci* 1991; **36**: 577-82.
- 13 Moshal MG. The treatment of duodenal ulcers with TDB: a duodenoscopic double-blind cross-over investigation. *Postgrad Med J* 1975; **51** (suppl 5): 36-40.
- 14 Martin DF, Hollanders D, May SJ, et al. Difference in relapse rates of duodenal ulcer healing with cimetidine or tripotassium dicitrato bismuthate. *Lancet* 1981; **i**: 7-10.
- 15 Lee FI, Samloff IM, Hardman J. Comparison of Tripotassium dicitrato bismuthate tablets with ranitidine in healing and relapse of duodenal ulcers. *Lancet* 1985; **290**: 1299-302.
- 16 Sidebotham RL, Batten JJ, Li K, Spencer J, Baron JH. Some effects of tripotassium dicitrato bismuthate on gastric secretion and the mucus barrier. *Aliment Pharmacol Therap* 1990; **4**: 535-46.
- 17 Logan RPH, Polson RJ, Baron JH, Misiewicz JJ. Follow-up after anti-Helicobacter pylori treatment. *Lancet* 1991; **337**: 562-3.
- 18 Figura N, Guglielmetti A, Rossolini A, et al. Cytotoxin production by Campylobacter pylori strains isolated from patients with peptic ulcers and from patients with chronic gastritis only. *J Clin Microbiol* 1989; **27**: 225-6.
- 19 Cover TL, Dolley CP, Blaser MJ. Characterisation of and human serological response to proteins in Helicobacter pylori broth culture supernatants with vacuolising cytotoxin activity. *Infect Immun* 1990; **58**: 603-10.
- 20 Crabtree JE, Taylor JD, Wyatt JI, et al. Mucosal IgA recognition of Helicobacter pylori 120 kDa protein, peptic ulceration, and gastric pathology. *Lancet* 1991; **338**: 332-5.
- 21 Lee A. H pylori initiated ulcerogenesis: look to the host. *Lancet* 1993; **341**: 280-1.
- 22 Glupczynski Y, Burette A. Drug therapy for Helicobacter pylori infection: Problems and pitfalls. *Am J Gastroenterol* 1990; **85**: 1545-51.
- 23 Glupczynski Y, Burette A, De Koster E, et al. Metronidazole resistance in Helicobacter pylori. *Lancet* 1990; **335**: 976-7.