LETTERS TO THE EDITOR

Strongyloidiasis

EDITOR,—We were delighted to read Dr Grove's leading article on the protean manifestations of strongyloidiasis (*Gut* 1994; 35: 437–40) and the case report by Dr Delarocque Astagneau *et al* of a case presenting with biliary obstruction (*Gut* 1994; 35: 705–6), particularly with the reported increase in the incidence among migrant workers and holiday travellers to endemic areas. We recently submitted a case report of a West Indian man living in the United Kingdom since 1965 with disseminated strongyloidiasis who, similar to the patient in the case report, was also seropositive for human T lymphotropic virus (HTLV)-1.¹ These cases highlight the well recognised association between HTLV-1 infection and disseminated strongyloidiasis, particularly in areas of high prevalence in the West Indies and southwest Japan.

Thirty eight to 47.8% of patients from a Japanese series with disseminated strongyloidiasis were HTLV-1 positive.^{2–4} Infection with HTLV-1, commonly associated with the adult T cell leukaemia/lymphoma syndrome in Japan and HTLV-1 associated myelopathy/tropical spastic paraparesis in the West Indies, causes impaired cell mediated immunity without reducing T cell helper/suppressor ratios, and may cause a predisposition to disseminated helminth infections by reducing serum concentrations of IgE and IgE binding factors.⁵ Relapse rates after treatment for disseminated strongyloidiasis with a single agent are higher in HTLV-1 positive patients. Patients from the West Indies or Japan with disseminated strongyloidiasis should therefore be screened for HTLV-1 infection. Treatment with at least two agents is recommended with closer follow up for symptoms and signs of recrudescence or re-infection.^{6,7}

R G SHIDRAWI
The Rayne Institute,
St Thomas' Hospital,
London

D WESTABY
Chelsea & Westminster Hospital,
London

- Shidrawi RG, Shousha S, Westaby D. Systemic strongyloidiasis in an HTLV-1 positive male presenting as chronic pancreatitis with a pancreatico-biliary stricture. *Eur J Gastroenterol Hepatol* (in press).
- Zaha O. Investigation of clinical symptoms in slight strongyloidiasis. *J Jap Assoc Infect Diseases* 1992; 66: 1378–81.
- Hanada S, Uematsu T. The prevalence of human T-cell leukemia virus type 1 infection in patients with hematologic and nonhematologic diseases in an adult T-cell leukemia-endemic area of Japan. *Cancer* 1989; 64: 1290–5.
- Sato Y, Toma H. Seroprevalence studies on the concomitance of Strongyloidiasis with T-cell leukemia viral infection in Okinawa, Japan. *Japanese Journal of Parasitology* 1990; 39: 376–83.
- Matsumoto T, Miike T. Decreased serum levels of IgE and IgE-binding factors in individuals infected with HTLV-1. *Clin Exp Immunol* 1990; 81: 207–11.
- Testa J, Kizimandji-Coton G. Ivermectin in the treatment of strongyloidiasis, ascariasis and

ancylostomiasis in the Central African Republic of Bangui. *Med Afr Noire* 1990; 37: 283–4.

7 Naquira C, Jimenez G. Ivermectin for human strongyloidiasis and other intestinal helminths. *Am J Trop Med Hyg* 1989; 40: 304–9.

Pancreas in end stage renal disease

EDITOR,—I read with interest the article by Lerch and colleagues (*Gut* 1994; 35: 401–7) describing the development of pancreatic insufficiency in experimental renal failure. I, however, was very surprised to see one of my papers¹ given as a reference for statements in the introduction and discussion of their manuscript to support claims that exocrine pancreatic insufficiency occurs in renal failure. In contrast, we showed increased trypsin and normal lipase secretion in patients with renal failure undergoing dialysis.¹ Confusion may have arisen because we reviewed the data of others published before our article and gave a number of references that reported normal or decreased pancreatic secretion in renal insufficiency.^{2–6}

Of possible interest, we also found that patients with renal insufficiency had raised plasma concentrations of hormones that affect pancreatic secretion, which led us to hypothesise that the raised hormone concentrations may cause morphological changes of the pancreas.^{1,7} One of the important problems of the rat model of renal failure discussed by Lerch and colleagues is that severe renal insufficiency is only transient; serum creatinine decreased to almost normal values by 56 days. It is entirely possible that severe renal insufficiency was not present long enough to cause prolonged physiological changes such as constantly increased hormonal concentrations to cause morphological abnormalities.

Another possible problem with the study of Lerch *et al* was use of questionable controls; comparisons were made to caerulein pancreatitis which, according to the authors, causes acute pancreatitis whereas the aim of their study was to determine if renal insufficiency causes chronic pancreatitis.

These problems raise the question of whether the study of Lerch *et al* provides much useful information to resolve the question of pancreatic functional and morphological abnormalities in human renal failure.

E P DiMAGNO,
Mayo Foundation,
Rochester,
Minnesota 55905,
USA

- Owyang C, Miller LJ, DiMagno EP, Mitchell III JC, Go VLW. Pancreatic exocrine function in severe human chronic renal failure. *Gut* 1982; 23: 357–61.
- Otte M, Stahlheber H, Forell MM, *et al*. Pancreatic secretion of patients with chronic renal insufficiency. *Klin Wochenschr* 1975; 53: 67–72.
- Bartos V, Melichar J, Erben J. The function of the exocrine pancreas in chronic renal disease. *Digestion* 1970; 3: 33–40.
- Poll VM, Werner J, Huber W, Kempmann G, Willig F. The exocrine pancreatic function in chronic renal insufficiency. *Z Gastroenterol* 1979; 17: 177–86.
- Gebhardt VW, Stein G, Brosseckert H, Graupner C, Noske A. Excretory pancreatic function in kidney failure. *Z Gesamte Inn Med* 1974; 29: 895–9.
- Wittich K-A, Schmidt H, Scheler F, Creutzfeldt W. Exocrine pancreatic function in chronic kidney failure. *Vehr Dtsch Ges Inn Med* 1968; 74: 1072–4.
- Owyang C, Miller LJ, Go VLW, DiMagno EP, Brennan Jr LA. The gastrointestinal hormone profile in renal insufficiency. *Mayo Clin Proc* 1979; 54: 769–73.

Reply

EDITOR,—Dr DiMagno raises some important questions regarding our study and the participation of the exocrine pancreas in chronic renal disease. The data in his earlier report¹ indeed show that the exocrine pancreas responds with increased digestive enzyme secretion (trypsin) to cholecystokinin (CCK) or meal stimulation in vivo and this finding is in accordance with the increased secretory response (amylase and lipase), which we found in vitro at the time interval where renal insufficiency was most severe in our animals (*Gut* 1994; 35: 401–7). His work was meant to be referenced at a different place in our paper and we apologise for this error.

Dr DiMagno further suggests that renal failure was only severe during the initial phase of our observation period, while, at the stage of developing glomerulosclerosis² uraemia was much milder. We agree that an animal model entailing total (in contrast with subtotal) nephrectomy combined with longterm haemodialysis would have more adequately reflected the situation in human end stage renal disease. Such an animal model, however, would be extremely challenging to establish in the laboratory and the course of uraemia in our setting was clearly severe enough to significantly affect pancreatic exocrine function, morphology, DNA, protein synthesis, cellular turnover, and regeneration. We therefore maintain that subtotal nephrectomy, although not truly reflecting end stage renal disease in humans, is a valuable tool to study the effects of renal insufficiency on the exocrine pancreas.

Dr DiMagno reminds us that the pathophysiology as well as the sequelae of uraemic pancreopathy are still a matter of controversy and ongoing study.³ As indicated in the abstract of our report it was previously unclear whether uraemic pancreopathy 'should be regarded as a manifestation of chronic pancreatitis, arising from recurrent attacks of acute pancreatitis, or represents a distinct entity'. To consider these questions we have used for comparison a standard rat model of pancreatitis that combines most of the biochemical and cell biological characteristics reported in other existing animal models of pancreatitis. The changes affecting the pancreas in renal failure bear no resemblance with any of the different features of pancreatitis we investigated. They did, however, resemble toxic damage of the pancreas and we therefore concluded that uraemic pancreopathy is distinct from pancreatitis and probably does not result from recurrent attacks of pancreatitis. The initial hypothesis that recurrent attacks of acute pancreatitis might participate in the changes of the uraemic pancreas is not as improbable as Dr DiMagno suggests because the incidence of acute pancreatitis is significantly increased in patients with end stage renal disease.⁴ We agree that it would have been very interesting to include an additional non-invasive control model to represent chronic pancreatitis in our study (compared with the acute changes associated with the caerulein model we used). Unfortunately no such animal model exists.

The hypothesis proposed by Dr DiMagno that a chronic increase of circulating CCK^{1,5} is responsible for the changes associated with uraemic pancreopathy is intriguing, and, although we have not measured CCK in our rat model, is not excluded by our data. The same increase in