Gut ischaemia

U Haglund

Abstract
Intestinal mucosal injury that results from local ischaemia can be detected by early increases in gut permeability, followed by later morphological, histological, and biochemical abnormalities. Local adaptive mechanisms (for example increased oxygen extraction) can cope with reductions in blood flow of up to 50%, as may occur during episodes of septic shock or cardiac tamponade. Why then does hypoxic injury develop? The peculiar vascular anatomy of the villi allows for oxygen short circuiting to occur at their base, when blood flow is low. Although overall oxygen extraction efficiency may be high, regional hypoxia at the villus tip may, paradoxically occur. The severity of reperfusion injury depends on the duration of preceding hypoxia. Free radical generation through the hypoxanthine xanthine oxidase system is important in mediating cellular damage. In addition, luminal aggressive factors (for example, pancreatic proteases) may cause mucosal damage, as suggested by earlier studies. More recent studies in pigs suggests that pancreatic duct ligation merely delays, but does not prevent development of gut reperfusion injury. Enteral nutrition should benefit patients with the ischaemic intestine because in comparison with total parental nutrition, it stimulates regional blood flow, and attenuates mucosal injury. There are no randomised trials to verify this, but use of tonometry to monitor local ischaemia may help resolve the issue.

(Gut 1994; supplement 1: S73–S76)

The intact gut mucosa serves as a barrier between the non-sterile lumen and the sterile interior of the body. Micro-organisms may sometimes penetrate this barrier – a process referred to as translocation.¹ According to present knowledge, translocation in the normal healthy subject causes only a limited bacteraemic/endotoxaemic challenge, which can be cleared by the defence systems (mesenteric lymph nodes and the hepatic reticuloendothelial system). During pathological states, however, leakage of bacteria through the more permeable gut mucosa may be increased, resulting in a significant septic condition. After trauma, in critical illness or septic states, the intestinal mucosal barrier may be impaired or possibly broken down completely. As a consequence, the possibility for micro-organisms to invade the body is greatly enhanced. In addition, other toxic substances such as myocardial depressant substances and

proteolytic enzymes might be released from the intestine.² Mucosal ischaemia is believed to be a common denominator for the processes that cause enhanced mucosal barrier permeability. The participation of mucosal ischaemia in intestinal mucosal injury and the possible influence by enteral nutrition on these conditions are currently being studied.

Ischaemic intestinal mucosal injury (Table)

The first detectable sign of intestinal mucosal injury in ischaemia is increased capillary permeability. With longterm ischaemia, the mucosal permeability will be further increased with subsequent mucosal epithelial cell injury becoming morphologically detectable. It is important to consider the intestinal mucosal injury process as a continuum of injury ranging from increased permeability to tissue destruction (Fig 1). The extent and duration of ischaemia will determine the depth of the tissue injury³ (Fig 2).

Intestinal ischaemia as seen, for example, in the septic, critically ill patient probably does not cause damage to the gut beyond the superficial part of the mucosa. Intestinal strangulation, mesenteric vascular obstruction, and so called non-occlusive intestinal ischaemia might cause additional injury to the deeper layers of the intestinal wall. In the patient suffering with shock, mucosal injury is confined to the villi. The first morphological sign of tissue injury is the lifting of the epithelial cells at the very tip of the villi.⁴ With more severe or more prolonged ischaemia, or both, the epithelial cells along the sides of the villus are lifted until the epithelial

Mechanism of intestinal mucosal injury

Ischaemic injury
Decreased oxygen delivery
Reduced intestinal (mucosal) blood flow
Increased short circuiting of oxygen in the villus counter-current exchanger
Increased needs of oxygen
Reperfusion injury

Figure 1: The continuum of intestinal ischaemic injury from normal mucosa, permeability changes to transmural infarction.

Department of Surgery, University Hospital, Uppsala, Sweden
U Haglund

Correspondence to:
U Haglund, Department of Surgery, University Hospital, S-751 85 Uppsala, Sweden.

Figure 2: Intestinal ischaemia is characterised by its extent and its duration.

cells are finally lost and the villus core disintegrates. Chiu *et al* describe an injury related grading system that has been widely used⁴ because their various grades have shown to be proportional to the extent and duration of the ischaemia. Even in situations where the villi are severely damaged, healing might occur quite rapidly if measures to provide adequate mucosal tissue oxygenation were instituted. Thus, Park *et al*⁵ showed that epithelial lifting returns to normal within six hours. Villi that had lost their epithelial cells were shown to be covered and to maintain a fairly normal appearance within 18 hours.⁵

Ischaemic mucosal injury

During ischaemia, the intestinal mucosa may be injured as a result of several mechanisms acting alone or in concert. Ischaemia may cause intestinal injury, which may increase oxygen demands or decrease oxygen supply through reduced intestinal blood flow or oxygen short circuiting in the villus countercurrent exchange. Additionally, ischaemia may cause reperfusion injury, which can also damage the intestinal mucosa.

After hypovolaemia, or reduced cardiac output for other reasons (such as cardiac tamponade), the splanchnic blood supply might be insufficient to meet the tissues needs. It has been shown that the small intestinal blood supply has to be reduced by more than 50% to induce detectable tissue injury.⁶ If the blood flow reduction is less than 50%, compensatory mechanisms will be enough to avert tissue injury. These mechanisms include increased oxygen extraction and oxygen redistribution within the intestinal wall to those areas with a high metabolic demand.

In haemorrhage or cardiac tamponade, or both the reduced delivery to the intestine may create sufficient hypoxia in the superficial part of the gut mucosa to cause tissue injury.⁷ Where ischaemia, however, was induced regionally in the small intestine by lowering the arterial inflow pressure, a very significant redistribution of intramural blood flow provides the mucosa with more oxygen.⁸ Moreover, in sepsis, blood flow to the superficial part of the mucosa has also been shown to be maintained well in animals developing intestinal mucosal injury.⁹ Furthermore, in regional intestinal ischaemia and in septic shock, the intestinal mucosal injury could be prevented by supplying the gut mucosa with small amounts of extra oxygen by intraluminal perfusion with oxygenated saline. Perfusion with nitrogen saturated saline had no such effect.^{9 10}

The unusual situation where the blood supply to the superficial part of the mucosa is fairly well maintained but hypoxic injury still develops in this area within 1–2 hours of shock, may be explained by the peculiar vascular anatomy of the intestinal mucosa. The villus is supplied with a centrally located arterial vessel, which does not arborise until it reaches the tip of the villus. The blood flow is then drained through a subepithelial network of capillaries and venules. The distance between these two

sets of vessels with mainly opposite flow directions is less than 20 μ , which will permit diffusion equilibrium to take place for easily diffusible substances such as oxygen. A limiting factor is the time allowed to obtain diffusion equilibrium. In shock, the blood flow velocity is significantly reduced,¹¹ and the prerequisite for a very effective short circuiting of oxygen at the base of the villi is present. A certain degree of short circuiting of oxygen in the villus countercurrent exchanger has been shown in the resting intestine,¹² and a very low resting pO_2 has been recorded at the tip of the villus.¹³ Thus, an increased effectiveness of the countercurrent exchanger during hypotensive states might well explain an almost anoxic environment at the tip of the villi despite a rather well maintained volume of blood flow to this region.

During recent years an additional mechanism causing impaired tissue oxygenation to the gut during sepsis has been described. Dahn *et al*¹⁴ provided strong indications for an increased oxygen demand of the splanchnic organs during sepsis compared with trauma not complicated by sepsis. Arvidsson *et al*¹⁵ could measure oxygen consumption in the gastrointestinal tract and in the liver during sepsis and haemorrhage. It could be shown that during sepsis induced by faecal peritonitis in pigs, the splanchnic oxygen consumption almost doubled within a couple of hours. For the intestine, the oxygen delivery was judged to be sufficient to meet the total oxygen demands of the tissue, under these experimental conditions. In sepsis, however, the hepatic needs for oxygen exceeded the delivery, and flow dependent hypoxia was expressed as increased lactate production.¹⁵ Recently, it was shown, using the tonometric technique for monitoring intramucosal pH (see later) that mucosal ischaemia occurred rapidly after sepsis despite maintained or increased oxygen delivery and increased oxygen consumption of the gut.¹⁶

The cause of the increased oxygen demand in the splanchnic area in sepsis is not fully understood. From a theoretical point of view, activated resident macrophages and invading white blood cells might account for at least part of the increased demand.

To summarise, in the critically ill, tissue oxygenation of the splanchnic organs is threatened by several simultaneous mechanisms causing intestinal ischaemia. Blood flow delivery to the area might be impaired. The mucosal countercurrent exchanger causes a further decrease of oxygen delivery to the superficial part of the mucosa. In the septic state there is, in addition, an increased demand for oxygen in the splanchnic area. Moreover, sepsis causes decreased extraction and utilisation of oxygen by the tissue.

Reperfusion injury

During the past 10 years, there has been increasing evidence that, in addition to the injury taking place during ischaemia, intestinal injury also takes place at reperfusion.^{17 18} The

- 2 Haglund U. Gastro-intestinal mucosal injury in shock. In: Vincent JL, ed. *Update in intensive care and emergency medicine*. Berlin, Heidelberg: Springer-Verlag, 1991: 154-60.
- 3 Haglund U, Bulkley GB, Granger DN. On the pathophysiology of intestinal ischemic injury. *Acta Chir Scand* 1987; 153: 321-4.
- 4 Chiu CJ, McArdle AH, Brown R, Scott HJ, Gurd FN. Intestinal mucosal lesion in low-flow states. *Arch Surg* 1970; 101: 478-83.
- 5 Park PO, Haglund U. Regeneration of small bowel mucosa after intestinal ischemia. *Crit Care Med* 1992; 20: 135-9.
- 6 Bulkley GB, Kvietys PR, Parks DA, Perry MA, Granger DN. Relationship of blood flow and oxygen consumption to ischemic injury in the canine small intestine. *Gastroenterology* 1985; 89: 852-7.
- 7 Bailey RW, Bulkley GB, Hamilton SR, Morris JB, Haglund UH. Protection of the small intestine from nonocclusive mesenteric ischemic injury due to cardiogenic shock. *Am J Surg* 1987; 153: 108-16.
- 8 Redfors S, Hallbäck DA, Haglund U, Jodal M, Lundgren O. Blood flow distribution, villous osmolality and fluid and electrolyte transport in the cat small intestine during regional hypotension. *Acta Physiol Scand* 1984; 121: 193-209.
- 9 Falk A, Redfors S, Myrvold HE, Haglund U. Small intestinal mucosal lesions in feline septic shock; a study on the pathogenesis. *Circ Shock* 1985; 17: 327-37.
- 10 Haglund U, Abe T, Åhrén C, et al. The intestinal mucosal lesions in shock: I. Studies on the pathogenesis. *Eur Surg Res* 1984; 8: 435-47.
- 11 Haglund U, Jodal M, Lundgren O. The small bowel in arterial hypotension and shock. In: Shephard AP, Granger DN, eds. *Physiology of the intestinal circulation*. New York: Raven Press, 1984: 305-19.
- 12 Kampp M, Lundgren O, Nilsson NJ. Extravascular shunting of oxygen in the small intestine of the cat. *Acta Physiol Scand* 1968; 72: 396-403.
- 13 Bohlen HG. Intestinal tissue pO₂ and microvascular responses during glucose exposure. *Am J Physiol* 1980; 238: H164-71.
- 14 Dahn MS, Lange MP, Lodell K, Hans B, Jacobs LA, Mitchell RA. Splanchnic and total body oxygen consumption differences in septic and injured patients. *Surgery* 1987; 101: 69-80.
- 15 Arvidsson D, Rasmussen I, Almqvist P, Niklasson F, Haglund U. Splanchnic oxygen consumption in septic and hemorrhagic shock. *Surgery* 1991; 2: 190-7.
- 16 Rasmussen I, Haglund U. Early gut ischemia in experimental fecal peritonitis. *Circ Shock* 1992; 38: 22-8.
- 17 Granger DN, Rutili G, McCord JM. Superoxide radicals in feline intestinal ischemia. *Gastroenterology* 1981; 81: 22-9.
- 18 Schoenberg MH, Fredholm B, Haglund U, et al. Studies on the oxygen radical mechanism involved in the small intestinal reperfusion damage. *Acta Physiol Scand* 1985; 124: 581-9.
- 19 Park PO, Haglund U, Bulkley GB, Fält K. The sequence of development of intestinal tissue injury following strangulation ischemia and reperfusion. *Surgery* 1990; 107: 574-80.
- 20 Haglund U, Gerdin B. Oxygen-free radicals (OFR) and circulatory shock. *Circ Shock* 1991; 34: 405-11.
- 21 Bounous G. Acute necrosis of the intestinal mucosa. *Gastroenterology* 1982; 82: 1457-67.
- 22 Parks DA, Williams TK, Beckman JS. Conversion of xanthine dehydrogenase to oxidase in ischemic rat intestine: a reevaluation. *Am J Physiol* 1988; 254: G768-74.
- 23 Montgomery A, Borgström A, Haglund U. Pancreatic proteases and intestinal mucosal injury after ischemia and reperfusion in the pig. *Gastroenterology* 1992; 102: 216-22.
- 24 Fiddian-Green RG. Studies in splanchnic ischemia and multiple organ failure. In: Marston A, Bulkley GB, Fiddian-Green RG, Haglund U, eds. *Splanchnic ischemia and multiple organ failure*. London: Edward Arnold, 1989: 350-63.
- 25 Antonson JB, Boyle CC, Kruithoff KL, Wang H, Sacristan E, Rothchild HR, et al. Validation of tonometric measurement of gut intramural pH during endotoxemia and mesenteric occlusion in the pig. *Am J Physiol* 1990; 259: G519-23.
- 26 Nobumichi H, Masaharu N, Manabu N, et al. Structural and functional alterations in the gut of parenterally or enterally fed rats. *J Surg Res* 1989; 47: 129-33.
- 27 Alverdy JC, Aoys E, Moss GS. Total parenteral nutrition promotes bacterial translocation from the gut. *Surgery* 1988; 104: 185-90.
- 28 Moore FA, Moore EE, Jones TN, et al. TEN versus TPN following major abdominal trauma - reduces septic morbidity. *J Trauma* 1989; 29: 916-23.
- 29 Hwang TL, O'Dwyer ST, Smith RJ, et al. Preservation of the small bowel mucosa using glutamine-enriched parenteral nutrition *Surg Forum* 1986; 37: 56-8.
- 30 Rombeau JL. A review of the effects of glutamine-enriched diets on experimentally induced enterocolitis. *JPEN* 1990; 14: 100S-5S.