INTRODUCTION

Some of the most important research into enteral nutrition in the 1970s and early 1980s concerned the definition of nutritional requirements of starved and injured patients and the most appropriate techniques for giving diets.

These studies showed that the energy requirements of starved and injured patients had previously been overestimated^{1,2} while nitrogen requirements had been underestimated.³ In disease states, it became accepted that the recommended daily allowances for vitamins, trace elements, and haematinics were inadequate and that in the patients requiring predigested chemically defined elemental diets more attention needed to be focused on sodium intakes.⁴ Current research has focused on qualitative aspects of lipid requirements, specifically in regard of the omega 6 to omega 3 fatty acid ratios.

Advances in administration technique led to design changes in nasogastric and nasoenteral feeding tubes,⁵ incorporation of drip chambers in giving sets,⁶ the use of 'closed system' diet reservoirs of up to two litres in volume.⁷ These have led to increased efficiency of enteral feeding as well as to a reduction in side effects or complications. The abandonment of 'starter regimens'⁸ in most patients, is to be welcomed because as a technique it led to reduced efficacy of enteral feeding, particularly in respect of the amount of nutrients given.

A further important advance has undoubtedly been the development of percutaneous endoscopic gastrostomy (PEG) tube placement. Patients requiring longterm enteral feeding (who previously had to suffer the discomforts of longterm nasogastric tube placement) can now be fed more simply and efficiently by gastrostomy. This has been of particular benefit to those patients receiving home enteral nutrition. We anticipate the development of more sophisticated and manageable, narrower bore PEG feeding tubes.

Much of the research described above, was done during a time when enteral nutrition was perceived to be supportive treatment. What is meant by this (in clinical terms) is that provision of nutritional support in itself by preserving the nutritional state and thus preventing malnutrition related complications⁹ gains time for the primary care physician to treat the underlying disease.

Research in the field of enteral nutrition is currently undergoing an interesting change. Investigators now perceive that there are a number of new substrates that are of potential benefit to patients receiving enteral nutrition. While the metabolic response to sepsis and trauma is the framework within which many new substrates are being tested, the fact that the gut is not just an inert tube for digestion and assimilation of food is fundamental to the understanding of the potential of these novel substrates. These include glutamine, ornithine alpha ketoglutarate (OKGA), omega 3 polyunsaturated fatty acids, short chain fatty acids, and nucleosides.

A unifying feature of these novel enteral substrates is their tissue specific effects. In the context of enteral nutrition it is appropriate that they all maintain the integrity of the gastrointestinal tract, either directly (glutamine, short chain fatty acids) or indirectly (OKGA, polyunsaturated fatty acids) by moderating the effects of sepsis or trauma on this organ. Work that has defined the potential benefits of the novel substrates has therefore refocused the attention of investigators to the gut. There is abundant evidence in published works that starvation results in atrophy of the

villous architecture. Similar findings have been described during longterm parenteral nutrition in animals,¹⁰ and in the segments of small intestine surgically bypassed for obesity.¹¹ The lack of luminal nutrition plays a pivotal part in producing these changes, it is assumed, by inhibiting production of factors trophic to the gut. The fact that absence of luminal nutrition has a deleterious effect on intestinal morphology has provided a powerful argument for providing nutritional support wherever possible, by the enteral, rather than the parenteral, route. This, however, does not complete the story. Changes in intestinal morphology have been shown to be associated with changes in barrier function. Thus not only does parenteral nutrition adversely affect the intestinal morphology of experimental animals but it is also associated with the passage of viable bacteria from the gastrointestinal tract to otherwise sterile tissues, such as mesenteric lymph nodes, liver, and spleen.¹² This process of bacterial translocation is not seen to the same extent in animals receiving enteral feeding.¹⁰

Bacterial translocation has now been shown to occur in experimental animals submitted to a pathological stimuli such as burns, trauma, sepsis, mesenteric ischaemia, and intestinal bacterial overgrowth.¹³ Bacterial translocation can be potentiated by protein malnutrition¹⁴ and specifically by essential fatty acid deficiency.¹⁵

There is now experimental evidence to show that partial loss of intestinal barrier function may be reversed by luminal nutrition either in the form of oral nutrition¹⁶ or as suggested in some experiments, by glutamine alone.¹⁷

Although the clinical significance of bacterial translocation is unclear, translocating bacteria have been hypothesised as a potential source of sepsis/sepsis syndrome in critically ill patients. If this hypothesis is correct then the protective effects of luminal nutrition, as seen so far, may turn out to be of unparalleled importance and raise the status of enteral nutrition, at least in critically ill patients, from supportive to primary treatment.

Clearly though, several important questions remain unanswered. The first is that the route by which translocating bacteria reach the mesenteric lymph nodes is uncertain. It follows also that we still have inadequate knowledge as to how or why translocating bacteria exert their deleterious effects. Finally we need to learn much more about the mechanisms participating in the protective effect of novel substrates or luminal nutrition, or both.

Routes taken by translocating bacteria to reach the reticuloendothelial system (RES).

The Figure summarises the various routes by which translocating bacteria reach the reticuloendothelial system. Do the factors associated with bacterial translocation (such as malnutrition, lack of luminal nutrition, sepsis or trauma of different kinds) lead to structural changes within the intestinal epithelium at paracellular or transcellular sites, which in turn lead to a failure of mechanical barrier function? Alternatively, is bacterial translocation a normal phenomenon such that under the above conditions, tissue macrophages that engulf bacteria and transport them to the mesenteric lymph nodes fail to accomplish intracellular killing? The first two sessions of the meeting brought together international experts in gut morphology, function, and immunology who attempted to provide answers to those questions. If, as it has been proposed, normal intestinal bacteria can initiate systemic disease as a consequence of translocating the gut mucosal barrier to the reticuloendothelial system, it is important to discover the mechanisms by which this occurs. In this regard, there would seem to be much interest in the role of macrophage secreted cytokines, not only in the pathogenesis of the sepsis/sepsis syndrome but also in the initiating process of intestinal bacteria translocation. Professor van Leeuwen considered these points in his paper discussing the 'clinical significance of bacterial translocation'.

One of the clinically most pertinent questions arising from the recent work in experimental animals, is that intestinal bacterial translocation can be significantly attenuated or prevented by different forms of nutritional support. It is not only important to find out if 'standard' enteral nutrition or food itself has a protective effective, but also if this effect can be significantly enhanced by exposing the luminal, or indeed serosal aspects of the gut to 'unphysiological' loads (concentration $\times$ rate) of nutritional or 'novel' substrates. Authors in the third session discussed evidence as to whether short chain fatty acids, glutamine, medium chain triglyceride arginine, and its metabolites and nucleosides can influence gut morphology and function.

In other clinical areas, however, the case for enteral nutrition as primary therapy is proved, no more so than in the management of Crohn's disease and the short bowel syndrome. Not discussed here are also the benefits of enteral nutrition as a primary therapy of chronic porto-systemic encephalopathy in patients with chronic liver disease.¹⁸ The reader will also benefit from the advances that have been made in the use of enteral nutrition as primary therapy in radiation injury and gut ischaemia as

well as in cancer – the latter perhaps more demanding of a conference of its own.

The drawing together of small groups of multi-disciplinary international experts occurs far too seldom these days. We hope that the proceedings of this meeting will stand the test of time.

D B A SILK
G K GRIMBLE

Department of Gastroenterology and Nutrition,
Central Middlesex Hospital NHS Trust,
Acton Lane, London NW10 7NS

This conference was initiated by Dr Sam Hull of Abbot Ross.

- MacFie J. Active metabolic expenditure of gastroenterological patients receiving intravenous nutrition. *JPEN* 1984; 8: 371–6.
- Jeejeebhoy KN. Energy metabolism in the critically ill. In: Garrow JS, Halliday D, eds. *Substrate and energy metabolism in man*. London: John Libby, 1985; 1: 93–101.
- Rees RGP, Cooper TM, Beetham R, Frost PG, Silk DBA. Influence of energy and nitrogen contents of enteral diets on nitrogen balance; a double blind prospective controlled clinical trial. *Gut* 1989; 30: 123–9.
- Spiller RC, Jones BJM, Silk DBA. Jejunal water and electrolyte absorption from five proprietary enteral feeds in man: importance of sodium content. *Gut* 1987; 28: 681–7.
- Silk DBA, Rees RG, Keohane PP, Attrill H. Clinical efficacy and design changes in 'Fine Bore' nasogastric feeding tubes: a seven year experience involving 109 intubations in 403 patients. *JPEN* 1987; 11: 378–84.
- Payne-James JJ, Rana SK, Bray MJ, McSwiggan DA, Silk DBA. Retrograde (ascending) bacterial contamination of enteral diet administration systems. *JPEN* 1992; 16.4: 369–73.
- Rees RGP, Ryan J, Attrill H, Silk DBA. Clinical evaluation of two-litre prepackaged enteral delivery system: a controlled trial. *JPEN* 1988; 12: 274–7.
- Keohane PP, Attrill H, Love M, Frost P, Silk DBA. Relation between osmolality of diet and gastrointestinal side effects in enteral nutrition. *BMJ* 1984; 288: 678–80.
- Rana SK, Bray J, Menzies-Gow N, Jameson J, Payne-James JJ, Frost P, et al. Short term benefits of post operative oral dietary supplements in surgical patients. *Clinical Nutrition* 1992; 11: 337–44.
- Johnson LR, Copeland EM, Dudrick SJ, et al. Structural and hormonal alterations in the gastrointestinal tract of parenterally fed rats. *Gastroenterology* 1975; 68: 1177–83.
- Daly JM, Castro GA, Akhtar M, Dudrick SJ. Morphologic and biochemical intestinal changes after jejuno-ileal by-pass. [Abstract]. *Gastroenterology* 1977; 72: 1042.
- Alverdy JC, Aoy E, Moss GS. Total parenteral nutrition promotes bacterial translocation from the gut. *Surgery* 1988; 104: 185–90.
- Wells CL, Maddaus MA, Simmons RL. Proposed mechanism for the translocation of intestinal bacteria. *Rev Infect Dis* 1988; 10: 958–79.
- Deitch EA, Winterton J, Li M, et al. The gut as a portal of entry for bacteremia: role of protein malnutrition. *Ann Surg* 1987; 205: 681–92.
- Barton RG, Cerra FB, Wells CL. Effect of a diet deficient in essential fatty acids on the translocation of intestinal bacteria. *JPEN* 1992; 16: 122–8.
- Wells CL, Jechorek RP, Evlansden SG, et al. The effect of dietary glutamine and dietary RNA on ileal flora, ileal histology and bacterial translocation in mice. *Nutrition* 1990; 6: 70–5.
- Souba WW, Klimberg VS, Hantakeaki RD, et al. Oval glutamine reduces bacterial translocation following abdominal radiation. *J Surg Res* 1990; 48: 1–5.
- Horst D, Grace ND, Conn HO, et al. Comparison of dietary protein with an oral, branched chain-enriched amino acid supplement in chronic portal systemic encephalopathy. A randomised controlled trial. *Hepatology* 1984; 4: 279–87.