analogous to that in the posterior nerve root ganglion in herpes zoster, came closer to the mark than have gastroenterologists since.

MICHAEL ATKINSON

Department of Surgery, University of Nottingham,
University Hospital,
Queen's Medical Centre,
Nottingham NG7 2UH

- 1 Hurst AF. Essays and addresses on digestive and nervous diseases and on Addison's anaemia and asthma. London: Heinemann, 1924.
- 2 Rake GW. On the pathology of achalasia. *Guy's Hospital Reports* 1927; 77: 141-50.
- 3 Cassella RR, Brown AL, Sayre GP, Ellis FH Jr. Achalasia of the esophagus: pathologic and etiologic considerations. *Ann Surg* 1964; 160: 474-86.

- 4 Smith B. The neurological lesion in achalasia of the cardia. *Gut* 1970; 11: 388-91.
- 5 Elder JB, Gillespie G. The vagus and achalasia. *Gut* 1969; 10: 1045.
- 6 Atkinson M, Ogilvie AL, Robertson CS, Smart HL. Vagal function in achalasia. *Q J Med* 1987; 63: 297-303.
- 7 Raju GS, Rao PN, Broor SL. Effect of sham feeding on gastric acid secretion in patients with achalasia. *Gastroenterology* 1989; 96: 989.
- 8 Holloway RH, Dodds WJ, Helm JF, et al. Integrity of cholinergic innervation to the lower esophageal sphincter in achalasia. *Gastroenterology* 1986; 90: 924-9.
- 9 Westley CR, Herbst JJ, Goldman S, Wisek WC. Infantile achalasia: inherited as an autosomal recessive disorder. *J Pediatr* 1975; 87: 243-6.
- 10 Mayberry JF, Atkinson M. A study of swallowing difficulties in first degree relatives of patients with achalasia. *Thorax* 1985; 40: 391-3.
- 11 Robertson CS, Martin BAB, Atkinson M. Varicella-Zoster virus DNA in the oesophageal myenteric plexus in achalasia. *Gut* 1993; 34: 299-302.
- 12 Benjamin SB, Castell DO. Achalasia and Hodgkin's disease: a chance association? *J Clin Gastroenterol* 1981; 3: 175-8.

International gastroenterology

Gastroenterology in China

With the establishment of New China in 1949, a nationwide educational programme in public health and sanitation was adopted. This was not confined to metropolitan cities and towns; it was even more intensively carried through in rural areas where measures were instituted to control prevalent parasitic diseases. The first important campaign, beginning as early as December 1949, was against schistosomiasis and this showed the wretched conditions then prevailing in rural districts. Our central government emphasised the importance of this campaign by the presence of the minister of public health in person, together with the governors of 13 provinces and one cosmopolitan city (Shanghai), on a leadership board to organise, firstly, the anti-schistosomiasis campaign and, secondly, the eradication of three other important parasitic diseases: kala-azar, filariasis, and malaria and, at the same time, to improve sanitation in the countryside. It was this board that rapidly appreciated the extent to which the vast farming districts suffered from a dire shortage of medical personnel. It was clear that this could not be remedied by relocation of the medical practitioners who were, at that time, mostly working in large cities, and it had become obvious that the small numbers of trained doctors were unequal to the task in hand.

Therefore, the board decided to enrol young people, who had graduated from primary and junior middle schools in each of the provinces, to be trained on a one year course on the use of anti-parasitic drugs, mainly against schistosomiasis, and also to survey sanitation facilities. This 'army' of young people was known as the 'barefoot doctors'. It has to be remembered that, in the period between 1950 to 1956, there was a continuing shortage of well trained doctors especially in the field of parasitic disease and it was this shortage that was filled by the 'barefoot' personnel. There are no more 'barefoot doctors', but the work done by these young people is recognised and honoured to this day. The campaign against parasitic diseases was a success, as judged by a considerable reduction in the prevalence of schistosomiasis, filariasis, and malaria and the near complete eradication of kala-azar.

Before 1975, there was no organised society of gastroenterology in China but gastroenterology, as a speciality, was only recognised as a branch of the Chinese Society of

International Medicine (CSIM). Through the great efforts and wise counsel of the late distinguished Professor Xiao-Qian Zhang, of Peking Union Medical College, the Chinese Gastroenterology Society was established in 1978. Since then, the Society has flourished and prospered so rapidly that almost every province as well as important cosmopolitan cities in China will have a subsidiary society and one or more institutes or centres that are affiliated to the Society where clinical gastroenterology and research are carried out; membership of the Chinese Gastroenterological Society has now reached a total of about 5000. There is a national assembly every three years to highlight and discuss the work being done in various parts of China. Since 1981, there has been an official journal of the Society named *The Chinese Journal of Digestion*, originally a quarterly and now a bimonthly journal. Within the last five years, there have been several successful large international symposia on gastroenterology in China.

The era of modern gastroenterology in China began with the introduction of endoscopy in the 1970s. There are now several endoscopes of Chinese manufacture that have been distributed to large and small hospitals in the nation. Every hospital has a gastrointestinal clinic with a specialist staff and all young specialists are familiar with diagnostic upper gastrointestinal endoscopy. Therapeutic endoscopy is also practised but mainly in the gastroenterology centres in large cities.

Gastrointestinal diseases are much influenced by geography, personal habits, customs, and ethnic origin. In China, some diseases are more prevalent such as chronic gastritis resulting from *Helicobacter pylori*, gastric cancer, viral hepatitis (especially hepatitis B), and hepatoma but inflammatory bowel disease, colorectal cancer, and pancreatic cancer are less common than in Western countries. Because gastric cancer and hepatoma threatens the lives of so many Chinese people, there is considerable effort by Chinese gastroenterologists on research into the treatment and, more important, the prevention of these malignant tumours.

SHAO-JI JIANG

Shanghai Institute of Digestive Disease,
145 Shan-Dong Zhong Road,
Shanghai 200001,
China