CASE REPORTS

Avulsion of short gastric arteries caused by vomiting

N Hayes, P D Waterworth, S M Griffin

Abstract

A case is presented describing a new, potentially life threatening complication of vomiting after a 21 year old man presented in shock with a haemoperitoneum caused by violent, self induced emesis.

(Gut 1994; 35: 1137-1138)

Forceful or prolonged retching, from whatever cause, can lead to life threatening oesophago-gastric complications. The commonest, a Mallory-Weiss tear,¹ often presenting as an upper gastrointestinal haemorrhage, arises from a breach of the mucosa around the oesophagogastric junction. Occasionally there is a full thickness perforation, usually of the left wall of the oesophagus several centimetres above the cardia, which was first described by Boerhaave in the eighteenth century.² We describe a hitherto unreported serious local complication of vomiting occurring after self induced emesis.

Case report

A 21 year old man was referred as an emergency by his general practitioner with acute abdominal pain. The preceding evening he had consumed eight pints of lager followed by a take away Chinese meal. At 7 am on the day of admission, he awoke feeling uncomfortable and bloated, so he forced himself to vomit by putting his fingers down his throat. The resulting retching was surprisingly violent, and epigastric pain developed shortly afterwards. When he later visited his doctor, the pain had radiated to both shoulders and he was referred to casualty. On admission, the patient recounted the above history and was clear that he had suffered no recent external trauma. Examination showed the patient to be afebrile, but pale and distressed with a pulse rate of 112 bpm and blood pressure 75/35 mm Hg. His hypotension responded to an intravenous 'fluid challenge' with 1000 ml colloid solution. There was tenderness in both the epigastrium and the right iliac fossa. He had a white cell count of $20.4 \times 10^9/l$, haemoglobin of 121 g/l, and haematocrit of 36.1%. Serum urea, amylase, and electrolyte tests were normal. Although subphrenic gas was absent on an erect chest film, a diagnosis of perforated viscus was made.

At laparotomy, a haemoperitoneum of about 1.5 litres was discovered. There was a short gastrosplenic ligament which, with the greater omentum, was torn at its reflection with the stomach, effectively mobilising the stomach along the greater curvature (Figure). Four short gastric arteries and branches of the gastroepiploic artery had been torn, these vessels were in the main sealed by a fresh clot. After controlling the damaged vessels with polyglactin ligatures, inspection showed no damage to the stomach muscle and as no blood was seen in aspirates from the nasogastric tube, endoscopy was not performed. The abdomen was closed leaving a drain at the splenic hilum. The patient made an uneventful recovery.

Discussion

Injury to the upper gastrointestinal tract from external violence can take a number of forms,³ but recognised lesions caused by vomiting are mainly partial or full thickness mural lacerations.^{1,2} In this case, there was no clinical

Department of
Surgery, Newcastle
General Hospital,
Newcastle upon Tyne
N Hayes
P D Waterworth
S M Griffin

Correspondence to:
Mr N Hayes, Newcastle
General Hospital, Westgate
Road, Newcastle upon Tyne
NE4 6BE.

Accepted for publication
23 November 1993

Schematic representation of extent of injury.

evidence of injury to any aspect of the oesophageal or gastric wall and yet a potentially life threatening laceration of the gastrosplenic omentum resulted. We believe that the mechanism of injury may have been either an intussusception or partial volvulus of the greater curve of stomach caused by violent retching, with movement hampered by a short gastrosplenic ligament that subsequently tore.

Although the case described is unusual, there are previously reported cases of spontaneous haemoperitoneum.⁴ We believe that if a patient presents with a history of forceful retching followed by abdominal pain and clinical signs of haemorrhage, the possibility of a tear in the gastrosplenic omentum with intraperitoneal haemorrhage should be

considered. There seems therefore to be three serious gastro-oesophageal complications of vomiting: (a) Mallory-Weiss tear, (b) Boerhaave's syndrome, and (c) avulsion of the short gastric vessels.

The authors wish to thank Alan Waller of the University of Newcastle upon Tyne Audio Visual Centre for preparing the illustration.

- 1 Mallory G, Weiss S. Hemorrhages from lacerations of the cardiac orifice of the stomach due to vomiting. *Am J Med Sci* 1929; 178: 506-15.
- 2 Barrett N. Spontaneous perforation of the oesophagus. Review of the literature and report of three new cases. *Thorax* 1946; 1: 48-70.
- 3 Hulewicz B. Gastric trauma following cardiopulmonary resuscitation. *Med Sci Law* 1990; 30: 149-52.
- 4 Sanderson PL, Davies MFP, Howat JMT. Idiopathic spontaneous haemoperitoneum. *Postgrad Med J* 1988; 64: 322-4.