LETTERS TO THE EDITOR

Hypolactasia and protection against *Plasmodium* spp infection in *Homo sapiens*

EDITOR,—Anderson and Vullo's hypothesis (*Gut* 1994; 35: 1487–9) is based on the contention that hypolactasia is causatively associated with riboflavin deficiency (consequent upon low milk intake), which has evolutionarily produced a survival advantage against human *Plasmodium* spp infection. Available evidence (quoted by the authors) indicates, however, that such a deficiency exerts only a very mild influence on the metabolism of the malarial parasite.

The overwhelming consensus of current opinion is that the hypolactasia phenotype represents the normal state for *Homo sapiens*¹ (and virtually all other mammals), and that persistence of the enzyme into adult life in certain groups before 'artificial' mass population movements (for example, northern Europeans to the USA and Australia) resulted from a significant survival advantage; by involving malaria as the major 'driving force', Anderson and Vullo are thus interpreting the scenario in 'reverse' fashion.

Plasmodium spp are extremely ancient organisms²; before they infected the earliest mammals, they utilised more primitive organisms, including reptiles, birds, etc, as reservoir hosts. The Anderson and Vullo hypothesis therefore implies that the very widespread prevalence of the hypolactasia phenotype in the animal kingdom developed as an exceedingly early protective mechanism against *Plasmodium* spp, and persistence of the enzyme into adult life only developed very much later in areas where *Plasmodium* spp was not a major environmental hazard affecting survival. The Hamitic tribes of east, west, and central Africa (most adults possess persistence of the enzyme into their adult life³) presumably migrated into northern Africa comparatively recently (within the last few thousand years) – and were thence 'unprotected' against *Plasmodium* spp. How then, did these Hamitic 'pastoralists' survive *P falciparum* infection? I know of no evidence that Hamites are, at the present day, more prone to *P falciparum* infection compared with Bantus with the exception that they lack certain major genetic diseases^{4,5} (see later)! *P vivax* infection was certainly a problem in northern Europe until recent times; however, it never caused the widespread acute mortality associated with human *P falciparum* infection – which probably originated in Africa also comparatively recently.

The approximate parallel outlined in the hypothesis between the geographical areas incorporating the hypolactasia phenotype and those in which major haemoglobinopathies (which clearly possess a selective advantage against *P falciparum* infection) are comparatively common, is of interest.^{4,5} But even here the degree of protection is only partial; riboflavin deficiency has probably exerted a comparatively minimal, and certainly a lesser, effect in terms of selective Darwinian pressure.

Therefore, although this hypothesis seems superficially attractive, I have grave doubts

that it explains the human lactase polymorphism more satisfactorily than several of the previous theories.¹

G C COOK

Hospital for Tropical Diseases,
St Pancras Way,
London NW1 0PE

- 1 Cook GC. Hypolactasia: geographical distribution, diagnosis, and practical significance. In: Chandra RK, ed. *Critical reviews in tropical medicine* 2. New York: Plenum Publishing, 1984: 117–39.
- 2 Bruce-Chwatt LJ. History of malaria from pre-history to eradication. In: Wernsdorfer WH, McGregor IA, eds. *Malaria: principles and practice of malariology*. Edinburgh: Churchill Livingstone, 1988: 1–59.
- 3 Cook GC, Kajubi SK. Tribal incidence of lactase deficiency in Uganda. *Lancet* 1966; i: 725–30.
- 4 Fleming AF. Hematologic diseases. In: Strickland GT, ed. *Hunter's tropical medicine*. 7th ed. Philadelphia: WB Saunders, 1991: 36–64.
- 5 Bruce-Chwatt LJ. Paleogenetics and paleoepidemiology of primate malaria. *Bull WHO* 1965; 32: 363–87.

Reply

EDITOR,—We thank Dr G C Cook for his interest in our hypothesis and would like to reply to some of his points.

Dr Cook says that 'available evidence (quoted by the authors) indicates, however, that such a deficiency exerts only a very mild influence on the metabolism of the malarial parasite'. We believe that the crucial point is not whether the influence of flavin deficiency on the parasite metabolism is 'mild' or 'very mild', but whether its effect is sufficient to prevent excessive multiplication of the parasite as it has been shown in vivo and quoted in our paper, so protecting the host from death.

Dr Cook also says that our hypothesis implies that 'persistence of the enzyme into adult life only developed very much later in areas where *Plasmodium* spp was not a major environmental hazard affecting survival'. It is possible that we have not made it clear that the basis of our hypothesis is that the 'persistent' phenotype was the 'wild type' existent initially, and therefore we did not envisage that it 'developed' later but that it continued to predominate in those areas.

Even if, as it seems today, there is no evidence that the Hamite tribe (who possess the 'persistent' phenotype) are more prone to malaria than the Bantus, this does not necessarily show that our hypothesis is wrong, because they could be protected by other factors, known or unknown.

We hope that the validity of our hypothesis might be supported if further detailed demographic studies are done on the 'persistent' and hypolactasia phenotypes in relation to malaria within suitable countries. The main purpose of publishing our hypothesis was to encourage such studies.

C VULLO

B ANDERSON

Divisione Pediatrica,
Arcispedak S Anna, via Savonarola 15,
44100 Ferrara, Italy

In vitro dissolution of cholesterol biliary stones

EDITOR,—We read with great interest the article by Bergman *et al* (*Gut* 1994; 35: 1653–8) who reported on in vitro cholesterol gall stone dissolution with methyl-*tert*-butyl ether (MTBE) and ethyl propionate in the presence of bile with or without dimethylsulphoxide (DMSO). We would like to

comment further on the important role of DMSO on stone dissolution.

We have experience with the clinical use of different solvents containing DMSO. Based on previous studies by Montet's group in which it was shown that the addition of DMSO to MTBE improved the dissolution rate of cholesterol¹ or brown pigment stones,² we have recently studied the bile duct stone dissolving capacity of multicomponent solvents in humans.³ In these mixtures, DMSO was present both in solvent 1 (aqueous glycine-NaOH buffer solution of EDTA, sodium deoxycholate and 30% DMSO) and in solvent 2 (DMSO/MTBE 70/30). Solvent mixtures were infused continuously and alternately for 16–24 hours through a nasobiliary catheter placed into the common bile duct over the stone(s). In 20 of 22 patients with pigment or mixed biliary duct stone(s), or both, which were too large to be removed after endoscopic sphincterotomy, a stone free state (nine of 22) or good disintegration of stones (11 of 22) was achieved with these solvents. Only mild toxic side effects were seen during treatment. These mixtures containing DMSO proved effective enough to dissolve not only cholesterol but also pigment or mixed stones, or all three.

Our recent in vitro investigation in the field of contact dissolution therapy of bile duct stones provides further evidence on the beneficial effect of DMSO.⁴

T TAKACS

First Department of Medicine,
Albert Szent-Györgyi Medical University,
Szeged, Hungary

J C MONTET

INSERM, 46 Boulevard de la Gaye,
13009, Marseille, France

- 1 Dai KY, Montet JC, Zhao XM, Amic J, Choux R. Dissolving agents of human mixed cholesterol stones. *Gastroenterol Clin Biol* 1988; 12: 312–9.
- 2 Dai KY, Montet JC, Zhao XM, Amic J, Montet AM. Dissolution of human brown pigment biliary stones. *J Hepatol* 1989; 9: 301–11.
- 3 Takacs T, Hajnal F, Nagy I, Montet JC, Lonovics J. Effective dissolution therapy of bile duct stones with a new multicomponent solvent. *European Journal of Gastroenterology and Hepatology* 1993; 35: 867–70.
- 4 Guitaoui M, Montet AM, Takacs T, Montet JC. Contact solvents for common bile duct stones. Studies in an in vitro system. *Liver* (in press).

Reply

EDITOR,—We thank Drs Takacs and Montet for their comments. Although our interest with the clinical use of topical dissolution therapy has mainly focused on the treatment of gall bladder stones, we would like to comment on side effects and efficacy of topical dissolution for bile duct stones.

We do agree with Takacs and Montet that the multilayered composition of bile duct stones make the combined use of DMSO, MTBE, or EDTA, or all three, a logical choice. Indeed, the use of DMSO for dissolving bile duct stones in humans has been described before but reports have been anecdotal.^{1–3} We are still very reluctant to infuse potentially toxic agents into human bile ducts. In contrast with topical dissolution of gall bladder stones, where the gall bladder provides a reservoir enabling actual extraction of the solvent after infusion, topical dissolution of ductal stones lacks the opportunity to recover the solvent from the patient. This increases the risk of toxic side effects. We and others⁴ have found MTBE to produce much more local (haemorrhagic duodenitis) and systemic side effects (somnolentia,