LETTERS TO THE EDITOR

Association of *Helicobacter pylori* infection, lymphoid follicles, and lymphocytic gastritis: a risk factor for the development of primary gastric lymphoma?

EDITOR,—A Miettinen *et al* have published (*Gut* 1995; 37: 471–6) an interesting article on the prevalence of lymphocytic gastritis in primary gastric lymphoma (32% of the cases studied). The authors ended by suggesting that the lymphocytic gastritis is a possible precursor state of gastric lymphoma.

This hypothesis can be proposed because the intraepithelial lymphocytes could represent an overflow of tumour associated T lymphocytes, necessary for the growth of the lymphoma.¹ Alternatively, they might play a part in negative regulation of the immune response.

In the light of this and other findings,² we suggest that the 'triad' formed by lymphocytic gastritis, lymphoid follicles, and *H. pylori* infection, could represent a potential risk factor for the development of primary gastric lymphoma. On the other hand, in such cases the sensitive polymerase chain reaction method for immunoglobulin heavy chain rearrangement could be useful to detect a possible monoclonal cell population (histologically undetectable) among reactive inflammatory cells.

Furthermore, as it is possible to obtain the disappearance of lymphoid follicles after anti-*H. pylori* treatment,³ the behaviour of *H. pylori* positive lymphocytic gastritis after antimicrobial treatment should be further investigated. This is important to establish whether an anti-*H. pylori* treatment is necessary to eliminate the risk factor represented by the 'triad', to prevent the development of primary gastric MALT-lymphoma.

G CAMMAROTA

A TURSI

G FEDELI

G GASBARRINI

Department of Internal Medicine,
Catholic University, Rome

- 1 Hussell T, Isaacson PG, Crabtree JE, Spencer J. The response of cells from low-grade B-cell gastric lymphoma of mucosa associated tissue to *Helicobacter pylori*. *Lancet* 1993; 342: 571–4.
- 2 Wotherspoon AC, Ortiz-Hidalgo C, Falzon MF, Isaacson PG. *Helicobacter pylori*-associated gastritis and primary B-cell gastric lymphoma. *Lancet* 1991; 338: 1175–6.
- 3 Cammarota G, Tursi A, Montalto M, Papa A, Branca G, Vecchio FM, *et al*. Prevention and treatment of low-grade B-cell primary gastric lymphoma by anti-*H. pylori* therapy. *J Clin Gastroenterol* 1995; 21: 118–22.

Colonocyte metabolism

EDITOR,—It was a great pleasure to read Drs Clausen and Mortensen's paper (*Gut* 1995; 37: 684–9) on colonocyte metabolism, particularly as Dr Clausen spent some time in our laboratory working with colonocytes. Careful reading of Dr Clausen's results makes it difficult to conclude that butyrate oxidation is not defective in ulcerative colitis (UC).

Their results of butyrate oxidation in UC (at 2 mmol/l butyrate) indicates a mean rate of about 190 nmol/min/g (dry weight) of ¹⁴CO₂ formation, which is similar to values originally reported in UC,¹ which varied between 90 nmol/min/g (dry weight) in acute colitis to 560 nmol/min/g (dry weight) in quiescent colitis. At complete odds with our published results are Clausen's control values of butyrate oxidation in cases unaffected by disease. Clausen's values, again at 2 mmol/l butyrate are about 175 nmol/min/g (dry weight) whereas our original values were 1090 nmol/min/g (dry weight). Control values of similar magnitude have been consistently published from our laboratory over a number of years. Clausen's values, which are about 6–23 times lower than ours, suggest inadequately metabolising cells, the reason for which is not immediately evident from their paper. The values of ATP produced by colonocytes also are low and parallel the low values of ¹⁴CO₂ recorded by Clausen and Mortensen. A number of points are important in colonocyte preparation: (a) constituents and pH of electrolyte solutions, (b) the presence of calcium and albumin, (c) the concentration of dithiothreitol, (d) the warm ischaemia time of collected tissue, (e) the extent of trauma in disaggregating cells, (f) a critical weight ±10–15 mg (dry weight) of colonocytes in each experiment, and (g) the specific activities of radiolabelled substrates must not be too high. If these factors were observed, we are at a loss to understand the poor oxidative metabolic performance now reported by Clausen and Mortensen. Under optimal oxidative conditions a defect in butyrate metabolism is detectable in UC. To date, two reports^{1,2} indicate a metabolic defect of butyrate oxidation in UC and two (ref 3 and Clausen's paper) 'fail' to show such a change. Metabolic experimentation requires rigorous attention to detail and familiarisation with general biochemical methodology. We have repeatedly checked for failure of butyrate oxidation in UC and have been able to show such a defect in all colonic tissue obtained at operation.

W E W ROEDIGER

S MILLARD

Cell Physiology Laboratory,

Department of Surgery,

University of Adelaide at The Queen Elizabeth

Hospital,

Adelaide, SA 5011, Australia

- 1 Roediger WEW. The colonic epithelium in ulcerative colitis: an energy deficiency disease. *Lancet* 1980; ii: 712–5.
- 2 Chapman MAS, Grahn MF, Boyle MA, Hutton M, Rogers J, Williams NS. Butyrate oxidation is impaired in the colonic mucosa of sufferers of quiescent ulcerative colitis. *Gut* 1994; 35: 73–6.
- 3 Finnie IA, Taylor BA, Rhodes JM. Ileal and colonic epithelial metabolism in quiescent ulcerative colitis: increased glutamine metabolism in distal colon but no defect in butyrate metabolism. *Gut* 1993; 34: 1552–8.

Reply

EDITOR,—We thank Dr Roediger for his comments on our publication on human colonocyte metabolism of short chain fatty acids (SCFAs) in which we were unable to confirm a defective oxidation of butyrate in ulcerative colitis (UC). Dr Roediger has previously shown that butyrate oxidation to CO₂ is decreased in colonocytes obtained from patients with acute and quiescent UC, and that this impairment correlates with the

state of the disease.¹ In reference to luminal concentrations of butyrate, the experiments by Dr Roediger were all performed in the presence of 10 mmol/l butyrate¹ whereas our experiments were carried out in the presence of concentrations ranging from 0.125 to 2 mmol/l. The resultant curves obeyed Michaelis-Menten kinetics, and the kinetic parameters, V_{max} (maximum rate of metabolism), and K_m (the substrate concentration at which the reaction has half its maximum value) were calculated by computer fitting of the data to a Michaelis-Menten plot. As pointed out by Dr Roediger our maximum rates (V_{max}) of CO₂ production from butyrate in controls (187 (24) nmol/min/g dry weight; mean (SEM)) are approximately sixfold lower than the control values reported in his original study,¹ and Dr Roediger speculates that this may be due to inadequately metabolising cells.

In 1990, Dr Roediger generously invited us to work with colonocytes in his laboratory, and we have used the same method of preparing isolated colonic epithelial cells from resected bowel specimens as originally described by Drs Roediger and Truelove² with the exception that hyaluronidase was omitted. Our values of CO₂ production from butyrate in controls ranged from 114 to 333 nmol/min/g dry weight in controls and from 90 to 400 nmol/min/g dry weight in patients with UC. If the lower values of butyrate oxidation in controls compared with the values found by Dr Roediger were due to inadequately metabolising cells, a greater variation might have been expected. Regarding the dry weight of epithelial cells, our cell preparations contained 9.1 (0.9) mg/ml (mean (SEM)), and no significant correlation between V_{max} values of CO₂ production from butyrate and dry weight of epithelial cells was found ($p=0.19$; Spearman's rank correlation test). The warm ischaemia time of collected tissue is difficult to determine exactly, especially in UC patients undergoing colectomy where preselection of a specific colonic section is more troublesome, but CO₂ production from SCFAs and glucose in our study was linear over 60 minutes in cell suspensions obtained from both UC patients and controls indicating efficient metabolic performance during the incubation period. In rats, we have evaluated the time of ischaemia on colonocyte metabolism by removing the colons for 30 minutes before cell preparation and found no effect on CO₂ production from butyrate (unpublished data). Two further studies have investigated the colonic epithelial metabolism of butyrate in UC. As these studies used a different technique (biopsy specimens), the numerical values cannot be directly compared with the values obtained by use of isolated colonocytes, but one study was in favour of a defective oxidation of butyrate in UC³ and one was not.⁴ Further experimental work in this area seems to be required to evaluate the role of colonic metabolism of SCFAs in the pathogenesis of UC.

M R CLAUSEN

P B MORTENSEN

Department of Medicine A 2102,

Division of Gastroenterology,

Rigshospitalet, University of Copenhagen,

Blegdamsvej 9, DK-2100 Copenhagen Ø,

Denmark

- 1 Roediger WEW. The colonic epithelium in ulcerative colitis: an energy-deficiency disease? *Lancet* 1980; ii: 712–5.
- 2 Roediger WEW, Truelove SC. Method of preparing isolated colonic epithelial cells (colonocytes) for metabolic studies. *Gut* 1979; 20: 484–8.