Changes in the enterocyte cytoskeleton in newborn rats exposed to ethanol in utero

J F Montes, G Estrada, M D López-Tejero, J García-Valero

Abstract

Background—Cytoskeletal changes after longterm exposure to ethanol have been described in a number of cell types in adult rat and humans. These changes can play a key part in the impairment of nutrient assimilation and postnatal growth retardation after prenatal damage of the intestinal epithelium produced by ethanol intake.

Aims—To determine, in the newborn rat, which cytoskeletal proteins are affected by longterm ethanol exposure in utero and to what extent.

Animals—The offspring of two experimental groups of female Wistar rats: ethanol treated group receiving up to 25% (w/v) of ethanol in the drinking fluid and control group receiving water as drinking fluid.

Methods—Single and double electron microscopy immunolocalisation and label density estimation of cytoskeletal proteins on sections of proximal small intestine incubated with monoclonal antibodies against actin, α -tubulin, cytokeratin (polypeptides 1, 5, 6, 7, 8, 10, 11, and 18), and with a polyclonal antibody anti- β 1,4-galactosyl transferase as trans golgi (TG) or trans golgi network (TGN) marker, or both. SDS-PAGE technique was also performed on cytoskeletal enriched fractions from small intestine. Western blotting analysis was carried out by incubation with the same antibodies used for immunolocalisation.

Results—Intestinal epithelium of newborn rats from the ethanol treated group showed an overexpression of cytoskeletal polypeptides ranging from 39 to 54 kDa, affecting actin and some cytokeratins, but not tubulin. Furthermore, a cytokeratin related polypeptide of 28–29 kDa was identified together with an increase in free ubiquitin in the same group. It was noteworthy that actin and cytokeratin were abnormally located in the TG or the TGN, or both.

Conclusions—Longterm exposure to ethanol in utero causes severe dysfunction in the cytoskeleton of the developing intestinal epithelium. Actin and cytokeratins, which are involved in cytoskeleton anchoring to plasma membrane and cell adhesion, are particularly affected, showing overexpression, impaired proteolysis, and mislocalisation.

(Gut 1996; 38: 846–852)

Longterm exposure to ethanol in utero results in postnatal growth retardation in the rat,¹ which could be related to prenatal damage^{2,3} induced by the high ethanol concentration found in the fetal gastrointestinal lumen.⁴ However, the mechanism by which ethanol impairs nutrient assimilation in the neonates is not well characterised. Enteral nutrition is critically dependent on the functioning of intestinal epithelium, based on (a) the organisation of membrane domains, (b) the transport pathways across these membranes, and (c) the maintenance of epithelial sheet integrity by cell adhesion. Appropriate nutrient transport in epithelial cells requires the development and maintenance of precise and polarised distributions of proteins and lipids, as well as the organisation and maintenance of specialised cell-cell and cell-substratum contacts.^{5–7}

Ethanol and its metabolic derivatives interact with proteins and lipids, modifying their functional properties. Several ethanol induced molecular and cellular phenomena have been described in alcoholic adults and cell cultures, such as decreased membrane bound enzyme activities,^{8,9} intracellular accumulation of secretion proteins,^{9,10} changed gene expression,^{10–13} inhibition of proteolysis,¹⁴ covalent binding with lipids^{15–17} and proteins,^{18,19} and inhibition of cell-cell adhesion.²⁰ As cytoskeletal integrity is a prerequisite for many of these cell functions, several authors have investigated the interactions between ethanol and the cytoskeletal proteins.^{21,22} Pinazo-Durán *et al*²³ found ethanol induced changes in cytoskeletal proteins such as GFAP in astrocytes, with occasional disorganisation of the entire cytoskeleton. Reddy *et al*⁸ reported changes in actin associated proteins such as myosin and Adickes *et al*²⁴ described cytoskeletal dysgenesis in myocytes. So far, the ethanol induced changes seen in actin cytoskeleton have been related to acetaldehyde, the main metabolite of ethanol oxidation, as studies carried out in vitro by Xu *et al*¹⁸ reported that G actin forms stable covalent adducts with acetaldehyde.

Given the significance of the cytoskeleton for enterocyte function at birth, and its importance to postnatal nutrition in mammals, the aim of this study is to determine in the newborn rat which cytoskeletal proteins are affected by longterm ethanol exposure in utero and to what extent.

Methods

Animals and treatments

The experiments were performed on the offspring of female Wistar rats from our own

Departament de Bioquímica i Fisiologia, Facultat de Biologia, Universitat de Barcelona, Barcelona, Spain
J F Montes
G Estrada
M D López-Tejero
J García-Valero

Correspondence to:
Dr José García-Valero,
Unitat de Biologia Cel·lular,
Departament de Bioquímica
i Fisiologia, Facultat
de Biologia, Avda Diagonal,
645, 08028 Barcelona,
Spain.

Accepted for publication
13 December 1995

Keywords: actin, cytokeratin, fetal alcohol syndrome, trans golgi network, tubulin, ubiquitin.

Figure 1: Actin immunolocalisation in the microvillus domain of control (A, $\times 58\,000$) and prenatally ethanol exposed enterocytes (B, $\times 52\,000$). Tubulin immunolocalisation in the cytoplasm of control (C, $\times 52\,000$) and prenatally ethanol exposed enterocytes (D, $\times 54\,000$). Bars = $5\,\mu\text{m}$.

colony, maintained at 25°C , 75% of relative humidity, and 12 hour light-dark cycles. The mothers, weighing 150–160 g, were placed in plastic cages distributed in groups of three rats per cage and were fed ad libitum with standard rodent chow (Panlab, Barcelona, Spain). Four weeks before pregnancy, the animals were divided into two experimental groups and subjected to different nutritional treatments²⁵: ethanol treated group received chow and 10%, 15%, 20%, and 25% (w/v) of ethanol in the drinking fluid in successive weeks before fecundation (ethanol adaptation period). After mating with untreated males, pregnant rats were kept on chow and 25% ethanol in drinking fluid until delivery (gestational period). The control group were handled in the same way as the ethanol treated group, fed ad libitum with the normal standard diet and water as drinking fluid. All rats were allowed to deliver spontaneously. Immediately after birth, pups were removed from their mothers and kept in a thermostatically controlled humidity chamber at 37°C until death.

Specimen preparation and immunocytochemistry
Three neonates from different litters in each experimental group were killed by decapitation

and the total small intestine (from the pylorus to the ileocaecal junction) was perfused gently, in situ, with cold NaCl 0.9% solution. The proximal small intestine (duodenum and proximal jejunum) were cut into small blocks of tissue and were fixed in 4% (w/v) paraformaldehyde and 0.1% (v/v) glutaraldehyde in 0.1 M phosphate buffered saline (PBS), pH 7.4, for two hours at 4°C . The samples were then cryo-protected by infusion with 2.1 M sucrose in PBS, mounted, and frozen in liquid nitrogen. Ultrathin cryosections were obtained using a cryo system (Reichert FC-40) and transferred to grids in a drop of 2.3 M sucrose in PBS.

Before labelling, sections were rinsed twice with 0.1 M glycine in PBS (gly PBS) for 10 minutes and incubated with 2% ovalbumin in PBS for 20 minutes. For single immunolabelling,²⁶ the grids were incubated during 30 minutes with monoclonal antibodies (mAbs) against α -tubulin (Amersham, Arlington Heights, IL), cytokeratin (polypeptides 1, 5, 6, 7, and 8 of the basic and 10, 11, and 18 of the acidic cytokeratins; ICN Biomedicals, Costa Mesa, CA), actin (Amersham), and with a polyclonal anti- β 1,4-galactosyl transferase serum (gift of Dr S Vilaró, Facultat de Biologia, Barcelona). After washing three times with gly PBS for 15 minutes, bound mAbs were visualised with 10 nm colloidal gold conjugated goat antimouse Ig (Amersham) and bound anti- β 1,4-galactosyl transferase serum was detected with 10 nm protein A gold (pAg) (Sigma Chemical, St Louis, MO). For double immunolabelling,²⁷ sections were first labelled for anti- β 1,4-galactosyl transferase serum with 10 nm pAg, then stabilised with 1% glutaraldehyde in PBS for five minutes, and incubated with the mAbs against actin or cytokeratin followed by unlabelled rabbit antimouse Ig (Amersham) and detected with 16 nm pAg (Sigma). The ultrathin cryosections were rinsed with PBS and double-distilled water before staining with uranyl acetate and embedding in methyl cellulose.²⁸ Observations were carried out with a Hitachi H-600 AB transmission electron microscope.

Quantitative evaluation

Label density was estimated as the number of gold particles either per area of compartment

Label density expressed as number of gold particles (gp) per unit length or area for α -tubulin, actin, and cytokeratin of several cell compartment

	Control group	Ethanol treated group
Tubulin		
Cytoplasm (gp/ μm^2)	7.63 (1.10)	11.46 (0.74)
Actin		
Microvilli (gp/ μm^2)	95.22 (10.24)	121.43 (7.30)
Lateral interdigitations (gp/ μm)	2.48 (0.24)	4.76 (0.83)**
Vesicles (gp/ μm)	1.75 (0.23)	12.13 (1.38)**
Cytokeratin		
Bundles (gp/ μm^2)	233.33 (27.60)	484.48 (36.20)**
Vesicles (gp/ μm)	0.79 (0.19)	10.19 (0.99)***

Values are mean (SEM) from at least 30 measurements performed on three animals from each experimental group. The significance of differences between control and ethanol treated groups was evaluated using the Student's *t* test. ** $p < 0.01$ and *** $p < 0.001$ comparing values in both experimental groups.

Figure 2: Actin immunolocalisation in control (A and C) and prenatally ethanol exposed enterocytes (B and D). Vesicular elements from apical and lateral enterocyte cytoplasm showed no labelling in controls (A, $\times 55\,000$), whereas homologous areas in the ethanol exposed group were strongly labelled (B, $\times 53\,000$). The lateral interdigitations between adjacent enterocytes were little developed and poorly labelled in controls (C, $\times 57\,000$), whereas in the ethanol exposed group they were highly developed and labelled (D, $\times 62\,000$). Bars = $5\,\mu\text{m}$.

or per unit of membrane length.²⁹ Both parameters, area and length, were estimated by stereological methods³⁰ and significance of mean differences was tested by Student's *t* test.

Preparation of cytoskeletal-enriched fractions from small intestine

Enriched fractions of intestinal cell cytoskeleton were obtained according to Quaroni *et al.*³¹ Briefly, the samples were homogenised in a glass dounce homogeniser (Anorsa, Spain) with 20 mM TRIS-HCl, pH 7.4, containing a mixture of protease inhibitors. Extraction of cytoskeletal proteins was carried out by centrifugations at 11 000–50 000 *g* in non-ionic detergents and buffers of both low and high salt concentration containing a protease inhibitor mixture. The high salt extracted cytoskeletal pellets were washed with PBS containing 0.2 mM PMSF and 0.1 mM DTT, and stored at -35°C in PBS until use.

SDS-PAGE and western blotting

SDS-PAGE was performed in a Protean I slab

cell (Bio-Rad Laboratories, Richmond, CA) in 5–12% acrylamide gels under reducing conditions. Polypeptides were stained in the gels using Coomassie Blue R (Bio-Rad). Apparent molecular masses of polypeptides were determined from their relative mobilities compared with a standard of molecular mass protein (thyroglobulin, 330 kDa; bovine serum albumin, 67 kDa; catalase, 60 kDa; lactate dehydrogenase, 36 kDa, and ferritin 18.5 kDa).

After electrophoresis, polypeptides from SDS gels were equilibrated in 50 mM TRIS-HCl, pH 7.4, in 20% glycerol for one hour³¹ and then transferred to nitrocellulose membranes³² in carbonate blot buffer (10 mM NaHCO_3 , 3 mM Na_2CO_3 , pH, 9.9, in 20% MeOH). Transfer was achieved at 20 V overnight at 4°C . Membranes were blocked with 3% ovalbumin in 10 mM PBS for 30 minutes and then incubated either with the same mAb against cytokeratin used in the immunocytochemical techniques (diluted 1:20 000 in PBS with 1% ovalbumin) or with rabbit antiubiquitin serum (Sigma, diluted 1:300) overnight at 4°C . After three washes in 10 mM PBS 0.2% TWEEN 20 (Sigma) the membranes were incubated for two hours at room temperature with peroxidase conjugated rabbit antimouse (Ig or peroxidase conjugated swine antirabbit Ig (Dako, Glostrup, Denmark) depending on the primary antibody. The membranes were developed in a substrate solution of 3,3'-diaminobenzidine tetrahydrochloride (DAB, Sigma) as chromogen and recorded on Technical Pan film (Kodak, Hemel Hempstead, UK).

Results

Cytoskeletal immunolocalisation was performed in duodenum and proximal jejunum of both control and prenatally ethanol exposed newborn rats. Within each experimental group, absorptive cells from the duodenum were structurally similar to those of the proximal jejunum, and had the same organisation pattern as those of adults. However, prenatally ethanol exposed enterocytes showed greater development of endomembranes around the nucleus than controls. These membranes, which appeared in a vesicular form and associated with saccules of Golgi complex (see Figs 2, 3, and 4), were identified as trans golgi or trans golgi network (TGN) by specific labelling with anti- β 1,4-galactosyl transferase (see Fig 4).

Tubulin was similarly distributed throughout the cytoplasm of control and prenatally ethanol exposed neonates, mainly in microtubular structures. No atypical intracellular localisation was seen such as label association with microvilli or TGN (Fig 1 (C) and (D)). The label density of the tubulin associated with cytoplasm showed no significant differences between control and prenatally ethanol exposed neonates (Table).

Actin distribution in the apical domain of the enterocyte was similar in control and prenatally ethanol exposed neonates (Fig 1 (A) and (B)), and actin label density in the

Figure 3: Cytoke- ratin immunolocalisation in control (A and C) and prenatally ethanol exposed enterocytes (B and D). Golgi related vesicles were devoid of label in controls (A, $\times 63\ 000$), whereas in the ethanol exposed group they showed strong labelling associated with the inner face of vesicular membranes (B, $\times 45\ 000$). Cytoke- ratin bundles showed little label in controls (C, $\times 21\ 000$), whereas in the ethanol exposed group they showed abundant label (D, $\times 53\ 000$). Bars (A, B, D) = $5\ \mu\text{m}$; bar (C) = $2\ \mu\text{m}$.

microvilli showed no significant differences between experimental groups (Table). Likewise, the cytoplasmic bundles of actin filaments showed no quantitative or distribution differences between experimental groups (data not shown). However, the interdigitations in the lateral domain of plasma membrane were more developed in prenatally ethanol exposed neonates than controls (Fig 2 (C) and (D)), and the actin label density associated with these structures was significantly increased in ethanol exposed enterocytes (Table). Regarding the vesicular compartment, abundant actin label density was associated with the internal face of vesicular membranes in prenatally ethanol exposed enterocytes, whereas control vesicles were devoid of label, showing a label density similar to background (Fig 2 (A) and (B); Table).

Cytoke- ratin localisation at the terminal web and apical cytoplasm showed no significant differences between control and prenatally ethanol exposed enterocytes (Fig 3 (C) and (D)). However, the cytoke- ratin bundles from medial and basal cytoplasm showed a significant label density increase in prenatally ethanol exposed enterocytes, compared with controls

(Table). Furthermore, like actin, cytoke- ratin was associated with the vesicular compartment of prenatally ethanol exposed enterocytes, and absent in control vesicles (Fig 3 (A) and (B)).

Characterisation of this vesicular compartment with anti- β 1,4-galactosyl transferase (Fig 4 (A)) showed that both actin (Fig 4 (B)) and cytoke- ratin (Fig 4 (C)), but not tubulin, were associated with the trans golgi or TGN compartments, or both, after prenatal ethanol exposure.

The protein pattern of intestinal cytoskeletal enriched fractions was similar in control and prenatally ethanol exposed neonates, as shown by SDS-PAGE analysis. However, a significant increase in polypeptides of molecular mass ranging from 39 to 54 kDa was seen in prenatally ethanol exposed neonates (Fig 5 (A)), compared with controls. This result supports the data of immunogold quantifications (Table) as the increased protein range coincides with the molecular mass of most intestinal cytoskeletal proteins.

Western blotting of the intestinal cytoskeletal enriched fractions and subsequent analysis of cytoke- ratins with the same mAb as used for immunolocalisation, showed a similar cytoskeletal pattern for the two experimental groups within the range of 45–53 kDa (Fig 5 (B)). However, an additional band of an apparent molecular mass of 28–29 kDa was detected in the prenatally ethanol exposed neonates. It was noteworthy that none of the cytoke- ratins characterised so far in published reports have a molecular mass similar to 28–29 kDa.

The increased abundance of ubiquitin in prenatally ethanol exposed neonates, compared with controls, was shown by western blotting of the intestinal cytoskeletal enriched fractions (Fig 6). This increase corresponds basically to the free ubiquitin pool.

Discussion

This study shows the effects of ethanol on the major cytoskeletal proteins of the gut epithelium from rat neonates, which was exposed to higher concentrations in the gastrointestinal lumen than in other fetal compartments.⁴ Ethanol crosses the placental barrier and is distributed and accumulated in a concentration gradient.^{4 33} However, reports of the passage of acetaldehyde, a product of ethanol oxidation, across the placental barrier are contradictory.^{34–37} Although placenta and fetal liver contain little alcohol dehydrogenase,^{4 36} there is evidence during longterm alcoholism of other pathways of ethanol oxidation both from maternal and from fetal tissues,^{38 39} which could interact with the cytoskeleton. These derivatives of ethanol, such as acetaldehyde and hydroxyethyl radicals, are probably produced either by peroxisomal catalase or by the microsomal ethanol inducible cytochrome P450.⁴⁰ Furthermore, Lange⁴¹ reported the formation of ethyl esters by non-oxidative reactions associated with glutathione S-transferase.

Increased concentrations of actin and cytoke- ratins, but not tubulin, were first suggested

Figure 4: Positive control for the immunolocalisation of β 1,4-galactosyl transferase, a marker for trans golgi and TGN (A, $\times 58\,000$). Double immunostaining for β 1,4-galactosyl transferase (10 nm gold) and either actin (15 nm gold) (B, $\times 46\,000$) or cytokeratin (15 nm gold) (C, $\times 56\,000$). The colocalisation shows the association of both actin and cytokeratin epitopes with the TGN membranes. Section C illustrates both the dilatation of cytokeratin containing TGN vesicles and cytokeratin association with desmosomes (arrowheads). Bars = 5 μ m.

by the labelling densities obtained in immunolocalisations of prenatally ethanol exposed enterocytes, compared with controls. Furthermore, this increase was confirmed by the increased protein abundance in the range of 39 to 54 kDa found in SDS-PAGE analysis, as nearly all cytoskeletal proteins fall in this range. This cytoskeletal increase could be produced either by protein overexpression or by inhibition of protein degradation. With regard to the first possibility, Zern *et al*¹⁰ showed a considerable increase in albumin synthesis in hepatocytes of rats fed for one year with a diet in which 36% of energy came from ethanol, compared with control rats. This protein increase was caused by an increase in the steady state level of active albumin mRNA in the adult ethanol exposed liver. With regard to the small intestine of neonates, a 10-fold to 15-fold increase in the values of lactase and intestinal alkaline phosphatase mRNAs was found after prenatal ethanol exposure.⁴² However, this increase was not correlated with a similar increase in the protein mass.

The possibility of an ethanol induced inhibition of protein degradation has been studied in

Figure 5: (A) SDS-PAGE analysis of intestinal cytoskeleton enriched fractions (40 μ g/lane). Lane 1 corresponds to standard mass proteins. Lane 2 shows basal protein expression in control neonates. Lane 3 shows the overexpression in polypeptides ranging from 39–54 kDa induced by prenatal ethanol exposure. (B) Western blot analysis of specific cytokeratins performed from the previous SDS-PAGE as described in methods. Lane 1 shows control neonates; lane 2, prenatally ethanol exposed neonates. The blot confirms the overexpression of cytokeratins after prenatal ethanol exposure and shows a unique peptide of 28–29 kDa (arrowhead), which is a specific feature for the ethanol group.

the liver of adult alcohol fed rats.¹⁴ The volume densities of autophagosomes and autolysosomes in hepatocytes was lower in ethanol treated rats than controls. Moreover, the rate of proteolysis was decreased by 30% in ethanol treated rats, suggesting that ethanol feeding inhibits proteolysis in the adult liver by preventing the sequestration of protein into lysosomes. Western blotting of intestinal cytoskeletal enriched fractions showed a significant increase in ubiquitin in prenatally ethanol exposed neonates. Most of the ubiquitin increase can be attributed to the effect that ethanol exposure induces an increased gene expression of stress proteins.¹²

Figure 6: Western blot analysis of free ubiquitin (Ubq) in intestinal cytoskeletal enriched fractions.

On the other hand, Corsi *et al*⁴³ showed that the cytoskeleton is covalently modified by ubiquitination. Furthermore, Akamatsu *et al*⁴⁴ stated that ubiquitination is an important process in the degradation of intermediate filament proteins in the cytoplasm. However, our results showed that the increase in the cytoke-
 ratin ubiquitination was not correlated with an increase in the cytoke-
 ratin degradation. Moreover, the results obtained from the ethanol treated group – that is, the increase in cytoke-
 ratin amount and the presence of the cytoke-
 ratin related polypeptide of 28–29 kDa – suggest a defective degradation of the proteins. The increased reactivity that several cytoskele-
 tal proteins exhibit for ethanol derivatives, especially aldehydes, could block the specific binding of ubiquitin, as both bind to lysine residues.^{18 45 46}

Furthermore, we observed that enterocytes exposed to ethanol in utero showed deficiencies in cell-cell and cell-basal lamina adhesion, with occasional detachment of the epithelial sheet (data not shown). This was in agreement with data of neural cell experiments in which addition of ethanol to the culture induced cell detachment.²⁰ Ethanol in the culture medium interacted with the L1 and N-CAM molecules of neurones resulting in inhibition of cell adhesion. In our study, the increased cyto-
 skeletal components are actin and cytoke-
 ratins, which happen to be the elements that interact with the cell adhesion molecules of plasma membrane. Tubulin, on the other hand, is not increased after prenatal ethanol exposure, but it happens to be a cytoskeletal element unrelated to cell adhesion. We hypothesise that the presence of a high concentration of ethanol (an average of 30 mM) in the fetal gastrointestinal lumen⁴ would first result in covalent modifi-
 cations of the surface adhesion molecules of enterocytes. Secondly, in response to loss of adhesiveness, feed back mechanisms would increase the expression of cell adhesion mole-
 cules, which might cause the overexpression and rearrangement of cytoskeletal elements.

The atypical localisation of actin and cyto-
 keratin in the trans golgi or the TGN, or both, can only be explained by the molecular modifi-
 cations caused by ethanol or its derivatives, or both, in both the cytoskeletal elements and the membrane of these compartments. Ethanol is a substrate for phospholipase D,¹⁶ an enzyme that catalyses the hydrolysis of phospholipids, such as phosphatidylcholine, to phosphatidic acids. However, this enzyme also utilises ethanol and several short chain alcohols as substrates, resulting in the formation of specific phosphatidylalcohols, such as phosphatidylethanol. Phosphatidylethanol induces an increase in the fluidity of artificial and natural bilayers, considerably modifying the physicochemical properties of membranes.¹⁵ Moreover, the hydrophobic head group of phosphatidylethanol might change the mem-
 brane architecture by disrupting the ionic bond lattice extending over the surface of the mem-
 brane.⁴⁷ These molecular modifications of cellular membranes might facilitate the non-
 specific entrance of altered proteins, actin, and

cytoke-
 ratins, in the trans golgi or TGN, or both.

In conclusion, our findings suggest that ethanol in utero exerts severe changes in the cytoskeleton of the developing intestinal epithelium. Actin and cytoke-
 ratin are two affected components showing increased expression and atypical intracellular localisa-
 tion. In contrast, tubulin is not affected. Therefore, the effects of ethanol in utero are specific for the cytoskeletal elements related with the plasma membrane and thus, affecting cell adhesion. These findings may contribute to understanding the cellular and molecular basis of intestinal dysfunctions found in the offspring of alcoholic mothers, causing pre and postnatal growth retardation.

The authors are grateful to Dr Senén Vilaró (Unitat de Biologia Cel·lular, Facultat de Biologia) for the serum against β 1,4-galactosyl transferase, to the staff of Serveis Científic-Tècnics (Universitat de Barcelona), and to Mr Robin Rycroft and Ms Mireia Martín for linguistic advice.

- Buts JP, Sokal EM, Van Hoof F. Prenatal exposure to ethanol in rats, effects on postnatal maturation of the small intestine and liver. *Pediatr Res* 1992; 32: 574–9.
- López-Tejero MD, Arilla E, Colás B, Llobera M, Herrera E. Low intestinal lactase activity in offspring from ethanol-treated mothers. *Biol Neonate* 1989; 55: 204–13.
- Camps L, Kedinger M, Simon-Assmann P, López-Tejero MD. Effect of prenatal exposure to ethanol on intestinal development of rat fetuses. *J Pediatr Gastroenterol Nutr* (in press).
- Través C, Camps L, López-Tejero MD. Liver alcohol dehydrogenase activity and ethanol levels during chronic ethanol intake in pregnant rats and their offspring. *Pharmacol Biochem Behav* 1995; 52: 93–9.
- Gottlieb TA, Ivanov IE, Adesnik M, Sabatini DD. Actin microfilaments play a critical role in endocytosis at the apical but not the basolateral surface of polarized epithelial cells. *J Cell Biol* 1993; 120: 695–710.
- Peterson MD, Bement WM, Mooseker MS. An in vitro model for the analysis of intestinal brush border assembly: II changes in expression and localization of brush border assembly in Caco-2BBE cells. *J Cell Sci* 1993; 105: 461–72.
- Mays RW, Beck KA, Nelson WJ. Organization and function of the cytoskeleton in polarized epithelial cells: a component of the protein sorting machinery. *Curr Opin Cell Biol* 1994; 6: 16–24.
- Reddy YS, Beesley RC. Effects of acute and chronic ethanol on cardiac contractile protein ATPase activity of Syrian hamsters. *Biochem Med Metab Biol* 1990; 44: 259–65.
- Guasch R, Renau-Piqueras J, Guerri C. Chronic ethanol consumption induces accumulation of proteins in the liver Golgi apparatus and decreases galactosyltransferase activity. *Alcohol Clin Exp Res* 1992; 16: 942–8.
- Zern MA, Chakraborty PR, Ruiz-Opazo N, Yap SH, Shafritz DA. Development and use of a rat albumin cDNA clone to evaluate the effect of chronic ethanol administration on hepatic protein synthesis. *Hepatology* 1983; 3: 317–22.
- Mochly-Rosen D, Chang F-H, Cheevers L, Kim M, Diamond I, Gordon AS. Chronic ethanol causes heterologous desensitization of receptors by reducing messenger RNA. *Nature* 1988; 333: 848–50.
- Miles MF, Diaz JE, DeGuzman V. Ethanol-responsive gene expression in neural cell cultures. *Biochem Biophys Acta* 1992; 1138: 268–74.
- Hu G, Querimit LA, Downing LA, Charness E. Ethanol differentially increases β -adrenergic and muscarinic acetylcholine receptor gene expression in NG108-15 cells. *J Biol Chem* 1993; 268: 23441–7.
- Pösö AR, Hirsimäki P. Inhibition of proteolysis in the liver by chronic ethanol feeding. *Biochem J* 1991; 273: 149–52.
- Omodeo-Salé F, Lindi C, Palestini P, Masserini M. Role of phosphatidylethanol in membranes. Effects on membrane fluidity, tolerance to ethanol, and activity of membrane-bound enzymes. *Biochemistry* 1991; 30: 2477–82.
- Hoek JB, Thomas AP, Rooney TA, Higashi K, Rubin E. Ethanol and signal transduction in the liver. *FASEB J* 1992; 6: 2386–96.
- Doyle KM, Bird DA, Al-Salihi S, Hallaq Y, Cluette-Brown J, Gross KA, *et al*. Fatty acid ethyl esters are present in human serum after ethanol ingestion. *J Lipid Res* 1994; 35: 428–37.
- Xu DS, Jennett RB, Smith SL, Sorrell MF, Tuma DJ. Covalent interactions of acetaldehyde with the actin/microfilament system. *Alcohol Alcoholism* 1989; 24: 281–9.
- Niemelä O, Juvonen T, Parkkila S. Immunohistochemical demonstration of acetaldehyde-modified epitopes in human liver after alcohol consumption. *J Clin Invest* 1991; 87: 1367–74.
- Charness ME, Safrans RM, Perides G. Ethanol inhibits neural cell-cell adhesion. *J Biol Chem* 1994; 269: 9304–9.

- 21 Sáez R, Bursal M, Renau-Piqueras J, Marqués A, Guerri C. Evolution of several cytoskeletal proteins of astrocytes in primary culture: effect of prenatal alcohol exposure. *Neurochem Res* 1991; 16: 737-47.
- 22 Barbatis C, Morton J, Woods JC, Burns J, Bradley J, McGee J O'D. Disorganisation of intermediate filament structure in alcoholic and other liver diseases. *Gut* 1986; 27: 765-70.
- 23 Pinazo-Durán MD, Renau-Piqueras J, Guerri C. Developmental changes in the optic nerve related to ethanol consumption in pregnant rats: analysis of the ethanol-exposed optic nerve. *Teratology* 1993; 48: 305-22.
- 24 Adickes ED, Mollner TJ. Ethanol-induced cytoskeletal dysgenesis with dietary protein manipulations. *Alcohol Alcoholism* 1986; 21: 347-55.
- 25 Testar X, López-Tejero MD, Llobera M, Herrera E. Ethanol administration in the drinking fluid to pregnant rats as a model for the fetal alcohol syndrome. *Pharmacol Biochem Behav* 1986; 24: 625-30.
- 26 Griffiths G, Simons K, Warren G, Tokuyasu KT. Immunoelectron microscopy using thin, frozen sections: application to studies of the intracellular transport of Semliki Forest virus spike glycoproteins. *Method Enzymol* 1983; 96: 435-50.
- 27 Geuze HJ, Slot JW, van der Ley PA, Scheffer RCT, Griffith JM. Use of colloidal gold particles in double-labelling immunoelectron microscopy of ultrathin frozen tissue sections. *J Cell Biol* 1981; 89: 653-65.
- 28 Tokuyasu KT. Immunocytochemistry on ultrathin sections. *Histochem J* 1980; 12: 381-403.
- 29 Griffiths G, Hoppeler H. Quantitation in immunocytochemistry: correlation of immunogold labeling to absolute number of membrane antigens. *J Histochem Cytochem* 1986; 34: 1389-98.
- 30 Weibel ER. Stereological methods. Vol I. *Practical methods for biological morphometry*. New York: Academic Press, 1979.
- 31 Quaroni A, Calnek D, Quaroni E, Chandler JS. Keratin expression in rat intestinal crypt and villus cells. *J Biol Chem* 1991; 266: 11923-31.
- 32 Towin H, Staehelin T, Gordon J. Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proc Natl Acad Sci USA* 1979; 76: 4350-4.
- 33 Guerri C, Sanchis R. Acetaldehyde and alcohol levels in pregnant rats and their fetuses. *Alcohol* 1985; 2: 267-70.
- 34 Kesäniemi YA, Sippel HW. Placental foetal metabolism and acetaldehyde in rat. I. Content of ethanol acetaldehyde in placenta and foetus of the pregnant rat during ethanol intoxication. *Acta Pharmacol Toxicol* 1975; 47: 43-8.
- 35 Zorzano A, Herrera E. Disposition of ethanol and acetaldehyde in late pregnant rats and their fetuses. *Pediatr Res* 1989; 25: 102-6.
- 36 Sanchis R, Guerri C. Alcohol-metabolizing enzymes in placenta and fetal liver: effects of chronic ethanol intake. *Alcohol Clin Exp Res* 1986; 10: 39-44.
- 37 Gordon BHJ, Baraona E, Miyakawa H, Finkelman F, Lieber CS. Exaggerated acetaldehyde response after ethanol administration during pregnancy and lactation in rats. *Alcohol Clin Exp Res* 1985; 9: 17-22.
- 38 Cambon-Gros G, López-Tejero MD, Fernández Y, Vilarem G, Mitjavila S. Increase in the CCl₄ radical induced effects on the rat liver microsomes after chronic ethanol intake during gestation. In: Nordmann R, Ribière C, Rouach H, eds. *Advances in the Biosciences*. Vol 71. Great Britain: Pergamon Press, 1988: 159-63.
- 39 Terada N, Nakai T, Yamaguchi M, Hattai A, Arizono K, Ariyoshi T. Effects of maternal ethanol intake during pregnancy on fetal and maternal liver enzymes systems in Wistar rats. *J Pharmacobiodyn* 1982; 5: 49-54.
- 40 Lieber CS. Biochemical mechanisms of alcohol-induced hepatic injury. *Alcohol Alcoholism* 1991; suppl 1: 283-90.
- 41 Lange LL. Mechanism of fatty acid ethyl ester formation and biological significance. *Ann NY Acad Sci* 1991; 625: 802-5.
- 42 Estrada G, Krasinski SD, Montgomery RK, Grand RJ, García-Valero J, López-Tejero MD. Quantitative analysis of lactase-phlorizin hydrolase expression in the absorptive enterocytes of newborn rat small intestine. *J Cell Physiol* (in press).
- 43 Corsi D, Galluzzi L, Crinelli R, Magnani M. Ubiquitin is conjugated to the cytoskeletal protein α -spectrin in mature erythrocytes. *J Biol Chem* 1995; 270: 8928-35.
- 44 Akamatsu M, Hori S, Tsutsumi Y, Yoshiyuki R, Ohkido M. Ubiquitinated cytokeratin inclusions in lichen amyloidosis: an immunohistochemical analysis. *Pathol Int* 1995; 45: 116-22.
- 45 Gregori L, Marriott D, West CM, Chau V. Specific recognition of calmodulin from Dictyostelium discoideum by the ATP, ubiquitin-dependent degradative pathway. *J Biol Chem* 1985; 260: 5232-5.
- 46 Rechsteiner M. Ubiquitin-mediated pathways for intracellular proteolysis. *Ann Rev Cell Biol* 1987; 3: 1-30.
- 47 Boggs JM. Lipid intermolecular hydrogen bonding: influence on structural organization and membrane function. *Biochem Biophys Acta* 1987; 906: 353-7.