Rectal corticosteroids versus alternative treatments in ulcerative colitis: a meta-analysis

J K Marshall, E J Irvine

Abstract

Background—Clear strategies to optimise the use of corticosteroids in ulcerative colitis are lacking.

Aim—A meta-analysis was undertaken to examine critically the role of rectal corticosteroids in the management of active distal ulcerative colitis.

Methods—All reported randomised controlled trials were retrieved by searching the Medline and EMBASE databases and the bibliographies of relevant studies. Trials which met inclusion criteria were assessed for scientific rigour. Data were extracted by two independent observers according to predetermined criteria.

Results—Of 83 trials retrieved, 33 met inclusion criteria. Pooled odds ratios (POR) showed conventional rectal corticosteroids and rectal budesonide to be clearly superior to placebo. In seven trials, rectal 5-aminosalicylic acid (5-ASA) was significantly better than conventional rectal corticosteroids for inducing remission of symptoms, endoscopy, and histology with POR of 2.42 (95% confidence interval (CI) 1.72-3.41), 1.89 (95% CI 1.29-2.76), and 2.03 (95% CI 1.28-3.20), respectively. Rectal budesonide was of comparable efficacy to conventional corticosteroids but produced less endogenous cortisol suppression. Side effects, although inconsistently reported, were generally minor. A cost comparison of rectal preparations showed 5-ASA to be less expensive than corticosteroids.

Conclusions—Rectal 5-ASA is superior to rectal corticosteroids in the management of distal ulcerative colitis.

(Gut 1997; 40: 775-781)

Keywords: ulcerative colitis, corticosteroids, therapy, topical administration, enema, suppository.

Oral corticosteroids have been a well-accepted treatment for active ulcerative colitis since Truelove *et al* reported the efficacy of oral hydrocortisone over 40 years ago.¹ However, their long term use may be limited and patient compliance diminished by potential adverse effects. Administration of a corticosteroid liquid enema was first suggested to be efficacious in distal ulcerative colitis in 1956.² The proven efficacy of direct drug delivery to the site of inflammation has since led to widespread acceptance of rectal corticosteroid therapy.³

The ability of a rectal preparation to achieve a proximal distribution is determined by the type of vehicle. Liquid enemas can deliver

medication consistently to the splenic flexure,⁴⁻⁹ and a larger volume seems to allow more proximal delivery.¹⁰⁻¹¹ Rectal foam disseminates medication to the rectum and distal descending colon,¹²⁻¹⁶ whereas suppositories coat only the rectum.¹⁷⁻¹⁸

Although studies of rectally administered corticosteroids have reported fewer systemic adverse effects than with oral preparations, plasma concentrations of prednisolone were similar after administration of identical oral or rectal doses.¹⁹⁻²⁰ Suppression of the hypothalamic-pituitary-adrenal axis in association with rectal therapy has also been shown.²¹⁻²⁶

Newer topically active corticosteroids such as tixocortol, beclomethasone, prednisolone metasulphabenzoate, and budesonide, with restricted absorption or rapid hepatic metabolism have been developed to reduce the adverse effects associated with conventional corticosteroids.²⁷⁻²⁸

To examine critically the role of rectal corticosteroids in the treatment of active distal ulcerative colitis, we performed a meta-analysis of all reported randomised controlled trials.

Methods

Relevant clinical trials were identified by searching the Medline database from 1966 to 1996 and the EMBASE database from 1985 to 1996, using the MeSH terms "inflammatory bowel disease", "therapy", and "topical administration", "enema", or "suppository". Bibliographies of all relevant studies and recent review articles were scanned to identify further citations. Each paper was assessed by two independent observers (JKM, EJI) according to predetermined inclusion criteria. Studies were accepted if patients had active ulcerative colitis with a documented disease margin distal to the splenic flexure on radiographic studies or less than 60 cm from the anal verge at flexible sigmoidoscopy or colonoscopy. We required that patients had been randomly assigned to two or more treatment groups, with rectal corticosteroids in at least one treatment arm and a symptom score as one of the main outcome criteria. The minimum duration of therapy permitted was two weeks.

Trials which met the inclusion criteria were then evaluated quantitatively for scientific rigour using a 30 point scoring system.²⁹ Disagreements in scoring between the two observers were settled by consensus.

Data were extracted from each report using a predefined format. Recorded data points included the number of patients enrolled, number completing the study, sex, and disease distribution. The proportion of patients that improved or attained remission, or both, by

Division of Gastroenterology and Intestinal Disease Research Programme, McMaster University, Hamilton, Ontario, Canada
J K Marshall
E J Irvine

Correspondence to:
Dr E J Irvine,
Division of Gastroenterology,
Room 4W8, McMaster
University Medical Center,
1200 Main Street West,
Hamilton, Ontario
L8N 3Z5, Canada.

Accepted for publication
23 December 1996

symptomatic, endoscopic, and histological criteria were recorded from each trial using an intention to treat principle. The dose, frequency, duration, and formulation (enema, suppository or foam) of treatments were noted, as were any reported adverse effects of therapy.

We anticipated that the definitions of "improvement" and "remission" would vary considerably among the papers accepted. To overcome the problems of standardising endpoints, provided that an adequate definition of improvement or remission was offered, the authors' criteria for these outcomes were respected. All clinical symptom scores included stool frequency and rectal bleeding.

Studies were grouped for analysis according to the type of corticosteroid and control therapies. Corticosteroid doses were also converted to a hydrocortisone dose equivalent to permit testing for a dose response relation.³⁰ The statistical analysis was conducted using the method of DerSimonian and Laird.³¹ An odds ratio for each trial and a common odds ratio for each group with 95% confidence intervals (CI) were calculated according to Mantel-Haenszel. Unless otherwise stated, odds ratios below 1 favoured corticosteroid treatment, whereas odds ratios above 1 favoured the alternative treatment. Continuous data points were pooled and compared using a weighted mean difference.³² Homogeneity within groups of trials was confirmed using the Breslow-Day test.³³ Overall response rates for each drug were calculated by dividing the total number of patients reaching an endpoint by the total number of patients treated.

Results

ACCEPTANCE AND VALIDITY SCORING

In total, 83 relevant trials were identified by the search, of which 33 met our inclusion criteria. Reasons for excluding a trial included: lack of randomisation (36 trials), inclusion of patients with disease proximal to the splenic flexure (29 trials), lack of a predefined symptom score (nine trials), duplicate reporting of data (three trials), and inclusion of patients with Crohn's colitis (one trial).

Table I summarises the characteristics of the 33 trials accepted for analysis. The median validity score (out of 30) was 21 (range 9–26).

RESPONSE RATES

Table II lists the response rates for all treatments and placebo. Among patients receiving conventional rectal corticosteroids (hydrocortisone, prednisolone, or betamethasone), pooled improvement rates by symptomatic, endoscopic, and histological criteria were 77%, 66%, and 52%, whereas remission rates were 45%, 34%, and 29%, respectively. The pooled response rates for the topically active corticosteroids (budesonide, beclomethasone, or prednisolone metasulphobenzoate) were similar: 73% for symptoms, 69% for endoscopy, and 55% for histology, whereas 46%, 31%, and 23%, respectively, attained

remission. When corticosteroid doses were converted to their hydrocortisone equivalent,³⁰ no dose response relation was observed for either conventional or topical formulations. Similarly, no correlation was apparent between duration of treatment and response rate.

Aminosalicylates (4-ASA or 5-ASA), which were used most frequently as a comparative treatment, produced improvement in symptoms, endoscopy, and histology in, respectively, 81%, 75%, and 65% of patients. Remission with these endpoints was induced in 52%, 41%, and 32% of patients, respectively.

Across four trials, 34% of patients taking placebo improved symptomatically, and 38% improved endoscopically. Remission rates by symptomatic and endoscopic criteria were 9% and 17%.

RECTAL CORTICOSTEROIDS VERSUS PLACEBO

Two trials compared conventional rectal corticosteroids with placebo.^{34–35} The combined results clearly favoured corticosteroids, with a pooled odds ratios (POR) for symptomatic and endoscopic improvement of 0.21 (95% CI 0.07–0.71) and 0.27 (95% CI 0.10–0.77), respectively. The PORs for symptomatic and endoscopic remission were 0.07 (95% CI 0.02–0.29) and 0.34 (95% CI 0.10–1.20), respectively. Histological endpoints were not reported in these early trials.

One trial reported that 2.3 mg budesonide enemas were superior to placebo for improvement of symptoms, endoscopy, and histology,³⁶ whereas another found 2.0 mg or 8.0 mg daily superior to placebo in inducing combined symptomatic and endoscopic remission, with higher response rates at the larger dose.³⁷

RECTAL VERSUS ORAL CORTICOSTEROIDS

Rectal hydrocortisone 100 mg was compared with oral prednisolone 60 mg daily in one trial which showed oral treatment to be better for symptomatic improvement and remission.³⁹ A second trial compared low dose oral prednisolone (7.5 mg daily) with rectal prednisolone metasulphobenzoate 20 mg, and found rectal treatment to be more efficacious for inducing symptomatic improvement.³⁸ Because of substantial differences in oral dose, these results were not pooled.

RECTAL CORTICOSTEROIDS VERSUS RECTAL 5-ASA

Seven accepted trials compared rectal corticosteroids with rectal 5-ASA.^{40–46} The total daily dose of 5-ASA ranged from 1 to 4 g, whereas the hydrocortisone equivalent dose of corticosteroids ranged from 100 to 356 mg. One trial compared a hydrocortisone foam with a 5-ASA suppository,⁴³ whereas another compared hydrocortisone foam with 5-ASA foam.⁴⁵ All other trials compared liquid enema preparations of equal volume. Pooled odds ratios for symptomatic, endoscopic, and histological improvement among the trials reporting these data were 1.36 (95% CI 0.88–2.09), 1.06

TABLE I Characteristics of trials accepted for meta-analysis

Reference	Author (year)	Medication (dose and frequency)*	Duration (days)	Number of patients	Validity score (maximum=30)
A Rectal corticosteroids v placebo					
34	Lennard-Jones <i>et al</i> (1962)	Prednisolone (5 mg SUPP od)	21	39	16
35	Watkinson (1958)	Placebo Hydrocortisone (100 mg/100 ml od) Placebo	15	19	17
B Rectal budesonide v placebo					
36	Danielsson <i>et al</i> (1992)	Budesonide (2.3 mg/115 ml od)	28	41	23
37	Hanauer and Robinson (1995)†	Placebo Budesonide (0.5 mg or 2.0 mg or 8.0 mg od) Placebo	42	233	12
C Rectal v oral corticosteroids					
38	Hamilton <i>et al</i> (1984)	Prednisolone metasulphoenzoate (20 mg od) Prednisolone (7.5 mg ORAL od)	14	36	17
39	Lennard-Jones <i>et al</i> (1960)	Hydrocortisone (100 mg/150 ml) od Prednisone (variable dose ORAL od) Salazopyrin (4 g ORAL od)	21	60	15
D Rectal corticosteroids v rectal 5-ASA					
40	Bianchi Porro <i>et al</i> (1995)	Hydrocortisone (100 mg/60 ml od)	21	52	23
41	Campieri <i>et al</i> (1981)	5-ASA (1 g/100 ml od) Hydrocortisone (100 mg/100 ml od)	15	86	19
42	Danish 5-ASA Group (1987)	5-ASA (4 g/100 ml od) Prednisolone (25 mg/100 ml od)	14	123	25
43	Farup <i>et al</i> (1995)	5-ASA (1 g/100 ml od) Hydrocortisone (178 mg/60 ml FOAM bid)	28	79	23
44	Friedman <i>et al</i> (1986)	5-ASA (0.5 g SUPP bid) Hydrocortisone (100 mg/100 ml od)	21	18	24
45	Lee <i>et al</i> (1996)	5-ASA (4 g/100 ml od) Prednisolone (20 mg/30 ml FOAM od)	28	295	25
46	Mulder <i>et al</i> (1988)	5-ASA (2 g/120 ml FOAM od) Prednisolone (30 mg/40 ml od) 5-ASA (3 g/40 ml od)	28	29	24
E Rectal corticosteroids v rectal 4-ASA					
47	O'Donnell <i>et al</i> (1992)	Prednisolone (20 mg/50 ml od) 4-ASA (2 g/50 ml od)	42	45	24
48	Sharma <i>et al</i> (1992)	Prednisolone (20 mg/60 ml od) 4-ASA (2 g/60 ml od)	28	40	26
F Rectal corticosteroids v rectal budesonide					
24	Bianchi Porro <i>et al</i> (1994)	Methylprednisolone (20 mg/100 ml od) Budesonide (2 mg/100 ml od)	28	88	21
25	Danielsson <i>et al</i> (1987)	Prednisolone (31.25 mg/100 ml od) Budesonide (2 mg/100 ml od)	28	64	20
26	Danish Budesonide Study Group (1991)	Prednisolone (25 mg/100 ml od) Budesonide (1 or 2 or 4 mg/100 ml od)	14	139	19
49	Lofberg <i>et al</i> (1994)	Prednisolone (31.25 mg/125 ml od) Budesonide (2.3 mg/115 ml od)	56	100	24
50	Tarpila <i>et al</i> (1994)	Hydrocortisone (125 mg/125 ml od) Budesonide (2.3 mg/115 ml od)	28	72	20
G Rectal budesonide v rectal 5-ASA					
51	Lamers <i>et al</i> (1991)†	Budesonide (2 mg/100 ml od) 5-ASA (4 g/60 ml od)	28	62	13
52	Lemann <i>et al</i> (1995)	Budesonide (2 mg/115 ml od) 5-ASA (1 g/100 ml od)	28	92	22
H Other					
53	Cobden <i>et al</i> (1991)	Prednisolone metabenzoate (20 mg/100 ml bid) 5-ASA (0.8 g ORAL qid)	28	37	26
54	Grace <i>et al</i> (1987)	Prednisolone (20 mg/100 ml od)	56	70	18
55	Halpern <i>et al</i> (1991)‡	Sodium cromoglycate (600 mg/100 ml od) Beclomethasone dipropionate (0.5 mg/100 ml od)	28	40	22
56	Hanauer <i>et al</i> (1986)†	Betamethasone phosphate (5 mg/100 ml od) Tixocortol pivalate (250 mg od)	21	125	9
57	Lennard-Jones (1971)	Hydrocortisone (100 mg od) Betamethasone valerate (5 mg/100 ml od)	28	105	17
58	Mulder <i>et al</i> (1989)	Prednisolone (20 mg/100 ml od) Beclomethasone dipropionate (2 or 3 mg/40 ml od)	28	25	23
59	Mulder <i>et al</i> (1994)†	Prednisolone (30 mg/40 ml od) Beclomethasone dipropionate (3 mg/100 ml od) 5-ASA (1 g/100 ml od)	28	60	13
60	Riley <i>et al</i> (1989)	Beclomethasone and 5-ASA (3 mg/100 ml and 1 g/100 ml od) Prednisolone metasulphobenzoate (20 mg/100 ml od)	28	44	23
61	Ruddell <i>et al</i> (1980)	Sucralfate (4 g/100 ml od) Hydrocortisone (100 mg/60 ml bid)	14	30	16
62	van der Heide <i>et al</i> (1988)	Hydrocortisone (100 mg/5 ml FOAM bid) Beclomethasone dipropionate (1 mg/40 ml od)	28	18	24
63	van Outryve <i>et al</i> (1996)†	Prednisolone (30 mg/40 ml od) Ridogrel (300 mg/40 ml od) Prednisolone (30 mg/40 ml od)	28	40	17

*Medication in enema format unless otherwise stated. †Abstract only. ‡Total 32 patients with 40 treatment courses. SUPP=suppository; od=once daily; bid=twice daily; qid=four times daily.

(95% CI 0.61–1.85), and 2.27 (95% CI 1.22–4.27), with results clearly favouring 5-ASA only for histology (Figs 1 and 3). Using the stricter outcome of disease remission, 5-ASA was significantly better for all three criteria with PORs of 2.42 (95% CI 1.72–3.41) for symptoms, 1.89 (1.29–2.76) for endoscopy, and 2.03 (95% CI 1.28–3.20) for histology (Figs 2 and 3). When the two trials using foam preparations^{43 45} were excluded from the

analysis, recalculated PORs for remission endpoints still favoured 5-ASA significantly, despite wider confidence intervals as a result of the smaller sample size.

A single trial compared 5-ASA and beclomethasone enemas alone versus a combined 5-ASA/beclomethasone enema.⁵⁹ The combination surpassed monotherapy in inducing symptomatic or endoscopic improvement ($p<0.05$).

TABLE II Pooled response rates for rectal preparations (all trials)

Medication	Response rate across all trials (ratio, percentage)					
	Improvement			Remission		
	Symptomatic	Endoscopic	Histologic	Symptomatic	Endoscopic	Histological
Conventional corticosteroids						
Hydrocortisone	58/94 (62%)	61/109 (56%)	71/137 (52%)	50/119 (42%)	43/96 (45%)	14/51 (27%)
Prednisolone	272/344 (79%)	146/196 (74%)	73/144 (51%)	149/338 (44%)	127/387 (33%)	83/262 (32%)
Methylprednisolone	28/44 (59%)	20/44 (45%)	20/44 (45%)	16/44 (36%)	8/44 (18%)	6/44 (14%)
Betamethasone	61/71 (86%)	18/20 (90%)	16/20 (80%)	40/71 (56%)	6/20 (30%)	10/20 (50%)
Pooled	428/553 (77%)	245/369 (66%)	180/345 (52%)	255/572 (45%)	184/547 (34%)	113/377 (30%)
Topically active corticosteroids						
Prednisolone metasulphobenzoate	33/40 (83%)	15/22 (68%)	15/22 (68%)	15/22 (68%)	17/41 (41%)	10/22 (45%)
Beclomethasone	42/65 (65%)	44/65 (68%)	24/45 (53%)	9/20 (45%)	16/56 (29%)	5/20 (25%)
Budesonide	49/64 (77%)	163/237 (69%)	110/205 (54%)	53/137 (39%)	70/237 (30%)	26/137 (19%)
Pooled	124/169 (73%)	222/324 (69%)	149/272 (55%)	77/169 (46%)	103/334 (31%)	41/179 (23%)
Aminosalicylates						
5-ASA	230/282 (82%)	154/212 (73%)	93/141 (66%)	195/368 (53%)	140/384 (36%)	84/251 (33%)
4-ASA	37/47 (79%)	20/20 (100%)	29/47 (62%)	9/27 (33%)	18/20 (90%)	5/27 (19%)
Pooled	267/329 (81%)	174/232 (75%)	122/188 (65%)	204/395 (52%)	158/404 (39%)	89/278 (32%)
Placebo	18/53 (34%)	12/32 (38%)		3/32 (9%)	5/32 (17%)	

RECTAL CORTICOSTEROIDS VERSUS RECTAL 4-ASA

Rectal corticosteroids were compared with rectal 4-ASA in two trials.^{47, 48} Each compared 4-ASA 2 g with prednisolone 20 mg (hydrocortisone dose equivalent 80 mg). POR for symptomatic improvement was 3.88 (95% CI 1.29–11.64), favouring 4-ASA.

RECTAL CORTICOSTEROIDS VERSUS RECTAL BUDESONIDE

Five trials compared conventional rectal corticosteroids with rectal budesonide.^{24–26, 49, 50} The

budesonide dose ranged from 2.0 to 2.5 mg, whereas the corticosteroid dose ranged from 100 to 125 mg of hydrocortisone equivalent. Data from one trial could not be extracted adequately for meta-analysis.²⁶ One trial used a hydrocortisone foam.⁵⁰ All other medications were given as enemas.

The PORs for improvement by symptomatic, endoscopic, and histological criteria were 2.08 (95% CI 0.84–5.14), 1.40 (95% CI 0.87–2.25), and 1.23 (95% CI 0.80–1.91), respectively (Fig 4). PORs for symptomatic, endoscopic, and histological remission were 0.85 (95% CI 0.44–1.63), 1.14 (95% CI 0.69–1.88), and 0.68 (95% CI 0.28–1.67). All confidence intervals included 1.

Figure 1: Symptomatic improvement: rectal corticosteroids v rectal 5-aminosalicylic acid (5-ASA).

Figure 2: Symptomatic remission: rectal corticosteroids v rectal 5-aminosalicylic acid (5-ASA).

RECTAL BUDESONIDE VERSUS RECTAL 5-ASA

Two trials comparing rectal budesonide with rectal 5-ASA were evaluated.^{51, 52} Endoscopic improvement and remission data were reported in both trials, with a POR of 0.58 (95% CI 0.27–1.22) and 0.95 (95% CI 0.43–2.10), respectively, where an OR <1 favoured 5-ASA. In one of the trials 5-ASA exceeded budesonide for inducing symptomatic remission, with an OR of 0.41 (95% CI 0.18–0.94).⁵² The other reported similar symptomatic remission rates in both treatment arms, but the data provided did not permit calculation of an OR.⁵¹

Figure 3: Pooled odds ratios for all outcomes: rectal corticosteroids v rectal 5-aminosalicylic acid (5-ASA).

Figure 4: Pooled odds ratios for all outcomes: conventional rectal corticosteroids v rectal budesonide.

ADVERSE EFFECTS OF RECTAL CORTICOSTEROIDS

Adverse effects of treatment were inconsistently reported in the accepted trials. Nine of the 33 trials made no reference whatsoever to adverse effects, whereas a further 11 trials reported no drug related adverse effects in any treatment arm. Among the remaining 13 trials, seven dropouts for drug related effects were noted: four on 5-ASA, one on conventional corticosteroids, one on budesonide, and one

on 4-ASA. Other drug related adverse effects such as nausea, abdominal distension, fatigue, and perianal irritation were infrequent. Overall, 10 trials reported hypothalamic-pituitary-adrenal axis function before and after treatment. Three of these compared rectal budesonide with conventional rectal corticosteroids, noting mean cortisol concentrations after four weeks of treatment.^{25 49 50} Cortisol concentrations were consistently higher, indicating lesser suppression, in the budesonide group than in the group receiving conventional corticosteroids (Fig 5). The weighted mean difference between pooled treatment arms was 119.1 nmol/l (95% CI 70.3–167.9), confirming that this difference was statistically significant. Another trial reported similar data using an analog scale which could not be pooled.²⁶

COST COMPARISON

The costs for rectal steroid and 5-ASA enema and foam preparations available in Canada, in Canadian dollars, were obtained from the Chedoke-McMaster hospital pharmacy and are shown in Table III. Hydrocortisone 100 mg and 5-ASA 4 g enemas are comparable in cost, whereas 5-ASA 1 g and 2 g enemas cost considerably less. Budesonide enemas are marginally more expensive than hydrocortisone liquid enema, and hydrocortisone foam costs slightly less. Methylprednisolone enemas currently are not available in Canada.

Figure 5: Plasma cortisol concentrations and weighted mean difference (with 95% CI) after four weeks of treatment: conventional rectal corticosteroids v rectal budesonide.

Discussion

This study confirms that rectal corticosteroids are an effective treatment for active distal ulcerative colitis, with a therapeutic gain over placebo of approximately 30%, but suggests that rectal 5-ASA is significantly more efficacious for inducing disease remission. Placebo controlled data are becoming increasingly scarce in the recent literature, possibly because of the ethical concerns of treating patients with active ulcerative colitis with placebo. Two early placebo controlled trials^{34 63} confirmed the efficacy of conventional rectal corticosteroids for inducing improvement and remission by symptomatic and endoscopic criteria. Two recent trials^{36 37} also showed the topically acting corticosteroid budesonide to be superior to placebo using symptomatic, endoscopic, and histological endpoints. Our pooled placebo data demonstrated symptomatic and endoscopic improvement in 34% and 38%, and symptomatic and endoscopic remission in 9% and 17% of patients, respectively. These findings were similar to those of another overview which suggested that placebo benefited 30% of patients and produced remission in 10%.⁶⁴

Our meta-analysis suggests that rectal 5-ASA is as efficacious as rectal corticosteroids for improving disease and is better than rectal corticosteroids for inducing remission. Results were consistent, with similarly narrow confidence intervals, for symptomatic, endoscopic, or histological outcomes. The two trials which

Medication	Unit cost (Can \$; US\$; £) excluding dispensing fee	Cost of 14 day course (Can \$; US\$; £) excluding dispensing fee
Hydrocortisone enema 100 mg (Cortenema)	6.93; 4.89; 3.00	97.02; 68.50; 42.01
Hydrocortisone foam 80 mg (Cortifoam 14 dose cannister)	83.24; 58.77; 36.04	83.24; 58.77; 36.04
Betamethasone enema 5 mg (Betnesol)	8.68; 6.13; 3.76	121.52; 85.79; 52.62
Budesonide enema 2.3 mg (Entocort)	9.00; 6.35; 3.90	126.00; 88.96; 54.56
Tixocortol 250 mg (Rectovalone)	7.95; 5.61; 3.44	111.30; 78.58; 48.19
5-ASA enema 4 g (Salofalk)	7.08; 5.00; 3.07	99.12; 69.98; 42.92
5-ASA enema 2 g (Salofalk)	4.10; 2.89; 1.78	57.40; 40.52; 24.85
5-ASA enema 2 g (Quintasa)	4.35; 3.07; 1.88	60.90; 43.00; 26.37
5-ASA enema 1 g (Quintasa)	3.97; 2.80; 1.72	55.58; 39.24; 24.07

compared rectal 5-ASA with budesonide suggested that 5-ASA is at least as effective in producing disease improvement and remission. However, not all endpoints were noted.

Only two studies compared the role of rectal 4-ASA with corticosteroids, but supported a therapeutic gain in producing symptomatic improvement for 4-ASA.

Budesonide is a new topically active corticosteroid formulation with a high glucocorticoid receptor affinity, and significant first pass hepatic metabolism. The pooled results for accepted trials failed to demonstrate significant therapeutic benefit of either budesonide or conventional corticosteroids. However, budesonide caused significantly less endogenous cortisol suppression, based on pooled mean cortisol concentrations. Although these data are promising, no data have been reported regarding differences in steroid specific adverse effects, such as osteopenia or Cushingoid facies.

Other factors which may influence the efficacy of a treatment formulation include the proximal extent of the disease. Although our pooled data did not permit subgroup analysis, foam and suppositories did produce higher response rates in patients with more distal disease,⁴³ supporting the findings of radiological and radionuclide studies of a more distal distribution of medication.¹²⁻¹⁸ Patient preference for foam or suppository preparations also may augment compliance, and hence the effectiveness of therapy.⁶¹ Similarly, the volume of enema or foam preparations may influence treatment distribution and efficacy.¹⁰ Although most trials used equivalent volumes in both treatment arms, Lee *et al*⁴⁵ compared a 30 cc prednisolone foam with a 120 cc 5-ASA foam, a bias which could favour the efficacy of 5-ASA. When these data were excluded from the pooled results, 5-ASA remained superior to corticosteroids in inducing remission, although confidence intervals were wider.

An important feature which potentially could confound results of this meta-analysis was duration of treatment, which ranged from 14 to 56 days among the trials accepted. Although pooled results of trial endpoints did not demonstrate a clear relation, individual trials which reported interim endpoints at different time intervals observed higher endoscopic and histological remission rates with prolonged treatment.^{25 36 42 43 49-51} Although longer treatment may potentially increase adverse effects or diminish compliance, treatment requiring endoscopic or histological remission has been associated with a lower relapse rate.^{65 66}

Adverse effects for all rectal preparations were under-reported, but seemed comparable. Drug costs using a two week treatment regimen were lower for rectal aminosalicylate products than for most corticosteroid preparations. However, longer term studies may be necessary to evaluate further the median time to remission before it can be concluded that 5-ASA enemas are more cost effective than corticosteroids. To facilitate our analysis, we accepted the authors' definitions of disease

response and remission. The ability to pool data effectively from several trials is limited by the variability in outcome criteria. As meta-analyses and overviews are updated, it is essential that methods of diagnosis, definitions of active or inactive disease, and criteria for symptomatic, endoscopic, and histological outcomes are standardised.⁶⁷ Increased attention should also be given to the potential confounding influence of proximal disease margin, formulation of delivery vehicle, volume of preparation, and duration of treatment. As the potential for adverse reactions often governs selection among equally effective agents, adverse effects also must be reported more rigorously.

We conclude that treatment with rectal 5-ASA is superior to treatment with rectal corticosteroids in the management of active distal ulcerative colitis. Rectal budesonide seems to be as effective as conventional rectal corticosteroids, but seems to cause less suppression of endogenous cortisol production. Conventional rectal corticosteroids may be regarded as an alternative rectal treatment for active distal ulcerative colitis once aminosalicylates have failed, or in patients allergic to 5-ASA.

- 1 Truelove SC, Witts LJ. Cortisone in ulcerative colitis: preliminary report on a therapeutic trial. *BMJ* 1954; 2: 375-8.
- 2 Truelove SC. Treatment of ulcerative colitis with local hydrocortisone. *BMJ* 1956; 2: 1267-72.
- 3 Mulder CJJ, Tytgat GNJ. Topical corticosteroids in inflammatory bowel disease. *Aliment Pharmacol Ther* 1993; 7: 125-30.
- 4 Chapman NJ, Brown ML, Phillips SF, Tremaine WJ, Schroeder KW, Dewanjee MK, *et al*. Distribution of mesalamine enemas in patients with active distal ulcerative colitis. *Mayo Clin Proc* 1992; 67: 245-8.
- 5 Tiel-van Buul MMC, Mulder CJJ, van Royen EA, Wiltink EHH, Tytgat GNJ. Retrograde spread of mesalazine (5-aminosalicylic acid)-containing enema in patients with ulcerative colitis. *Clin Pharmacokinet* 1991; 20: 247-51.
- 6 Jay M, Digenis GA, Foster TS, Antonow DR. Retrograde spreading of hydrocortisone enema in inflammatory bowel disease. *Dig Dis Sci* 1986; 31: 139-44.
- 7 Campieri M, Lanfranchi GA, Brignola C, Bazzocchi G, Gionchetti P, Minguzzi MR, *et al*. Retrograde spread of 5-aminosalicylic acid enemas in patients with active ulcerative colitis. *Dis Colon Rectum* 1986; 29: 108-10.
- 8 Kruis W, Bull U, Eisenburg J, Paumgartner G. Retrograde colonic spread of sulphasalazine enemas. *Scand J Gastroenterol* 1982; 17: 933-8.
- 9 Nymann-Pantelidis M, Nilsson A, Wagner ZG, Borga O. Pharmacokinetics and retrograde colonic spread of budesonide enemas in patients with distal ulcerative colitis. *Aliment Pharmacol Ther* 1994; 8: 617-22.
- 10 Swarbrick ET, Loose H, Lennard-Jones JE. Enema volume as an important factor in successful topical corticosteroid treatment of colitis. *Proc Royal Soc Med* 1974; 67: 753-4.
- 11 Van Bodegraven AA, Boer RO, Lourens J, Tuynman HARE, Sindram JW. Distribution of mesalazine enemas in active and quiescent ulcerative colitis. *Aliment Pharmacol Ther* 1996; 10: 327-32.
- 12 Farthing MJG, Rutland MD, Clark ML. Retrograde spread of hydrocortisone containing foam given intrarectally in ulcerative colitis. *BMJ* 1979; 2: 822-4.
- 13 Hay DJ. Spreading characteristics of proprietary rectal steroid preparations. In: Wilson CG, Hardy JG, eds. *Radionuclide imaging in drug research*. London: Croom Helm, 1982: 71-80.
- 14 Wood E, Wilson CG, Hardy JG. The spreading of foam and solution enemas. *Int J Pharmaceut* 1985; 25: 191-7.
- 15 Campieri M, Corbelli C, Gionchetti P, Brignola C, Belluzzi A, Di Febo G, *et al*. Spread and distribution of 5-ASA colonic foam and 5-ASA enemas in patients with ulcerative colitis. *Dig Dis Sci* 1992; 37: 1890-7.
- 16 Wilding IR, Kenyon CJ, Chauhan S, Hooper G, Marshall S, McCracken JS, *et al*. Colonic spreading of a non-chloro-fluorocarbon mesalazine rectal foam enema in patients with quiescent ulcerative colitis. *Aliment Pharmacol Ther* 1995; 9: 161-6.
- 17 Williams CN, Haber G, Aquino JA. Double-blind, placebo-controlled evaluation of 5-ASA suppositories in active distal proctitis and measurement of extent of spread using 99m-Tc-labelled 5-ASA suppositories. *Dig Dis Sci* 1987; 32: 71S-5S.

- 18 Jay M, Beihn RM, Digenis GA, Deland FH, Caldwell L, Mlodozeniec AR. Disposition of radiolabelled suppositories in humans. *J Pharm Pharmacol* 1985; **37**: 266-8.
- 19 Powell Tuck J, Lennard-Jones JE, May CS, Wilson CG, Paterson JW. Plasma prednisolone levels after administration of prednisolone-21-phosphate as a retention enema in colitis. *BMJ* 1976; **1**: 193-5.
- 20 Lee DAH, Taylor GM, James VHT, Walker G. Plasma prednisolone levels and adrenocortical responsiveness after administration of prednisolone-21-phosphate as a retention enema. *Gut* 1979; **20**: 349-55.
- 21 Farmer RG, Schumacher OP. Treatment of ulcerative colitis with hydrocortisone enemas: relationship of hydrocortisone absorption, adrenal suppression, and clinical response. *Dis Colon Rectum* 1970; **13**: 355-61.
- 22 Luman W, Gray RS, Pendek R, Palmer KR. Prednisolone metasulphobenzoate foam retention enemas suppress the hypothalamo-pituitary-adrenal axis. *Aliment Pharmacol Ther* 1994; **8**: 255-8.
- 23 Cann PA, Holdsworth CD. Systemic absorption from hydrocortisone foam enema in ulcerative colitis. *Lancet* 1987; **i**: 922-3.
- 24 Bianchi Porro G, Prantera C, Campieri M, Petrillo M, Campanini MC, Gionchetti P, et al. Comparative trial of methylprednisolone and budesonide enemas in the treatment of active distal ulcerative colitis. *Eur J Gastroenterol Hepatol* 1994; **6**: 125-30.
- 25 Danielsson A, Hellers G, Lyrenas E, Lofberg R, Nilsson A, Olsson O, et al. A controlled randomized trial of budesonide versus prednisolone retention enemas in active distal ulcerative colitis. *Scand J Gastroenterol* 1987; **22**: 987-92.
- 26 Danish Budesonide Study Group. Budesonide enema in distal ulcerative colitis. *Scand J Gastroenterol* 1991; **26**: 1225-30.
- 27 Spencer CM, McTavish D. Budesonide: a review of its pharmacological properties and therapeutic efficacy in inflammatory bowel disease. *Drugs* 1995; **50**: 854-72.
- 28 Brattsand R. Overview of newer glucocorticosteroid preparations for inflammatory bowel disease. *Can J Gastroenterol* 1990; **4**: 407-14.
- 29 Marshall JK, Irvine EJ. Rectal aminosalicilate therapy for distal ulcerative colitis: a meta-analysis. *Aliment Pharmacol Ther* 1995; **9**: 293-300.
- 30 Schimmer BP, Parker KL. Adrenocorticotrophic hormone; adrenocortical steroids and their synthetic analogs; inhibitors of the synthesis and actions of adrenocortical hormones. In: Wonsiewica MJ, McCurdy P, eds. *Goodman and Gilman's The pharmacological basis of therapeutics*. New York: McGraw Hill, 1996: 466.
- 31 DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986; **7**: 177-80.
- 32 Bracken MB. Statistical methods for analysis of effects of treatment in overviews of randomized trials. In: Sinclair JC, Bracken MB, eds. *Effective care of the newborn infant*. Oxford: Oxford University Press, 1992: 15-6.
- 33 Breslow NE, Day NE (eds). *Statistical methods in cancer research*. Lyon: International Agency for Research on Cancer, 1980: 136-46.
- 34 Lennard-Jones JE, Baron JH, Connell AM, Avery Jones F. A double blind controlled trial of prednisolone-21-phosphate suppositories in the treatment of idiopathic proctitis. *Gut* 1962; **3**: 207-10.
- 35 Watkinson G. Treatment of ulcerative colitis with topical hydrocortisone hemisuccinate sodium: a controlled trial employing restricted sequential analysis. *BMJ* 1958; **2**: 1077-82.
- 36 Danielsson A, Lofberg R, Persson T, Salde L, Schioler R, Suhr O, et al. A steroid enema, budesonide, lacking systemic effects for the treatment of distal ulcerative colitis or proctitis. *Scand J Gastroenterol* 1992; **27**: 9-12.
- 37 Hanauer S, Robinson M. Dose ranging study of budesonide enemas in active distal ulcerative colitis [abstract]. *Gastroenterology* 1995; **108**: A832.
- 38 Hamilton I, Pinder IF, Dickinson RJ, Ruddel WSJ, Dixon MF, Axon ATR. A comparison of prednisolone enemas with low-dose oral prednisolone in the treatment of acute distal ulcerative colitis. *Dis Colon Rectum* 1984; **27**: 701-2.
- 39 Lennard-Jones JE, Longmore AJ, Newell AC, Wilson CWE, Avery Jones F. An assessment of prednisone, salazopyrin, and topical hydrocortisone hemisuccinate used as outpatient treatment for ulcerative colitis. *Gut* 1960; **1**: 217-22.
- 40 Bianchi Porro G, Ardizzone S, Petrillo M, Fasoli A, Molteni P, Imbesi V. Low Pentasa dosage versus hydrocortisone in the topical treatment of active ulcerative colitis: a randomized double blind study. *Am J Gastroenterol* 1995; **90**: 736-9.
- 41 Campieri M, Lanfranchi GA, Bazzocchi G, Brignola C, Sarti F, Franzin G, et al. Treatment of ulcerative colitis with high-dose 5-aminosalicylic acid enemas. *Lancet* 1981; **ii**: 270-1.
- 42 Danish 5-ASA Study Group. Topical 5-aminosalicylic acid versus prednisolone in ulcerative proctosigmoiditis. *Dig Dis Sci* 1987; **32**: 598-602.
- 43 Farup PG, Hovde O, Halvorsen FA, Raknerud N, Brodin U. Mesalazine suppositories versus hydrocortisone foam in patients with distal ulcerative colitis. *Scand J Gastroenterol* 1995; **30**: 164-70.
- 44 Friedman LS, Richter JM, Kirkham SE, DeMonaco HJ, May RJ. 5-Aminosalicylic acid enemas in refractory distal ulcerative colitis: a randomized controlled trial. *Am J Gastroenterol* 1986; **81**: 412-8.
- 45 Lee FI, Jewell DP, Mani V, Keighley MRB, Kingston RD, Record CO, et al. A randomised trial comparing mesalazine and prednisolone foam enemas in patients with acute distal ulcerative colitis. *Gut* 1996; **38**: 229-33.
- 46 Mulder CJJ, Endert E, Van der Heide H, Houthoff HJ, Wiersinga W, Wiltink EHH, et al. Comparison of 5-aminosalicylic acid (3g) and prednisolone phosphate sodium (enemas (30mg) in the treatment of distal ulcerative colitis: a prospective, randomized, double-blind trial. *Scand J Gastroenterol* 1988; **23**: 1005-8.
- 47 O'Donnell LJD, Arvind AS, Cameron D, Talbot IC, Jewell DP, Lennard-Jones JE, et al. Double blind, controlled trials of 4-aminosalicylic acid and prednisolone enemas in distal ulcerative colitis. *Gut* 1992; **33**: 947-9.
- 48 Sharma MP, Dufhare HV, Dasarthy S. A prospective randomized double blind trial comparing prednisolone and 4-aminosalicylic acid enemas in acute distal ulcerative colitis. *J Gastroenterol Hepatol* 1992; **7**: 173-7.
- 49 Lofberg R, Ostergaard Thomsen O, Langholz E, Schioler R, Danielsson A, Suhr O, et al. Budesonide versus prednisolone retention enemas in active distal ulcerative colitis. *Aliment Pharmacol Ther* 1994; **8**: 623-9.
- 50 Tarpila S, Turunen U, Seppala K, Aukee S, Pikkariainen P, Elomaa I, et al. Budesonide enema in active haemorrhagic proctitis: a controlled trial against hydrocortisone foam enema. *Aliment Pharmacol Ther* 1994; **8**: 591-5.
- 51 Lamers C, Meijer J, Engels L, Bos L, Van Hozegand R, Driessen W, et al. Comparative study of the topically acting glucocorticoid budesonide and 5-aminosalicylic acid enema therapy of proctitis and proctosigmoiditis [abstract]. *Gastroenterology* 1991; **100**: A223.
- 52 Lemann M, Galian A, Rutgeerts P, van Heuvelzwijn R, Cortot A, Viteau JM, et al. Comparison of budesonide and 5-aminosalicylic acid enemas in active distal ulcerative colitis. *Aliment Pharmacol Ther* 1995; **9**: 557-62.
- 53 Cobden I, Al-Mardini H, Zaitoun A, Record CO. Is topical therapy necessary in acute distal ulcerative colitis? Double-blind comparison of high-dose oral mesalazine versus steroid enemas in the treatment of active distal ulcerative colitis. *Aliment Pharmacol Ther* 1991; **5**: 513-22.
- 54 Grace RH, Gent AE, Hellier MD. Comparative trial of sodium cromoglycate enemas with prednisolone enemas in the treatment of ulcerative colitis. *Gut* 1987; **28**: 88-92.
- 55 Halpern Z, Sold O, Baratz M, Konikoff F, Halak A, Gilat T. A controlled trial of beclomethasone versus betamethasone enemas in distal ulcerative colitis. *J Clin Gastroenterol* 1991; **13**: 38-41.
- 56 Hanauer SB, Kirsner JB, Barrett WE. The treatment of left-sided ulcerative colitis with tixocortol pivalate [abstract]. *Gastroenterology* 1986; **90**: A1449.
- 57 Lennard-Jones JE. Betamethasone 17-valerate and prednisolone 21-phosphate retention enemas in proctocolitis. *BMJ* 1971; **3**: 84-6.
- 58 Mulder CJJ, Endert E, Van der Heide H, Houthoff HJ, Wiersinga W, Wiltink EHH, et al. Comparison of beclomethasone dipropionate (2 and 3mg) and prednisolone sodium phosphate enemas (30mg) in the treatment of ulcerative proctitis: an adrenocortical approach. *Neth J Med* 1989; **35**: 18-24.
- 59 Mulder CJ, Fockens P, Van der Heide H, Tytgat GNJ. A controlled randomized trial of beclomethasone dipropionate (3mg) versus 5-ASA (1g) versus the combination of both (3mg/1g) as retention enemas in active distal ulcerative colitis [abstract]. *Gastroenterology* 1994; **106**: A739.
- 60 Riley SA, Gupta I, Mani V. A comparison of sucralfate and prednisolone enemas in the treatment of active distal ulcerative colitis. *Scand J Gastroenterol* 1989; **24**: 1014-8.
- 61 Ruddell WSJ, Dickinson RJ, Dixon MF, Axon ATR. Treatment of distal ulcerative colitis (proctosigmoiditis) in relapse: comparison of hydrocortisone enemas and rectal hydrocortisone foam. *Gut* 1980; **21**: 885-9.
- 62 Van der Heide H, Van den Brandt-Gradel, Tytgat GNJ, Endert E, Wiltink EHH, Schipper MEI, et al. Comparison of beclomethasone dipropionate and prednisolone 21-phosphate enemas in the treatment of ulcerative proctitis. *J Clin Gastroenterol* 1988; **10**: 169-72.
- 63 Van Outryve M, Huble F, Van Eeghem P, De Vos M. Comparison of ridogrel versus prednisolone, both administered rectally, for the treatment of active ulcerative colitis [abstract]. *Gastroenterology* 1996; **110**: A1035.
- 64 Ilnycki A, Shanahan F, Anton PA, Cheang M, Bernstein CN. The placebo response in ulcerative colitis [abstract]. *Gastroenterology* 1996; **110**: A929.
- 65 Meyers S, Janowitz HD. The natural history of ulcerative colitis: an analysis of the placebo response. *J Clin Gastroenterol* 1989; **11**: 33-7.
- 66 Riley SA, Mani V, Goodman MJ, Dutt S, Herd ME. Microscopic activity in ulcerative colitis: what does it mean? *Gut* 1991; **32**: 174-8.
- 67 Irvine EJ. Assessing outcome in randomized clinical trials: inflammatory bowel disease. *Can J Gastroenterol* 1993; **7**: 561-7.