High proportion of mutant K-ras gene in pancreatic juice of patients with pancreatic cystic lesions

K Tateishi, M Tada, M Yamagata, H Isayama, Y Komatsu, T Kawabe, Y Shiratori, M Omata

Abstract

Background/Aims—It was recently reported that the quantitative analysis of mutant K-ras gene in pancreatic juice could be useful for the diagnosis of pancreatic cancer. This methodology was applied to patients with pancreatic cystic lesions.

Methods—DNA was extracted from pancreatic juice collected at the time of endoscopy with injection of secretin. The ratio of the K-ras mutant allele to the wild-type allele was measured by two methods to detect and semiquantify mutant K-ras gene: polymerase chain reaction/preferential homoduplex formation assay and enriched polymerase chain reaction/enzyme linked mini-sequence assay.

Results—A high frequency of K-ras mutation was detected (more than 2% of all K-ras genes) in six of 14 patients (43%) with pancreatic cysts. This frequency was similar to those detected in patients with pancreatic adenocarcinoma and in intraductal papillary neoplasms of the pancreas. In contrast, the frequency of mutation was low (less than 2%) in patients without either pancreatic cysts or pancreatic neoplasms.

Conclusions—K-ras gene mutation may be derived from duct cells in the pancreas with a high potential for tumorigenesis. Therefore careful follow up of patients with a pancreatic cyst is recommended if they are found to have a high level of the mutant gene in the pancreatic juice.

(Gut 1999;45:737-740)

Keywords: pancreas; cyst; K-ras; pancreatic adenocarcinoma; quantitative analysis

We and others have previously reported the presence of K-ras mutations in pancreatic secretions as a molecular marker for pancreatic carcinoma because this cancer is known for their high incidence.¹⁻⁴ The main problem in these studies was the occasional presence of these mutations in normal or inflammatory pancreatic duct cells.⁵⁻⁸ Thus we decided to quantify cells carrying mutations in order to solve the problem of duct cells derived from non-malignant tissue, and we showed that quantitative analysis could explain the difference between pancreatic neoplasms and non-neoplasms.⁹

Unlike qualitative detection of *ras* gene mutation, quantification of the mutant in pancreatic juice proved to be helpful for differential diagnosis of pancreatic tumours, as there was a higher amount of K-ras mutant associated with pancreatic carcinoma than in non-neoplastic cases. In a previous study, however, we found two cases that were exceptions: a high level of the mutant gene (more than 1%) was found in the pancreatic juice from two patients without pancreatic carcinoma or chronic pancreatitis.⁹ When we carefully re-examined the clinical features of these two cases, we noticed that the patients had cystic lesions in the pancreas that had no apparent relation to pancreatic neoplasms or chronic pancreatitis.

The most common cystic lesions of the pancreas are pseudocysts, accounting for more than half of pancreatic cystic lesions.^{10, 11} The other cysts are neoplastic ones such as mucinous and serous cystadenoma (carcinoma), solitary or multiple retention cysts, and congenital cysts.¹¹ Pancreatic cysts have been reported to co-exist in the pancreas of patients with carcinoma in situ, although the nature of the cysts is not clear.^{12, 13}

Therefore, in this study, we semiquantitatively analysed the mutant *ras* gene in pancreatic juice from patients with cystic lesions in the pancreas without pancreatic neoplasms. Two useful assay kits, enriched polymerase chain reaction (PCR)/enzyme linked mini-sequence assay (ELMA) and PCR/preferential homoduplex formation assay (PHFA), were used.

Patients and methods

PATIENTS

A total of 76 patients undergoing endoscopic retrograde cholangiopancreatography (ERCP) at Tokyo University Hospital were selected for quantitative analysis of *ras* gene mutation in the pancreatic juice. Of these, 14 patients with cystic lesions in the pancreas were analysed (table 1). These patients consisted of eight men and six women; their age ranged from 52 to 80 (mean 65). Pseudocysts or pancreatic cysts associated with pancreatic carcinoma were excluded. These patients had no history of pancreatitis, and the character of the pancreatic cyst was as follows: no obvious pancreatic neoplasms in the pancreas, no papillary structure

Abbreviations used in this paper: ERCP, endoscopic retrograde cholangiopancreatography; PHFA, preferential homoduplex formation assay; ELMA, enzyme linked mini-sequence assay; IPN, intraductal papillary neoplasms of the pancreas.

Department of
Gastroenterology,
University of Tokyo,
Tokyo, Japan
K Tateishi
M Tada
M Yamagata
H Isayama
Y Komatsu
Y Shiratori
M Omata

Department of
Endoscopy and
Endoscopic Surgery,
University of Tokyo,
Tokyo, Japan
T Kawabe

Correspondence to:
Dr M Tada, Department of
Gastroenterology, University
of Tokyo, 7-3-1 Hongo
Bunkyo-Ku, Tokyo, Japan
(113-8655).

Accepted for publication
11 May 1999

Table 1 Characteristics of patients with pancreatic cystic lesions

Patient	Sex	Age (y)	No of cysts	Size (mm)	Mutant <i>ras</i> gene
1	M	58	2	16	(3+)
2	F	71	3	17	(2+)
3	F	80	1	15	(2+)
4	F	62	4	20	(2+)
5	F	63	2	17	(2+)
6	M	69	5	25	(2+)
7	F	73	1	18	(1+)
8	M	61	2	10	(1+)
9	M	59	2	10	(1+)
10	M	53	1	15	(+/-)
11	M	73	2	30	(-)
12	F	62	>10	3	(-)
13	M	52	1	40	(-)
14	M	61	2	27	(-)

Semiquantitative analysis of the mutant gene was scored as (3+), (2+), (1+), (+/-), (-) according to the percentage of mutant *ras* gene; (3+), (2+), (1+), (+/-), and (-) represented more than 20%, 2–20%, 0.2–2%, less than 0.2%, and 0% (not detected) of the mutant respectively.

in the cyst, no tendency to grow, no dilatation of the main pancreatic duct, and normal papilla of Vater. The maximum size of the cysts ranged from 3 to 40 mm in diameter, with an average of 19 mm. The number of cysts ranged from one to more than 10 and the average was two. Twenty patients with pancreatic adenocarcinoma, 15 with intraductal neoplasms of the pancreas, six with chronic pancreatitis, and 21 without either pancreatic neoplasms or pancreatic cysts were also analysed for comparison (table 2).

METHODS

Pure pancreatic juice was collected from the pancreatic duct during ERCP with the injection of secretin. DNA was extracted from the pancreatic juice by the standard phenol/chloroform method.¹⁴ We recently obtained similar results to the semiquantitative assay of mutant *K-ras* gene using our modified allele specific PCR⁹ and two other methods: (a) PCR/PHFA (Wakunaga Pharmaceutical Corporation, Hiroshima, Japan and SRL Corporation, Tokyo, Japan); (b) enriched PCR/ELMA (BML Corporation, Tokyo, Japan).¹⁵ The two assays are available as handy kits. Therefore mutant *ras* gene was analysed by these two methods in this study.

Briefly, PCR/PHFA is based on strand competition during hybridisation between a double labelled amplicon, which was prepared from biotin and dinitrophenyl labelled primers, and an unlabelled amplicon.¹⁶ Under a precisely controlled temperature gradient, the preferential formation of a homoduplex over a heteroduplex occurred. After annealing, the identical sequence of the double labelled and unlabelled amplicons resulted in a low popula-

tion of rehybridised double labelled double stranded DNA as the result of strand exchange between them. Even when the two differed by a single base substitution, the double labelled molecule was regenerated efficiently because of preferential homoduplex formation. The regenerated double labelled molecules were captured on a streptavidin coated microtitre plate and quantified enzymatically with a chromogenic substance. This technique has been applied to HLA typing,¹⁶ and was also introduced for the semiquantitative analysis of mutant *K-ras* gene.

Enriched PCR/ELMA was based on the enrichment procedure of mutant *K-ras* gene described previously,¹⁷ and this was followed by the incorporation of biotin labelled nucleotide specific for the mutant gene, which was then quantified enzymatically with a chromogenic substance.¹⁸ Firstly, *K-ras* gene was amplified by PCR using a primer containing an artificial base substitution in order to mediate a restriction enzyme site that was limited to the wild type *K-ras* gene. After amplification, the amplicon was digested with a specific restriction enzyme in order to cleave a fragment of wild type allele and to select a fragment of the mutant allele. Second round PCR was performed to amplify the mutant gene selectively.¹⁷ Then the amplification product was captured by the oligomer which was designed to hybridise adjacent to the mutation site and was immobilised in a microtitration well. In a one nucleotide primer elongation step catalysed by *Taq* polymerase, a single biotin labelled nucleotide corresponding to a specific mutation was incorporated; this disclosed the presence of the mutation.^{19, 20} Finally, the biotin labelled extended oligomer was semiquantified enzymatically with a chromogenic substance.¹⁸

The results of the semiquantitative analysis were scored as (3+), (2+), (1+), (+/-), and (-) according to the percentage of mutant *ras* gene. All materials were measured by both assays. Because similar amounts of mutant gene were obtained to those previously reported,¹⁵ the amount obtained by enriched PCR/ELMA was used for the results. The scores (3+), (2+), (1+), (+/-), and (-) represented approximately more than 20%, 2–20%, 0.2–2%, less than 0.2%, and 0%—that is, not detected—of the mutant respectively according to the manufacturer. All codons (there are six types of mutation at codon 12 of *K-ras* gene) detected by the two methods were counted and added together in order to examine the proportion of GAT or GTT mutations, which were the most commonly detected (95%) in pancreatic carcinoma,¹⁹ because there was a difference in the type of mutant gene detected by the two methods.

Results

Mutant *ras* gene was detected in 10 of 14 patients with pancreatic cysts (fig 1). It was detected in high amounts ((2+) or more) in six of the 14 cases (43%). In cases of pancreatic carcinoma and intraductal papillary neoplasms of the pancreas (IPN), it was detected in high amounts ((2+) or more) in 10 of 20 cases (50%) and in 9 of 15 cases (60%)

Table 2 Clinical features of patients selected for quantitative analysis of *ras* gene mutation in pancreatic juice

	Total no patients	Male	Female	Age (years)	
				Mean	Range
Pancreatic cyst	14	8	6	65	52–80
Pancreatic carcinoma	20	10	10	65	40–82
Intraductal neoplasm of the pancreas	15	8	7	65	40–80
Chronic pancreatitis	6	3	3	59	31–70
Others	21	12	9	64	47–91

Figure 1 Amount of mutant *ras* gene according to patient classification. Mutant gene was detected in high amounts ((2+) or more) in patients with pancreatic carcinoma (Cancer), intraductal papillary neoplasms of the pancreas (IPN), and pancreatic cysts. In contrast, it was detected in low amounts ((1+) or less) in patients without these features.

Figure 2 Pancreatic cyst. Two typical cases of pancreatic cysts are shown by endoscopic ultrasonography. The maximum size of the cysts in all the positive patients ranged from 10 to 30 mm in diameter, with an average of two lesions per patient. Mutant *ras* gene was detected in high amounts ((2+) or more) in these patients.

respectively. In contrast, it was detected in low amounts ((1+) or less) in all the 27 patients without pancreatic neoplasms, pancreatic cysts, or chronic pancreatitis. This result was consistent with our previous report that the relative amount of K-*ras* mutant in the pancreatic juice from patients with pancreatic carcinoma and IPN was higher than in that of other patients.⁹ It was concluded that the proportion of mutant K-*ras* gene in the pancreatic juice from patients with pancreatic cystic lesions was as high as that from patients with pancreatic neoplasms.

With regard to the clinical characteristics of patients with pancreatic cysts, no correlation was found between high amount of *ras* gene mutant and age, sex, size of cysts, or number of cysts (table 1, fig 2). The average age of the six patients, two men and four women, was 67

Figure 3 Types of mutant *ras* gene according to patient classification. The proportion of codons GAT or GTT, which were the most commonly detected in pancreatic carcinoma, was analysed according to patient classification. All the codons detected by the two methods were counted and added together. The proportion of GAT or GTT was high in patients with pancreatic cysts and pancreatic neoplasms. It was relatively low in the other cases. Cancer, pancreatic carcinoma; IPN, intraductal papillary neoplasms of the pancreas.

years. The average number of cysts was three and the average maximum diameter was 18 mm.

As for the type of mutated codons, the numbers of codons detected by the two assays were 11 GAT, 10 GTT, three CGT, one TGT, one GCT, and one AGT, in patients with pancreatic cysts. The total number of codons was more than twice the number of patients with the mutation, because different kinds of mutants were detected in the same patient in some cases. The proportion of GAT or GTT was 78% (21 of all the 27 mutations detected by the two different assays) in this study (fig 3). It was 89% (41 of 46) in patients with pancreatic cancer and 84% (32 of 38) in those with IPN. It was relatively low (50%, six of 12) in the other cases. Therefore it was evident that the frequency of GAT or GTT in patients with pancreatic cysts was also high, similar to that in patients with pancreatic neoplasms.

Discussion

There are many kinds of cysts in the pancreas, more than two thirds being pseudocysts. Pseudocysts are formed when the duct system is ruptured, resulting in leakage of pancreatic juice. Most of them are usually preceded by a history of pancreatitis or trauma, and there has also been a rare case caused by a primary neoplasm of the pancreas.¹⁰ In this study, pseudocysts were excluded by clinical history and radiological findings, although our series may include cystic lesions associated with mild chronic pancreatitis, which was not diagnosed by imaging modalities. Other common pancreatic cystic lesions are neoplastic, retention, and congenital cysts.¹¹ Cystic neoplasms are divided into serous and mucinous adenomas (or adenocarcinomas). Serous cystadenoma is characterised by numerous tiny cysts ranging from 0.1 to 5 mm with a central scar.²² In our study, patient 12 had these characteristics, but mutant *ras* was not detected. In fact, it was reported that *ras* gene mutation was absent when serous cystadenoma tissue was analysed.²³ Mucinous cystadenoma is a cystic pancreatic tumour that occurs almost exclusively in women in the pancreatic body or tail. The tumours are solitary, ranging from 20 to 35 mm in diameter.¹¹ In our study, two patients (nos 3 and 7) with (2+) and (1+) amounts of mutant *ras* gene may be considered to belong to this category. The other kinds of cyst are solitary or multiple retention and congenital cysts.¹¹ Most congenital cysts are associated with von Hippel-Lindau disease or hereditary polycystic kidney disease.¹¹ In our series, there were no patients with clinical features of these diseases. Retention cysts occur secondary to obstruction of the pancreatic duct system and most of them are small.¹¹ Therefore most of the cystic lesions in patients other than nos 3, 7, and 12 in this study could be retention cysts.

A high rate of mutation of the K-*ras* gene in the pancreatic juice was detected unexpectedly in most patients with pancreatic cysts as well as in patients with pancreatic carcinoma and IPN.

The fact that the frequency of the mutational type of codon was also similar to that of pancreatic neoplasms may suggest that some neoplastic change had already occurred in pancreas with these cystic lesions, although such a lesion was not radiologically detected. The patients in our study showed no clinical or radiological features of pancreatic carcinoma or chronic pancreatitis. However, retention cysts are usually formed in association with chronic pancreatitis or pancreatic cancer.¹¹ Small and multiple cysts were found during 15% of 100 autopsies of patients with pancreatic cancer.²⁴ Another study reported that pancreatic cysts co-existed in the pancreas of patients with carcinoma in situ, and that this was the earliest feature of pancreatic cancer.^{12 13} These findings may indicate that small cystic lesions in the pancreas reflect the premalignant state of the pancreas.

An alternative possibility is that they represent early lesions of IPN. IPN is another type of pancreatic neoplasm. There is no definite name for this tumour, and IPN, intraductal papilloma, intraductal papillary mucinous tumour, mucin-producing cancer, mucin-hypersecreting neoplasm, and ductectatic cystadenoma or cystadenocarcinoma are all included in this category.^{25 26} These neoplasms, although heterogeneous with regard to their epithelial differentiation, form a tumour group that is morphologically and biologically distinct from ductal adenocarcinoma, mucinous cystic tumour, and ductal papillary hyperplasia. In IPN, there is a cystic dilatation of the main pancreatic duct or one of its branches. A widely dilated ampulla of Vater filled with mucoid material is often observed.¹¹ *Ras* gene mutation has been detected in this type of pancreatic tumour.^{25 27} Although typical cases of IPN were excluded by selecting cystic lesions without papillary structure in the cystic wall, without a tendency to grow, without dilatation of the main pancreatic duct, and with normal papilla of Vater, there is still a possibility that these cystic lesions are precursors of IPN before they actually show the characteristic clinical signs.

Of course, it is too early to draw any definite conclusions at this point. The number of patients is too small, and long term follow up of such patients will be required. However, little is known about the risk factors for pancreatic cancer, which makes it particularly difficult to find carcinoma of the pancreas at an early stage.²⁸ New radiological techniques such as endoscopic ultrasonography, which is superior to ERCP for the detection of small lesions in the pancreas, have been introduced,²⁹ and it has become easier to detect small cysts. Thus we should search carefully for small cystic lesions in the pancreas, and quantify *K-ras* mutation in the pancreatic juice if the presence of a pancreatic cyst is discovered. If the mutant gene is detected in high amounts, the cyst should be followed up carefully because it may undergo malignant changes.

- 1 Tada M, Omata M, Kawai S, et al. Detection of *ras* gene mutations in pancreatic juice and peripheral blood of patients with pancreatic adenocarcinoma. *Cancer Res* 1993;53:2472-4.
- 2 Kondo H, Sugano K, Fukayama N, et al. Detection of point mutations in the *K-ras* oncogene at codon 12 in pure pancreatic juice for diagnosis of pancreatic carcinoma. *Cancer* 1994;73:1589-94.
- 3 Trumper LH, Burger B, von Bonin F, et al. Diagnosis of pancreatic adenocarcinoma by polymerase chain reaction from pancreatic secretions. *Br J Cancer* 1994;70:278-84.
- 4 Van Laethem JL, Vertongen P, Deviere J, et al. Detection of *c-Ki-ras* gene codon mutations from pancreatic tumors. *Gut* 1995;36:781-7.
- 5 Tada M, Ohashi M, Shiratori Y, et al. Analysis of *K-ras* gene mutation in hyperplastic duct cells of the pancreas without pancreatic diseases. *Gastroenterology* 1996;110:227-31.
- 6 Yanagisawa A, Ohtake K, Ohnishi K, et al. Frequent *c-Ki-ras* oncogene activation in mucous cell hyperplasia of pancreas suffering from chronic inflammation. *Cancer Res* 1993;53:953-6.
- 7 Tada M. *Ki-ras* mutations in chronic pancreatitis: which relevance? *Gastroenterology* 1996;111:566-7.
- 8 Kondo H, Sugano K, Fukayama N, et al. Detection of *K-ras* gene mutations at codon 12 in the pancreatic juice of patients with intraductal papillary mucinous tumors of the pancreas. *Cancer* 1997;79:900-5.
- 9 Tada M, Teratani T, Komatsu Y, et al. Quantitative analysis of *ras* gene mutation in pancreatic juice for diagnosis of pancreatic adenocarcinoma. *Dig Dis Sci* 1998;43:15-20.
- 10 Hastings PR, Nance FC, Becker WF. Changing patterns in the management of pancreatic pseudocysts. *Ann Surg* 1975;181:546-51.
- 11 Solcia E, Capella C, Kloppel G. Tumors of the exocrine pancreas. *Atlas of tumor pathology, 3rd series, fascicle 20*. Washington, D.C.: Armed Forces Institute of Pathology, 1997.
- 12 Uehara H, Nakaizumi A, Tatsuta M, et al. Diagnosis of carcinoma in situ of the pancreas by peroral pancreatoscopy and pancreatic cytology. *Cancer* 1997;79:454-61.
- 13 Kimura W, Nagai H, Kuroda A, et al. Analysis of small cystic lesions of the pancreas. *Int J Pancreatol* 1995;18:197-206.
- 14 Sambrook J, Fritsch EF, Maniatis T. *Molecular cloning: a laboratory manual*. Cold Spring Harbor Laboratory, NY: Cold Spring Harbor Laboratory Press, 1989.
- 15 Tada M, Tateishi K, Yamagata M, et al. Quantitative analysis of *ras* gene mutation in pancreatic secretions for diagnosis of pancreatic carcinoma: comparison with three different assays. *Gastroenterology* 1998;114:G2047.
- 16 Oka T, Matsunaga H, Tokunaga K, et al. A simple method for detecting single base substitutions and its application to HLA-DPB1 typing. *Nucleic Acids Res* 1994;22:1541-7.
- 17 Levi S, Urbano-Ispizua A, Gill R, et al. Multiple *K-ras* codon 12 mutations in cholangiocarcinomas demonstrated with a sensitive polymerase chain reaction technique. *Cancer Res* 1991;51:3497-502.
- 18 Yamaguchi T, Yun S. Development of new assay for detecting mutant *K-ras* gene and its application for diagnosis of pancreatic cancer: enriched-PCR/ELMA. *Vita* 1997;14:50-54. (In Japanese.)
- 19 Brunetto MR, Randone A, Ranki M, et al. Quantitative analysis of wild-type and HBeAg minus hepatitis B viruses by a sequence-dependent primer extension assay. *J Med Virol* 1994;43:310-15.
- 20 Hawkins AE, Gilson RJC, Beath SV, et al. Novel application of a point mutation assay: evidence of transmission of hepatitis B virus with precore mutations and their detection in infants with fulminant hepatitis B. *J Med Virol* 1994;44:13-21.
- 21 Hruban RH, van Mansfield AD, Offerhaus GJ, et al. *K-ras* oncogene activation in adenocarcinoma of the human pancreas: a study of 82 carcinomas using a combination of mutant-enriched polymerase chain reaction analysis and allele-specific oligonucleotide hybridization. *Am J Pathol* 1993;143:545-54.
- 22 Yamaguchi K, Enjoji M. Cystic neoplasms of the pancreas. *Gastroenterology* 1987;92:1934-43.
- 23 Ishiwata T, Nakao A, Nomoto S, et al. Immunochemical and molecular biological studies of serous cystadenoma of the pancreas. *Pancreas* 1998;16:40-4.
- 24 Mikal S, Campbell AJ. A carcinoma of the pancreas. *Surgery* 1950;28:963-9.
- 25 Sessa F, Solcia E, Capella M, et al. Intraductal papillary-mucinous tumors represent a distinct group of pancreatic neoplasms: an investigation of tumour cell differentiation and *K-ras*, *p53* and *c-erbB-2* abnormalities in 26 patients. *Virchows Arch* 1994;425:357-67.
- 26 Loftus Jr EV, Olivares-Pakzad BA, Batts KP, et al. Intraductal papillary-mucinous tumors of the pancreas: clinicopathological features, outcomes, and nomenclature. *Gastroenterology* 1996;110:1909-18.
- 27 Tada M, Omata M, Ohto M. *Ras* gene mutations in intraductal papillary neoplasms of the pancreas. *Cancer* 1991;67:634-7.
- 28 Warshaw AL, Fernandez-del Castillo C. Pancreatic carcinoma. *N Engl J Med* 1992;326:455-65.
- 29 Nakaizumi A, Uehara H, Ishii H, et al. Endoscopic ultrasonography in diagnosis and staging of pancreatic cancer. *Dig Dis Sci* 1995;40:696-700.