PAPERS

The role of polyamines in gastric mucus synthesis inhibited by cigarette smoke or its extract

L Ma, W P Wang, J Y C Chow, S K Lam, C H Cho

Abstract

Background—Cigarette smoking was shown to delay gastric ulcer healing and reduce synthesis of mucus, which is important for gastric ulcer protection and healing. Polyamines are important in these processes.

Aims—To study the effects of cigarette smoking on the synthesis of mucus and to investigate if such an effect is acting by interference with the polyamine pathway.

Methods—Gastric mucosal ornithine decarboxylase activity, mucous secreting layer thickness, and ulcer size were determined after different concentrations of cigarette smoke exposure (0, 2, or 4%) in intact animals and animals with ulcers. Synthesis of mucus and ornithine decarboxylase activity and mRNA expression were also assessed in cigarette smoke extract treated MKN-28 cells.

Results—Exposure to cigarette smoke significantly reduced the thickness of the mucous secreting layer and gastric mucosal ornithine decarboxylase activity in animals with or without ulcers. Spermidine not only reversed inhibition of mucus synthesis in both intact and ulcer bearing animals but also reversed the delay in ulcer healing. Cigarette smoke extract significantly reduced mucus synthesis and ornithine decarboxylase activity but not its mRNA expression in MKN-28 cells. The reduction in mucus synthesis was restored by spermidine.

Conclusions—Cigarette smoke and its extract repress mucus synthesis in vivo and in vitro, respectively. Reduction of ornithine decarboxylase activity in gastric mucosa is closely associated with this effect.

(Gut 2000;47:170-177)

Keywords: mucus; ornithine decarboxylase; cigarette smoking; ulcers

increase mucus synthesis accelerate recovery from ulceration.⁵⁻⁷

Polyamines have been associated with cell proliferation during ulcer healing.⁸⁻⁹ A number of studies have demonstrated that polyamines are involved in epidermal growth factor (EGF) mediated gastroprotection, ulcer healing, and inhibition of acid secretion.¹⁰⁻¹¹ Ornithine decarboxylase (ODC) is the first rate limiting enzyme for the biosynthesis of polyamines, including spermine, spermidine, and putrescine. Both its activity and mRNA expression were elevated by EGF.¹¹⁻¹² Furthermore, EGF has been demonstrated to stimulate gastric mucus synthesis.¹³⁻¹⁴ Therefore, we thought it would be interesting to determine if ODC and polyamines are also involved in gastric mucus synthesis.

A clinical study showed that smokers had a lower level of acid mucosubstances in the gastric epithelium than non-smokers.¹⁵ However, because of the lack of an appropriate animal model for smoking, most studies have focused on the effect of nicotine on mucus synthesis and secretion¹⁶⁻¹⁷ as this compound is one of the major components of cigarette smoke and is easily accessible. Consequently, these studies reflect only part of the effects of cigarette smoking and cannot represent the action of smoking as a whole. Hence it would be useful to understand the exact relationship between cigarette smoking and mucus synthesis. We hypothesised that cigarette smoking might decrease mucus synthesis and secretion by interfering with the polyamine pathway, but little is known of the relationship between cigarette smoking and polyamine production or ODC activity in the gastric mucosa. Hence the aims of this study were to investigate: (1) how exposure to cigarette smoke affects mucus synthesis using a well established rat smoking model¹⁸ and the human gastric epithelial MKN-28 cell line; and (2) if ODC is involved in this process.

Abbreviations used in this paper: DFMO, DL- α -difluoromethyl-ornithine; EGF, epidermal growth factor; MTT, 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide; ODC, ornithine decarboxylase; PBS, phosphate buffered saline; TCA, trichloroacetic acid; PAS, periodic acid-Schiff; RT-PCR, reverse transcription-polymerase chain reaction.

Department of
Pharmacology, Faculty
of Medicine,
University of Hong
Kong, 5 Sassoon Road,
Hong Kong, China

L Ma
W P Wang
J Y C Chow
C H Cho

Department of
Medicine, Faculty of
Medicine, University
of Hong Kong,
5 Sassoon Road, Hong
Kong, China
S K Lam

Correspondence to:
Professor C H Cho. Email:
chcho@hkusua.hku.hk

Accepted for publication
8 February 2000

Epidemiological and animal studies have shown that cigarette smoking delays ulcer healing.¹⁻³ However, the underlying mechanisms are not fully elucidated. The physiological role of mucus is important in mucosal protection and ulcer healing.⁴ Drugs which

Materials and methods

CHEMICALS AND DRUGS

Chemicals and drugs were purchased from Sigma (St Louis, Missouri, USA) and organic solvents were purchased from Merck (Darmstadt, Germany), unless otherwise stated. In the animal study, spermidine was dissolved in distilled water and given orally at a dose of 100 mg/kg before each exposure to cigarette smoke.

ANIMALS AND INDUCTION OF GASTRIC KISSING ULCERS

The study was approved by the Committee on the Use of Live Animals for Teaching and Research in the University of Hong Kong. Male Sprague-Dawley rats (weighing 180–200 g) were reared on a standard laboratory diet (Ralston Purina Co., Chicago, Illinois, USA) and given tap water. They were kept in a room where temperature ($22\pm1^{\circ}\text{C}$), humidity (65–70%), and day/night cycle (12 hours/12 hours) were controlled. Rats were deprived of food but had free access to tap water 24 hours before ulcer induction. Gastric kissing ulcers were produced by luminal application of an acetic acid solution, as described by Tsukimi and Okabe¹⁹ with some modifications.³ Thereafter, animals were fed a standard diet of laboratory chow and given tap water ad libitum.

CIGARETTE SMOKE EXPOSURE

Camel cigarettes, without filters (R J Reynolds, Winston-Salem, North Carolina, USA) containing 1.2 mg of nicotine and 18 mg of tar per cigarette were used throughout the study. Intact or ulcerated rats (24 hours after ulcer induction) were exposed to different concentrations of cigarette smoke (0, 2, or 4%, vol/vol) in a chamber (39 cm $\times$ 23.5 cm $\times$ 21 cm) for one hour each day for several days. Procedures for exposure to cigarette smoke and the equipment used have been described previously.¹⁸ We have previously shown that this type of exposure does not affect the normal physiological functions of rats, such as acid/base balance, O_2/CO_2 in blood, heart rate, or blood pressure.¹⁸

Twenty four hours after the final exposure to cigarette smoke, rats were killed under ether anaesthesia by cervical dislocation. After washing with a cold saline solution, a longitudinal section of the stomach along the greater curvature (including the ulcer base and both sides of the ulcer margin in ulcerated stomachs) was obtained and fixed in 4% buffered formalin for 24 hours at 4°C for further histological study. The remaining glandular mucosa around the ulcer was scraped with a glass slide on an ice cold dish and immediately frozen in liquid nitrogen. Mucosal samples were stored at -70°C until assay.

EXTRACTION OF SUBSTANCES FROM CIGARETTE SMOKE

The substances in cigarette smoke were extracted using a smoke perfusion system. Briefly, smoke was forced through a series of bottles containing either 500 ml of chloroform (2 $\times$) or 95% ethanol (4 $\times$). The substances in smoke were thus absorbed in the solvents. The

substances dissolved in ethanol were re-extracted with chloroform. All of the chloroform extractions were collected together and then concentrated by a RE 47 rotary evaporator (Yamato Scientific Co. Ltd, Tokyo, Japan) connected to a cooling system (Julabo F10, Laborthechik GmbH, Seelbach, Germany). Our previous study showed that the chloroform extract but not the ethanol fraction potentiated ethanol induced gastric ulceration. Also, such fractions did not contain nicotine.²⁰ The chloroform extract was subsequently used in examining the effect of cigarette smoke components on mucus synthesis *in vitro*.

DETERMINATION OF MUCUS SYNTHESIS IN MKN-28 CELLS

MKN-28 is a secretory type of human gastric carcinoma cell line derived from a moderately differentiated tubular adenocarcinoma.²¹ The rate of mucus synthesis was determined by measuring incorporation of D-[6- ^3H] glucosamine into gastric mucosal glycoprotein according to the method of Takahashi and Okabe.²² Cells (1×10^5) were incubated in 24 well culture plates for 16 hours; they were then washed twice with $\text{Ca}^{++}/\text{Mg}^{++}$ free phosphate buffered saline (PBS), followed by incubation with 0.5 ml of the medium containing [^3H]glucosamine HCl (Amersham, Little Chalfont, UK) in the presence of various substances (cigarette extract, polyamines, and DL- α -difluoromethyl-ornithine (DFMO)), and/or corresponding vehicle at 37°C in 5% CO_2 for six hours. At the end of incubation, medium was aspirated and discarded. It was found that the amount of mucus glycoprotein secreted from cells into the medium was very low over the six hour incubation period. The remaining cells were washed twice with $\text{Ca}^{++}/\text{Mg}^{++}$ free PBS, solubilised with 0.4 ml of 0.3 mol/l NaOH, and neutralised with 0.4 ml of 0.3 mol/l HCl. The resulting aliquot was transferred to an Eppendorf tube. Then 0.5 ml of 50% trichloroacetic acid (TCA) was added to precipitate protein. Pellets were washed twice with 10% TCA, once with chloroform-methanol solution (1:1, vol: vol), and finally dried. These dried pellets were then dissolved in 25 μl of DMSO and the fractions were subjected to Sepharose CL-4B column chromatography for isolation of mucin, as described by Takahashi and Okabe.²² Radioactivity in the void fractions was measured as the amount of mucus synthesised by the cells. Mucus synthesis was expressed as [^3H] glucosamine incorporation (cpm) per 1×10^5 cells.

ESTIMATION OF CELL VIABILITY

Cell viability was estimated using 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyl-tetrazolium bromide (MTT).²³ MNK-28 cells ($2\times 10^4/100\ \mu\text{l}$ medium) were seeded into a 96 well plate and incubated overnight for attachment. They were then incubated with cigarette extract, various drugs, and/or vehicle at 37°C for six hours. At the end of incubation, the medium was aspirated. The remaining cells were incubated further with 0.25 mg/ml of MTT for three hours. MTT was extracted with 0.04 mol/l

HCl/isopropanol, and the colour change in the extract was measured at 595 nm. Cell viability was expressed as OD₅₉₅ per 2×10⁴ cells.

DETERMINATION OF ORNITHINE DECARBOXYLASE (ODC) ACTIVITY IN THE GASTRIC MUCOSA AND MKN-28 CELLS

ODC activity was assessed by measuring the amount of ¹⁴CO₂ liberated from DL-[1-¹⁴C] ornithine.²⁴ Mucosal samples were collected as previously described and homogenised in 67 mmol/l sodium potassium phosphate buffer (pH 7.4). The homogenates were centrifuged at 20 000 g, 4°C, for 20 minutes. The supernatants were used for assay of enzyme activity. Cultured MKN-28 cells were scraped from the culture plate and placed in 10 mmol/l Tris HCl buffer (pH 7.4). The cells were then sonicated for 15 minutes under ice cold conditions and centrifuged at 2500 g, 4°C, for 10 minutes. Again the supernatants were used for assay of ODC activity. A 300 µl aliquot of the supernatant was incubated in a stoppered test tube in the presence of 2.5 mmol/l of L-[1-¹⁴C]ornithine for 15 minutes at 37°C. The ¹⁴CO₂ liberated from the decarboxylation of ornithine was trapped by a piece of filter paper impregnated with 20 µl of 2.0 mol/l NaOH.

The paper was placed in a well connected to the stopper and suspended above the reaction mixture. The reaction was terminated by addition of 0.3 ml of 10% TCA. The radioactivity of ¹⁴CO₂ trapped in the filter paper was measured by a liquid scintillation counter (LS-6500; Beckman Instruments, Fullerton, California, USA). Intra-assay and interassay coefficients of variation were 7.6% and 9.1%, respectively. Protein content of the supernatant was measured by the Lowry method.²⁵ Enzyme activity was expressed as picomoles of ¹⁴CO₂ liberated per milligram of protein per hour.

ASSESSMENT OF THE MUCOUS SECRETING LAYER

Sections were stained with the periodic acid-Schiff (PAS) technique and counterstained with Mayer's haematoxylin.²⁶ The mucous secreting layer is a cellular layer within the mucosa which was assessed by measuring the relative thickness of the layer of cytoplasmic PAS stained positive cells using an image analyser (Q500IW, Leica Imaging Systems Ltd, Cambridge, UK) at 200× magnification by an investigator unaware of the treatments. This method washed away some of the very superficial mucous layer adherent to the luminal surface of the gastric mucosa.^{27, 28} However, most of the mucous secreting layer was found inside the mucosa which was largely preserved during the 24 hour fixation period. In normal stomach sections, 10 fields of measurements were taken and the results averaged. In ulcerated sections, two corresponding fields starting from the ulcer margins were measured and averaged. The results were calculated by determination of the length of the mucous secreting layer over total mucosal thickness (measured from the surface of the epithelium of the glandular mucosa to the point just above the submucosa).

RNA ISOLATION

After incubating MKN-28 cells in the presence of different concentrations of cigarette smoke extract or its vehicle for six hours, cells were washed twice with PBS. Total RNA was isolated from 15×10⁶ cells/5 cm diameter dish using TRIZOL Reagent (GibcoBRL, Gaithersburg, Maryland, USA). The cells were lysed directly by adding 1 ml of TRIZOL Reagent. The RNA containing aqueous phase was transferred to a fresh tube and mixed with 0.5 ml of isopropyl alcohol. The sample was centrifuged at 12 000 g for 10 minutes at 4°C. The RNA precipitate was washed with 75% ethanol and dried, and finally dissolved in 50 µl of DEPC water for reverse transcription-polymerase chain reaction (RT-PCR). The concentration of RNA was measured spectrophotometrically.

DETERMINATION OF ODC mRNA BY RT-PCR

RT-PCR was used to semiquantitate mRNA for ODC. Firstly, 4.0 µg of total RNA was used to generate the first strand of complementary DNA by using reverse transcriptase (Gibco BRL, Gaithersburg, Maryland, USA) following the manufacturer's instructions. PCR was performed for ODC and β-actin from the same complementary DNA samples using the following primer pairs. For ODC, the sense primer (5'-CCTTCGTGCAGGCAATCTCT) which recognises a sequence in exon 7 and the antisense primer (5'-GCTGCATGAGTTCCCACGCA) which recognises a sequence at the junction of exons 10 and 11 of human ODC cDNA were used.²⁹ For β-actin, the sense and antisense primers were 5'-AAGATGACCCAGATCATGTTTGAG and 5'-AGGAGGAGCAATGAT-CTTGATCTT, respectively. PCR amplification was initiated using a commercial PCR Core Kit (Boehringer, Mannheim, Germany). After the last cycle of amplification, samples were incubated for seven minutes at 72°C. PCR products (636 bp ODC fragment and 647 bp β-actin fragment) were visualised by UV illumination after electrophoresis through 1% agarose gels containing 0.5 µg/ml ethidium bromide. The gel photographs were submitted to a Multi-Analyst (Bio-Rad Laboratories, Hercules, California, USA). Semiquantitative analysis of ODC mRNA was performed by comparing the intensity of the PCR product of ODC to that of β-actin.

STATISTICAL ANALYSIS

Results are expressed as mean (SEM). Statistical analysis was performed using analysis of variance (ANOVA) and the unpaired Student's *t* test. *p* values less than 0.05 were considered statistically significant.

Results

EFFECTS OF CIGARETTE SMOKE EXPOSURE ON THE MUCOUS SECRETING LAYER AND ODC ACTIVITY IN INTACT AND ULCERATED RATS

As shown in fig 1, the thickness of the gastric mucous secreting layer inside the mucosa in different measured fields was similar in intact rats. Exposure to cigarette smoke for one hour once daily for three, six, or nine days

Figure 1 Effect of exposure to cigarette smoke on the mucous secreting layer in intact rats. Rats were exposed to cigarette smoke (0, 2, or 4%) for one hour each day for three, six, or nine days. Values are mean (SEM) of 7–8 rats. ** $p < 0.01$, *** $p < 0.001$ v 0% exposure.

significantly decreased the thickness of the gastric mucous secreting layer in a dose dependent manner. In parallel, gastric mucosal ODC activity was also substantially reduced by exposure to cigarette smoke (fig 2A). In rats with ulcers, the thickness of the gastric mucous secreting layer four and seven days after ulcer induction was maximal at the ulcer margin, decreasing towards the distant normal area, and ultimately becoming similar to that of the normal mucosa (fig 3A, B). Exposure to cigarette smoke significantly reduced the thickness of the mucous secreting layer in all fields along the full length of the mucosal section (fig 3). Similarly, gastric mucosal ODC activity in ulcer bearing rats was significantly decreased by cigarette smoke exposure on days 4 and 7 after ulcer induction (fig 2B).

Figure 2 Effect of exposure to cigarette smoke on gastric mucosal ornithine decarboxylase (ODC) activity in (A) intact and (B) ulcerated rats. ODC activity was determined by the amount of $^{14}\text{CO}_2$ liberated. Both intact and ulcerated rats (24 hours after ulcer induction) were exposed to cigarette smoke (0, 2, or 4%) for one hour each day for three or six days. ODC activity was measured in the gastric mucosa. * $p < 0.05$, ** $p < 0.01$ v 0% exposure.

Figure 3 Effect of exposure to cigarette smoke on the mucous secreting layer in rats, four (A) and seven (B) days after ulcer induction. Twenty four hours after ulcer induction, rats were exposed to cigarette smoke (0, 2, or 4%) for one hour each day for three or six days. Values are mean (SEM) of 10–12 rats. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ v 0% exposure.

EFFECTS OF CIGARETTE SMOKE EXTRACT ON MUCUS SYNTHESIS, ODC ACTIVITY, AND mRNA EXPRESSION IN MKN-28 CELLS

The amount of ^3H glucosamine incorporation in the mucin fraction was estimated as 75.71% of the whole cell preparation after Sepharose column chromatography. We used this column to determine mucus synthesis in the MKN-28 cell line. Incubation of MKN-28 cells with different concentrations of cigarette smoke extract (50, 100, or 200 $\mu\text{g}/\text{ml}$) for six hours significantly and dose dependently suppressed mucus synthesis, as indicated by a lower level of [^3H] glucosamine incorporation in the cultures exposed to smoke extract compared with the vehicle control (fig 4). Similar results were found when the cells were incubated for the same period of time with DFMO, an ODC specific inhibitor (fig 4). Further study demonstrated that cigarette smoke extract at concentrations of 100 and 200 $\mu\text{g}/\text{ml}$, which were sufficient to decrease mucus synthesis, also significantly reduced ODC activity in MKN-28 cells (fig 5). However, ODC mRNA expression was not affected by cigarette smoke extract at either dose. The ratios of mRNA expression of ODC to β -actin in vehicle and smoke extract (100 and 200 $\mu\text{g}/\text{ml}$) groups were 0.80 (0.02), 0.81 (0.02), and 0.81 (0.02), respectively. Meanwhile, cell viability was unchanged by smoke extract or DFMO at the concentrations used in this study (table 1A).

EFFECTS OF POLYAMINES ON MUCUS SYNTHESIS IN MKN-28 CELLS

Only spermidine at 0.1 mmol/ml showed a significant enhancing effect on mucus synthesis when MKN-28 cells were incubated with different polyamines for six hours. Putrescine and spermine at the concentrations used did not significantly affect mucus synthesis (fig 6A). Thereafter, incubation of cells with spermidine (0.1 and 1 mmol/ml) in the presence of smoke extract (100 mg/ml) significantly reversed extract induced inhibition of

Figure 4 Effects of cigarette smoke extract and DL- α -difluoromethyl-ornithine (DFMO) on mucus synthesis in MKN-28 cells. Cells were incubated with cigarette smoke extract, DFMO, or vehicle for six hours. Mucus synthesis was determined by the amount of [^3H] glucosamine incorporation. Values are mean (SEM) of six samples. *** $p < 0.001$ v vehicle.

Figure 5 Effect of cigarette smoke extract on ornithine decarboxylase (ODC) activity in MKN-28 cells. Cells were incubated with cigarette smoke extract or vehicle for six hours. ODC activity was determined by the amount of $^{14}\text{CO}_2$ liberated. Values are mean (SEM) of six samples. ** $p < 0.01$, *** $p < 0.001$ v vehicle.

mucus synthesis at the higher concentration of 1 mmol/ml (fig 6B). Again, cell viability was unaffected by the polyamines at the concentrations used in this study (table 1B, C).

EFFECTS OF SPERMIDINE ON THE MUCOUS SECRETING LAYER AND ULCER HEALING

In the animal study, it was found that cigarette smoke exposure at 4% for a one hour once daily for three days significantly reduced the thickness of the mucous secreting layer in both intact and ulcerated rats (fig 7A, B). These findings support our previous in vitro study. Intragastric administration of spermidine (100 mg/kg) before each exposure to cigarette smoke completely reversed inhibition of the mucous secreting cell layer induced by cigarette smoke in both intact and ulcer bearing rats (fig 7A, B) and significantly promoted ulcer healing in ulcerated animals (fig 8). Interestingly, spermidine at the same dose did not affect either the mucous secreting layer or ulcer healing in non-smoking control animals.

Table 1 Effects of different forms of treatment on cell viability of MKN-28 cells

Treatment	OD ₅₉₃ /2×10 ⁴ cells
A	
Vehicle	0.57 (0.01)
Extract	
50 $\mu\text{g}/\text{ml}$	0.55 (0.02)
100 $\mu\text{g}/\text{ml}$	0.57 (0.02)
200 $\mu\text{g}/\text{ml}$	0.54 (0.01)
DFMO	
10 mmol/ml	0.55 (0.01)
50 mmol/ml	0.56 (0.01)
B	
Vehicle	0.55 (0.03)
Putrescine	
0.1 mmol/ml	0.55 (0.01)
1 mmol/ml	0.59 (0.02)
Spermidine	
0.01 mmol/ml	0.57 (0.02)
0.1 mmol/ml	0.55 (0.02)
1 mmol/ml	0.56 (0.02)
Spermine	
0.1 mmol/ml	0.55 (0.02)
1 mmol/ml	0.54 (0.02)
C	
Vehicle	0.56 (0.03)
Extract 100 $\mu\text{g}/\text{ml}$	0.54 (0.02)
Extract+spermidine (0.1 mmol/ml)	0.54 (0.02)
Extract+spermidine (1 mmol/ml)	0.54 (0.02)

Values are mean (SEM) of six samples for each group.

Figure 6 Effects of polyamines on mucus synthesis in MKN-28 cells in the absence (A) and presence (B) of cigarette smoke extract. Cells were incubated with vehicle, polyamines, cigarette smoke extract, or a combination of spermidine and smoke extract for six hours. Mucus synthesis was determined by the amount of [^3H] glucosamine incorporation. Values are mean (SEM) of six samples. ** $p < 0.01$, *** $p < 0.001$ v vehicle; ††† $p < 0.001$ v cigarette smoke extract alone.

Figure 7 Effect of spermidine on inhibition of the mucous secreting layer induced by cigarette smoking in intact (A) and ulcerated (B) rats. Both intact and ulcerated rats (24 hours after ulcer induction) were exposed to cigarette smoke (0, 2, or 4%) for one hour each day for three days. Spermidine was administered intragastrically (100 mg/kg) before each exposure to cigarette smoke. Values are mean (SEM) of 7–8 rats. ** $p < 0.01$ v 0% exposure; ††† $p < 0.01$ v 4% alone.

Discussion

Cigarette smoking has been reported to increase both the incidence³⁰ and recurrence^{31, 32} of gastric and duodenal ulcers and also to delay ulcer healing.^{1, 33} Our previous studies showed that both cigarette smoke and its extract derived from cigarette filters potentiated ethanol induced gastric mucosal damage and delayed the ulcer healing rate in rats.^{3, 20} In the present study, cigarette smoke and its extract decreased the thickness of the mucous secreting layer inside the mucosa and also mucus synthesis in vitro, respectively (figs 1, 3, 4), indicating that modulation of gastric mucus may be part of the mechanism by which cigarette smoking intensifies ulceration and

impairs ulcer healing. However, how cigarette smoking affects mucus synthesis is still unclear. Gastric mucus is secreted by epithelial cells and could be modulated by a number of factors secreted or synthesised locally in the gastric mucosa.^{34, 35}

Polyamines are involved in many EGF mediated bioactivities including cell proliferation and acid secretion in the stomach.^{11, 36} Sialoadenectomised rats had reduced levels of gastric mucin³⁷ indicating that salivary EGF is associated with gastric mucus synthesis. There is evidence that smokers have lower levels of salivary EGF than non-smokers,³⁸ suggesting that salivary EGF production is impaired by cigarette smoking. Therefore, it would be help-

Figure 8 Effect of spermidine on the delay in ulcer healing induced by cigarette smoking, four days after ulcer induction. Twenty four hours after ulcer induction rats were exposed to cigarette smoke (0, 2, or 4%) for one hour each day for three days. Spermidine was administered intragastrically (100 mg/kg) before each exposure. Values are mean (SEM) of 7–8 rats. ** $p < 0.01$ v 0% exposure; †† $p < 0.01$ v 4% alone.

ful to determine if polyamine synthesis is affected by cigarette smoking and the role of polyamines in the regulation of gastric mucus production. However, the relationships between cigarette smoking, polyamine production, and mucus synthesis in the gastric mucosa has never been investigated.

Using the present modified method to determine mucus synthesis in an isolated cell preparation, only spermidine showed a significant enhancing effect on mucus synthesis (fig 6A). This may be explained by the fact that different polyamines contributed differently to certain bioactivities.³⁹ Further study demonstrated that cigarette smoke extract decreased mucus synthesis, accompanied by a reduction in ODC activity in a dose dependent manner in MKN-28 cells (figs 4, 5). The former action was completely reversed by spermidine at a concentration of 1 mmol/l (fig 6B). DFMO, an ODC inhibitor, also induced a similar reduction in mucus synthesis (fig 4). These findings indicate that suppression of ODC activity is likely to be one of the mechanisms of inhibition of mucus synthesis induced by cigarette smoke extract. This effect was further supported by the in vivo study which showed that cigarette smoke exposure decreased the mucous secreting layer, accompanied by a reduction in mucosal ODC activity in both intact and ulcer bearing animals (fig 2). Oral administration of spermidine (100 mg/kg) markedly antagonised inhibition of the mucous secreting layer caused by cigarette smoke exposure (fig 7). Parallel to restoration of the mucous secreting layer by spermidine, the delay in ulcer healing was also substantially reversed (fig 8). Taken together, the in vitro and in vivo studies indicate that impairment of mucus synthesis as a result of suppression of mucosal ODC activity may

contribute to the delay in ulcer healing induced by cigarette smoking. Interestingly, the thickness of the gastric mucous secreting layer after ulcer induction was substantially increased at the ulcer margin (figs 3, 7B). This may protect the newly proliferated cell against injury from the aggressive factors in the lumen. Furthermore, it may also provide the correct pH for epithelial cell restitution⁴⁰ and binding of growth factors to their receptors at the ulcer margin.⁴¹ All of these actions could help in ulcer healing. Cigarette smoke exposure reduced the mucous secreting layer at the ulcer margin. This may be one of the mechanisms by which cigarette smoking delays ulcer healing. In the present study, ulcer induction itself significantly increased ODC activity in the gastric mucosa (fig 2). This is consistent with the finding that stress induced gastric mucosal damage activated ODC activity together with ulcer healing.^{8, 9} Unlike mucosa with depressed ODC activity in smoking animals, supplementation of spermidine at the same dose did not have any effect on mucus synthesis in non-smoking animals.

The finding that cigarette smoke extract did not affect ODC mRNA expression in MKN-28 cells was surprising. Nevertheless, this suggests that the decrease in ODC activity was not caused by suppression of ODC gene activation. It is not known if translation to ODC protein, structural modification, or degradation was affected and further investigation is needed. However, it is known that mammalian ODCs are highly unstable. Their activities are dependent on many factors, including the low molecular weight thiols, pyridoxal phosphate.⁴² It has been suggested that in the absence of thiols, purified ODC polymerise through thiol disulphide interchanges and that polymerised forms have much less or no catalytic activity.⁴³ It is likely that cigarette smoke extract could impair ODC activity through interfering with the interaction between ODC and its cofactors. Our study showed that the smoke extract used in this study did not contain alkaloid, such as nicotine²⁰ and yet markedly inhibited synthesis of mucus in gastric epithelial cells. This observation suggests that nicotine may not be the culprit for the acute adverse action of cigarette smoking on gastric ulceration.

In summary, although the present experimental conditions may not fully reflect the situation of active smoking, they are similar to passive smoking in humans. More importantly, we have demonstrated for the first time that short term exposure to cigarette smoke decreased mucosal ODC activity which subsequently impaired mucus synthesis. Administration of spermidine completely restored the inhibited gastric mucus synthesis and delayed ulcer healing. Because of the wide distribution of mucus in the gastrointestinal tract, and its pivotal role in protection and repair of the gastric mucosa, our findings may have substantial relevance in both basic and clinical settings. Hence our results have unveiled the possible mechanisms by which cigarette smoking adversely affects the gastrointestinal tract and may also have significant implications in the

treatment of ulcer disorders, especially those with a delayed healing rate, as well as other diseases where reduction of mucus secretion has been implicated.

The authors would like to thank the Hong Kong Research Grant Council for support and also Dr S Takahashi for advice on the determination of mucus synthesis in MKN-28 cells.

- 1 Doll R, Jones FA, Pygott F. Effect of smoking on the production and maintenance of gastric and duodenal ulcers. *Lancet* 1958;1:657-62.
- 2 Iwata F, Leung FW. Tobacco cigarette smoke aggravates gastric ulcer in rats by attenuation of ulcer margin hyperemia. *Am J Physiol* 1995;268:G153-60.
- 3 Ma L, Chow JYC, Cho CH. Cigarette smoking delays ulcer healing: role of constitutive nitric oxide synthase in rat stomach. *Am J Physiol* 1999;39:G238-48.
- 4 Allen A, Hutton DA, Leonard AJ, et al. The role of mucus in the protection of the gastroduodenal mucosa. *Scand J Gastroenterol* 1986;21(suppl 125):71-7.
- 5 Slomiany BL, Laszewicz W, Slomiany A. In vitro inhibition of peptic degradation of porcine gastric mucus glycoprotein by sucralfate. *Scand J Gastroenterol* 1985;20:857-60.
- 6 Dai S, Ogle CW, Cho CH. Effects of carbenoxolone sodium on gastric and duodenal mucus synthesis in mice. *Pharmacology* 1986;33:58-60.
- 7 Li KM. Effect of zinc sulfate on acetic acid-induced gastric ulceration in rats. *J Pharm Pharmacol* 1990;42:657-9.
- 8 Wang JY, Johnson LR. Role of ornithine decarboxylase in repair of gastric mucosal stress ulcers. *Am J Physiol* 1990;258:G78-85.
- 9 Brzozowski T, Konturek SJ, Majka J, et al. Epidermal growth factor, polyamines, and prostaglandins in healing of stress-induced gastric lesions in rat. *Dig Dis Sci* 1993;38:276-83.
- 10 Konturek SJ, Brzozowski T, Konturek PK, et al. Role of salivary glands and epidermal growth factor (EGF) in gastric secretion and mucosal integrity in rats exposed to stress. *Regul Pept* 1991;32:203-15.
- 11 Wojciechowski K, Trzeciak L, Konturek SJ, et al. Inhibition of acid secretory response and induction of ODC and its mRNA by TGF- α and EGF in isolated rat gastric glands. *Regul Pept* 1995;56:1-8.
- 12 Tsujikawa T, Bamba T, Josoda S. The trophic effect of epidermal growth factor on morphological changes and polyamine metabolism in the small intestine of rats. *Gastroenterol Jpn* 1990;25:328-34.
- 13 Kelly SM, Hunter JO. Epidermal growth factor stimulates synthesis and secretion of mucus glycoprotein in human gastric mucosa. *Clin Sci* 1990;79:425-7.
- 14 Yoshida S, Kasuga S, Hirao Y, et al. Effect of biosynthetic human epidermal growth factor on the synthesis and secretion of mucin glycoprotein from primary culture of rabbit fundal mucosal cells. *In Vitro Cell Dev Biol* 1987;23:460-4.
- 15 Hollanders D, Morrissey SM, Ward PM. Histochemistry of gastric mucins in smokers and non-smokers. *Br J Clin Pract* 1986;40:74-7.
- 16 Wong SH, Ogle CW, Cho CH. The influence of chronic or acute nicotine pretreatment on ethanol-induced gastric ulceration in the rat. *J Pharm Pharmacol* 1986;38:537-40.
- 17 Luk ISC, Ho J, Wong WM, et al. Influence of chronic nicotine intake and acute ethanol challenge on gastric mucus level and blood flow in rabbits. *Digestion* 1994;55:399-404.
- 18 Chow JYC, Ma L, Cho CH. An experimental model for studying passive cigarette smoking effects on gastric ulceration. *Life Sci* 1996;58:2415-22.
- 19 Tsukimi Y, Okabe S. Validity of kissing gastric ulcers induced in rats for screening of antiulcer drug. *J Gastroenterol Hepatol* 1994;9:S60-5.
- 20 Chow JYC, Ma L, Zhu M, et al. The potentiating actions of cigarette smoking on ethanol-induced gastric mucosal damage in rats. *Gastroenterology* 1997;113:1188-97.
- 21 Hojo H. Establishment of cultured cell lines of human stomach cancer origin and their morphological characteristics. *Niigata Igakukai Zasshi* 1977;91:731-50.
- 22 Takahashi S, Okabe S. Role of extracellular Ca⁺⁺ and calmodulin in roxatidine-stimulated secretion and synthesis of mucus by cultured rabbit gastric mucosal cells. *J Pharmacol Exp Ther* 1998;284:37-42.
- 23 Mosmann T. Rapid colorimetric assay for cellular growth and survival: Application to proliferation and cytotoxicity assays. *J Immunol Methods* 1983;65:55-63.
- 24 Russell D, Snyder SH. Amine synthesis in rapidly growing tissues: ornithine decarboxylase activity in regenerating rat liver, chick embryo and various tumors. *Proc Natl Acad Sci USA* 1968;60:1420-7.
- 25 Lowry OH, Rosebrough NJ, Farr AL, et al. Protein measurement with the folin phenol reagent. *J Biol Chem* 1951;193:265-75.
- 26 Boon ME, Drijver JS. *Routine cytological staining techniques— theoretical background and practice*. London: Macmillan Education Ltd, 1986.
- 27 Jordan N, Newton J, Pearson J, et al. A novel method for the visualization of the in situ mucus layer in rat and man. *Clin Sci* 1998;95:97-106.
- 28 Pullan RD, Thomas GAO, Rhodes M, et al. Thickness of adherent mucus gel on colonic mucosa in humans and its relevance to colitis. *Gut* 1994;35:353-9.
- 29 Hettinen J, Halmekytö M, Alhonen L, et al. Levels of ornithine decarboxylase genomic sequences, heterogeneous nuclear RNA and mRNA in human myeloma cells resistant to α -difluoromethylornithine. *Biochem J* 1991;278:871-4.
- 30 Kato I, Momura AMY, Stemmermann GN, et al. A prospective study of gastric and duodenal ulcer and its relation to smoking, alcohol, and diet. *Am J Epidemiol* 1992;135:521-30.
- 31 Korman MG, Hansky J, Eaves ER, et al. Influence of cigarette smoking on healing and relapse in duodenal ulcer disease. *Gastroenterology* 1983;85:871-4.
- 32 Sontag S, Graham DY, Belsito A, et al. Cimetidine, cigarette smoking, and recurrence of duodenal ulcer. *N Engl J Med* 1984;311:689-93.
- 33 Korman MG, Shaw RG, Hansky J, et al. Influence of smoking on healing rate of duodenal ulcer in response to cimetidine or high-dose antacid. *Gastroenterology* 1981;80:1451-3.
- 34 Lamont JT, Ventola AS, Maull EA, et al. Cysteamine and prostaglandin F2 beta stimulate rat gastric mucin release. *Gastroenterology* 1983;84:306-13.
- 35 Brown JF, Hanson PJ, Whittle BJ. Nitric oxide donors increase mucus gel thickness in rat stomach. *Eur J Pharmacol* 1992;223:103-4.
- 36 Konturek JW, Brzozowski T, Konturek SJ. Epidermal growth factor in protection, repair, and healing of gastroduodenal mucosa. *J Clin Gastroenterol* 1991;13(suppl 1):S88-97.
- 37 Sarosiek J, Bilski J, Murty VLN, et al. Role of salivary epidermal growth factor in the maintenance of physicochemical characteristics of oral and gastric mucosal mucus coat. *Biochem Biophys Res Commun* 1988;152:1421-7.
- 38 Konturek JW, Bielanski W, Konturek SJ, et al. Distribution and release of epidermal growth factor in man. *Gut* 1989;30:1194-200.
- 39 Pegg AE, McCann PP. Polyamine metabolism and function. *Am J Physiol* 243:C212-21.
- 40 Wallace JL, Granger DN. The cellular and molecular basis of gastric mucosal defense. *FASEB J* 1996;10:731-40.
- 41 Szabo S, Hollander D. Pathways of gastrointestinal protection and repair: Mechanisms of action of sucralfate. *Am J Med* 1989;86:23-31.
- 42 Bey P, Danzin C, Jung M. Inhibition of basic amino acid decarboxylase involved in polyamine biosynthesis. In: McCann PP, Pegg AE, Sjoerdsma A, eds. *Inhibition of polyamine metabolism—Biological significance and basis for new therapies*. London: Academic Press, 1987:1-27.
- 43 Jänne J, Williams-Ashman HG. On the purification of L-ornithine decarboxylase from rat prostate and effects of thiol compounds on the enzyme. *J Biol Chem* 1971;246:1725-32.