STOMACH

Bile reflux gastritis and intestinal metaplasia at the cardia

M F Dixon, N P Mapstone, P M Neville, P Moayyedi, A T R Axon

Gut 2002;51:351–355

See end of article for
authors' affiliations

Correspondence to:
Professor M F Dixon,
Department of
Histopathology, General
Infirmary at Leeds, Leeds
LS1 3EX, UK;
miked@pathology.leeds.ac.uk

Accepted for publication
28 January 2002

Background and aims: Intestinal metaplasia (IM) at the cardia is likely to be a precursor of cardia cancer. Previous work has shown that it is associated with chronic inflammation attributable to either gastro-oesophageal reflux disease (GORD) or *Helicobacter pylori* infection. An alternative aetiological factor is bile reflux. Duodenogastric reflux brings about histological changes in the gastric mucosa that can be graded and used to calculate a bile reflux index (BRI). We used the BRI to assess whether reflux of bile plays a part in the development of cardia IM.

Methods: Histological changes in simultaneous gastric antrum and cardia biopsies from 267 dyspeptic patients were independently graded by two pathologists. The association between cardia IM and age, sex, clinical group, *H pylori* status, increased BRI (>14), and inflammation at the cardia were evaluated using logistic regression.

Results: A total of 226 patients had adequate cardia and antral biopsies; 149 had GORD and 77 had non-ulcer dyspepsia. Cardia IM was present in 66 (29%) patients, of whom 28 (42%) had complete IM. Increasing age, male sex, chronic inflammation, and a high BRI emerged as significant independent associations with cardia IM. Clinical group and *H pylori* status were not independent risk factors.

Conclusions: Histological evidence of bile reflux into the stomach is associated with cardia IM. This could have an important bearing on carcinogenesis at this site.

The cardia is an area of controversy. Conventionally this "small and ill defined" region is described as a zone 0.5–3 cm long "just distal to the lower end of the oesophagus" where it is interposed between oxyntic mucosa and squamous epithelium.^{1,2} However, its existence as a normal anatomical structure has been challenged. Chandrasoma and colleagues³ claim that the cardia is absent in truly normal individuals so that oxyntic mucosa abuts directly onto squamous epithelium. They argue that cardia-type mucosa is always acquired and is a consequence of gastro-oesophageal reflux disease (GORD). Cardia mucosa closely resembles the gastric antrum and it is suggested that the "acquired" cardia is an example of gastric metaplasia in response to acid reflux.³

Intestinal metaplasia (IM) at the cardia has attracted further controversy. Its very frequency (15–22% of dyspeptic subjects with an endoscopically normal lower oesophagus^{4–6}) argues against it being closely associated with cardia cancer yet by analogy with the stomach, cardia IM is considered to be a premalignant condition.⁷ While gastric metaplasia can be readily understood as an adaptive response of squamous epithelium to acid injury, it seems likely that intestinal metaplasia is triggered by other factors. Previous work has implicated gastric *Helicobacter pylori* infection and the associated "carditis"⁸ but other investigators have dismissed this in favour of GORD.⁹ The latter conclusion does not necessarily implicate acid as the sole causative agent. There is increasing evidence that GORD is accompanied by duodenogastric reflux so that the oesophageal refluxate can contain bile and other components of pancreaticoduodenal secretion.¹⁰ It would be much more plausible for the change from gastric to intestinal phenotype at the cardia to be related to bile than to acid reflux alone.

To investigate the role of bile reflux in the causation of cardia IM, we used a histological indicator of bile injury to the antrum—the bile reflux index (BRI)¹¹—along with the extent of cardia inflammation and *H pylori* status to search for possible aetiological associations. Additionally, as the various subtypes of IM at the cardia are likely to carry different risks for progression to cancer, we sought to relate these to the same factors.

PATIENTS AND METHODS

Over a two year period, the Department of Pathology at Leeds General Infirmary received simultaneous biopsies of the gastro-oesophageal junction and gastric mucosa from 267 patients. The majority (192 cases) consisted of patients referred directly by their general practitioner to an open access dyspepsia clinic. Suitable patients were seen as part of an assessment procedure prior to possible recruitment into a study evaluating the symptomatic response of *H pylori* negative dyspepsia to pantoprazole. Patients aged 18–80 years were considered. Those taking non-steroidal anti-inflammatory drugs or proton pump inhibitors less than four weeks prior to endoscopy, those who had undergone prior gastric or oesophageal surgery, and patients who were known to have been *H pylori* positive in the past or had received *H pylori* eradication therapy were excluded. Prior to endoscopy, patients were assessed by an experienced gastroenterologist (PMN) and categorised on clinical grounds as having either non-ulcer dyspepsia (NUD) or GORD. The diagnosis of NUD was based on a clinical presentation of predominant epigastric pain or abdominal symptomatology (for example, bloating or nausea) with clinically insignificant or absent heartburn. The diagnosis of GORD was based on a clinical presentation of significant heartburn or acid reflux with or without epigastric pain. At endoscopy, four quadrantic biopsies were taken from immediately below the squamocolumnar junction (Z line), two biopsies from the gastric body (one on the lesser curve and one on the greater curve), one from the incisura angularis, and two from the prepyloric antrum. The local ethics committee gave approval for the biopsies with prior consent.

Classification of patients into NUD and GORD was modified after endoscopy. Patients with active ulcer disease, macroscopic evidence of Barrett's oesophagus, severe (>grade 3) oesophagitis, and neoplasia were excluded. Those patients

Abbreviations: IM, intestinal metaplasia; GORD, gastro-oesophageal reflux disease; NUD, non-ulcer dyspepsia; BRI, bile reflux index; AB/PAS, alcian blue/periodic acid-Schiff;

Figure 1 (A) Cardia biopsy (squamocolumnar junction) stained with alcian blue/periodic acid-Schiff for mucins showing widespread incomplete-type intestinal metaplasia. (B) Antral biopsy from the same patient (haematoxylin-eosin) showing widespread intestinal metaplasia (complete type) (grade=3), mild lamina propria oedema (grade=1), and mild chronic inflammation (grade=1). *Helicobacters* were not present in this case. Based on these findings the bile reflux index was 20.

who had been classified as NUD on clinical grounds who on endoscopy were found to have Savary grade 2–3 oesophagitis were reclassified as GORD on the grounds that significant endoscopic oesophagitis is a more powerful predictor of GORD than clinical assessment.

The remaining 75 patients had cardia biopsies taken for some clinical indication, for example as an adjunct to the his-

tological diagnosis of GORD or because an endoscopic abnormality (inflammation or nodularity) was seen at the junction. These patients also had no evidence of Barrett's oesophagus, active peptic ulceration, or previous gastric surgery, and at endoscopy at least two biopsies from immediately below the Z line, two corpus, and two antral biopsies were taken.

These 75 patients were not assessed prospectively but their notes were obtained and scrutinised retrospectively (PMN). Clinical histories had been taken at the time of endoscopy and were documented in the notes with the endoscopy report. These patients were categorised into GORD and NUD groups using the same criteria as for the prospectively recruited patients.

Histological assessment

Biopsies were fixed in 10% formalin, processed, sectioned at three levels, and stained by haematoxylin-eosin. Additional sections taken at the second level from all biopsies were stained with alcian blue/periodic acid-Schiff (AB/PAS) to demonstrate IM and by the modified Giemsa stain for *H. pylori*. The cardia and gastric biopsies were assessed separately by two pathologists (NPM and MFD, respectively) who were "blind" to each other's findings and to the clinical details. The "cardia" biopsies were categorised as either squamous, oxyntic, oxyntocardiac transitional zone, or cardia type. We accepted as cardia those biopsies composed of gastric-type foveolar and surface epithelium overlying either antral-like mucous glands devoid of parietal cells or mucous glands with occasional parietal cells. The transitional mucosa was characterised by an admixture of mucous and oxyntic glands. A case was only included when there was at least one cardia biopsy and at least one antral biopsy adequate for assessment.

The presence of acute inflammation (that is, neutrophil polymorph infiltration), chronic inflammation (CI)—lymphocytes and plasma cells—and IM in the cardia and antrum were graded on a 0–3 scale (absent, mild, moderate, and marked), while additionally in the antrum, lamina propria oedema (Oed) and *H. pylori* colonisation density (*Hp*) were graded. The grading approach was analogous to that described in the updated Sydney system¹² although small numbers of mononuclear cells at the cardia were considered "normal" (see below). The Giemsa stained corpus biopsies were also examined before a case was declared *H. pylori* negative. IM was defined by the finding of fully formed goblet cells in the AB/PAS stain. When detected, a new section was cut and stained with Gomori's aldehyde fuchsin technique to distinguish between type II and III (sulphomucin positive) IM.¹³

Table 1 Grades of acute and chronic cardia inflammation according to clinical group, *Helicobacter pylori* (*Hp*) status, and bile reflux index (BRI). Grades 1, 2, and 3 are equivalent to a mild, moderate, and marked increase. For acute inflammation grade 0=absence of polymorphs; for chronic inflammation 0="normal" numbers of lymphocytes and plasma cells (see text)

Group	n	Acute				Chronic			
		0	1	2	3	0	1	2	3
NUD	77	59	14	4	0	35	26	14	2
GORD	149	122	20	7	0	83	44	20	0
<i>Hp</i> +	44	10	24	10	0	2	15	25	2
<i>Hp</i> -	182	171	10	1	0	118	55	9	0
BRI ≤ 14	188	151	29	8	0	105	57	25	1
BRI > 14	38	30	5	3	0	16	13	9	1

GORD, gastro-oesophageal reflux disease; NUD, non-ulcer dyspepsia.

Acute and chronic inflammation of grades 2 and 3 were combined for all analyses.

NUD versus GORD: acute, Pearson $\chi^2=0.97$, df=2, p=0.62; chronic, Pearson $\chi^2=3.08$, df=2, p=0.21.

Hp+ versus *Hp*-: acute, Pearson $\chi^2=114.92$, df=2, p<0.0001; chronic, Pearson $\chi^2=94.95$, df=2, p<0.0001.

BRI ≤ 14 versus BRI > 14: acute, Pearson $\chi^2=0.98$, df=2, p=0.61; chronic, Pearson $\chi^2=4.72$, df=2, p=0.09.

Table 2 Association between intestinal metaplasia (IM) at the cardia and sex, increased bile reflux, *Helicobacter pylori* status, clinical group, and presence of acute and chronic inflammation at the cardia

Variable	Total No	% IM at cardia	Unadjusted odds ratio	95% CI	Adjusted odds ratio*	95% CI	p Value
Sex							
Female	130	24.6	1.00		1.00		
Male	96	35.4	1.68	0.94–2.99	2.67	1.36–5.27	0.004
BRI							
≤14	188	23.9	1.00		1.00		
>14	38	55.3	3.92	1.91–8.08	2.62	1.17–5.86	0.019
<i>H. pylori</i>							
Negative	182	25.8	1.00		1.00		
Positive	44	43.2	2.18	1.10–4.32	1.03	0.45–2.36	0.95
Group							
GORD	149	28.2	1.00		1.00		
NUD	77	31.2	1.15	0.63–2.10	1.06	0.53–2.09	0.87
Cardia CI							
No	119	18.5	1.00		1.00		
Yes	107	41.1	3.08	1.69–5.62	2.87	1.37–6.02	0.005
Cardia AI					N/A	N/A	N/A
No	181	24.3	1.00				
Yes	45	48.9	2.98	1.52–5.86			

*Factors mutually adjusted for each other and age in a logistic regression model. Age was statistically significant in the model (adjusted odds ratio=1.05/year (95% confidence interval (CI) 1.02–1.07); p=0.0007).

N/A, not applicable as acute and chronic inflammation at the cardia exhibited collinearity. Including chronic inflammation at the cardia resulted in the model with the most explanatory power.

GORD, gastro-oesophageal reflux disease; NUD, non-ulcer dyspepsia; CI, chronic inflammation; AI, acute inflammation.

Histological findings in the antrum were used to calculate the BRI. BRI was originally derived by stepwise logistic regression analysis of the histological grades found in antral biopsies from 350 subjects in whom gastric juice bile acid levels had been measured. Following analysis, an index comprising $(7 \times \text{Oed}) + (3 \times \text{IM}) + (4 \times \text{CI}) - (6 \times \text{Hp})$ gave the best prediction of a raised gastric juice bile acid concentration. An index above 14 had a sensitivity of 70% and a specificity of 85% for a bile acid level >1.00 mmol/l (the upper limit of “physiological” reflux).¹¹

Statistical analysis

The effect of clinical group (GORD or NUD), *H. pylori* status, and increased bile reflux on acute and chronic inflammation at the cardia was assessed using the Pearson χ^2 test. The association between IM at the cardia and sex, *H. pylori* status, increased bile reflux, clinical group, and the presence of acute and chronic inflammation at the cardia were evaluated using Fisher's exact test. The association between these factors and type of IM was assessed using Fisher's exact test while the association between age and IM was assessed using the Student's independent *t* test. Independent risk factors for IM at the cardia were evaluated using logistic regression with cardia IM as the independent variable. The proportion of the variance explained by the logistic regression model was calculated using Nagelkerke r^2 . A *p* value <0.05 was considered statistically significant for all analyses, and calculations were performed using SPSS version 9.0 (SPSS Inc. Chicago, Illinois, USA).

RESULTS

In 39 of 267 patients biopsied, no cardia-type mucosa was identified, and in a further two cases the antral biopsies were inadequate or missing. Thus there were 226 patients with cardia mucosa available (range 1–4 biopsies, median 2, mean 1.98) and at least one adequate biopsy of antral mucosa (see fig 1). Of these, 149 patients were considered to have GORD and 77 had NUD. The gastric biopsy findings from 47 of these NUD patients who were *H. pylori* negative were used in the control group of a previous study on bile reflux gastritis.¹⁴ With

regard to *H. pylori* status, 182 of 226 patients were negative and 44 (19.5%) were positive. There were no cases where the corpus biopsies were positive and the antral biopsies were negative for helicobacters. Concerning the BRI, 188/226 patients had a BRI ≤14 while 38 had a BRI >14. While inflammation at the cardia showed no significant differences between subjects with GORD or NUD, or with an elevated BRI, *H. pylori* positive cases were highly significantly different from negative patients for both acute and chronic inflammation (table 1).

IM was found at the cardia in 66 patients (29%); in 28 the cardia IM was predominantly (if not exclusively) type I, in 25 it was type II, and in 13 it was type III.

Unadjusted analyses revealed that *H. pylori* infection, BRI >14, and acute and chronic inflammation at the cardia were associated with IM (table 2). There was a trend for male sex to be associated with cardia IM but this did not reach statistical significance, while there was no difference between GORD and NUD patients (table 2). Patients with IM at the cardia were older (median age 57 years) than patients without IM (median 47 years) and the difference in mean age was 9.7 years (95% confidence interval (CI) 5.6–13.7). *H. pylori* infection and BRI >14 were also associated with increasing age and it is possible that these factors are acting as markers for older patients. To evaluate independent risk factors for IM at the cardia, age, sex, clinical group, BRI ≤14 or >14, *H. pylori* status, and chronic inflammation at the cardia were entered into a logistic regression model. Acute inflammation was omitted from the model as this exhibited strong collinearity with chronic inflammation. Age, sex, BRI >14, and chronic inflammation remained independent risk factors for cardia IM but *H. pylori* status became non-significant (table 2). The model explained 25.5% of the variance of the data. *H. pylori* status was strongly associated with chronic inflammation and the lack of a statistically significant relationship between *H. pylori* infection and cardia IM could be due to collinearity. This is not the case however as *H. pylori* was not associated with cardia IM in a logistic model that only included age (odds ratio 1.7; 95% CI 0.8–3.4; *p*=0.17).

The influence of *H. pylori* status, BRI >14, sex, clinical group, and chronic and acute inflammation on the type of cardia IM

Table 3 Association between incomplete intestinal metaplasia (types II and III IM), sex, bile reflux index (BRI), *Helicobacter pylori* status, clinical group, and chronic and acute inflammation at the cardia

Variable	Total No	% with incomplete IM	Unadjusted odds ratio	95% CI	p Value*
Sex					
Female	32	68.8	1.00		
Male	34	47.1	0.40	0.15–1.11	0.087
BRI					
≤14	45	51.1	1.00		
>14	21	71.4	2.39	0.79–7.28	0.18
<i>H. pylori</i>					
Negative	47	63.8	1.00		
Positive	19	42.1	0.41	0.14–1.22	0.17
Group					
GORD	42	57.1	1.00		
NUD	24	58.3	1.05	0.38–2.90	1.00
Cardia CI					
No	22	63.6	1.00		
Yes	44	54.5	0.69	0.24–1.96	0.60
Cardia AI					
No	44	65.9	1.00		
Yes	22	40.9	0.36	0.12–1.03	0.067

*Fisher's exact test

GORD, gastro-oesophageal reflux disease; NUD, non-ulcer dyspepsia; CI, chronic inflammation; AI, acute inflammation.

(complete (type I) or incomplete (types II and III)) was evaluated (table 3). There was a trend (>20% difference) for incomplete metaplasia to be associated with a BRI >14, *H. pylori* negative status, absence of acute inflammation at the cardia, and female sex. However, these factors did not reach formal statistical significance and the numbers of cases are small, making the data difficult to interpret (table 3).

Antral IM was found in 26 of 66 patients with cardia IM (39%) but there were an additional 16 patients with isolated antral IM. While the score for antral IM contributed to BRI, its presence did not predict high values. Equal numbers of patients with metaplasia in the stomach (that is, 21/42) had a BRI above and below 14.

DISCUSSION

The most striking finding in this study was that a histological index of bile reflux into the stomach was significantly associated with the presence of IM at the cardia. This is in keeping with a previous study in which we found that patients with Barrett's oesophagus (that is, IM in the oesophagus) had significantly higher BRI values than those with uncomplicated reflux oesophagitis or simple NUD.¹⁴ Interestingly, we found no statistically independent association between cardia IM and symptoms of GORD or with *H. pylori* infection. While it is evident that a lack of statistical significance does not equate to absence of effect, our finding that increasing age, male sex, and chronic inflammation, but not *H. pylori* infection, were factors most closely associated with cardia IM is clearly in keeping with the known epidemiology of cardia cancer. The additional observation that a high BRI is also a significant risk factor for cardia IM gives important new information which may have a bearing on pathogenesis.

We identified chronic inflammation at the cardia in 106 of 226 (47%) patients yet "carditis" has been claimed to be a universal finding whenever cardia-type mucosa is present and is indicative of GORD.^{15,16} While it is conceded that small numbers of lymphocytes and plasma cells are frequently, if not universally, present in cardia mucosa, this does not necessarily represent "chronic inflammation". The mere presence of mononuclear cells cannot be equated with inflammation as otherwise the entire mucosa of the intestinal tract would be in a state of inflammation. It seems to us that some degree of

lymphocytic infiltration, especially close to the squamocolumnar junction, is likely to be "physiological". As Weinstein puts it¹⁷ "these changes (carditis) could represent the consequence of gastric contents lapping at the shores of the oesophagus *in health*" (italics added). If it is acknowledged that the cardia has a "normal" complement of mononuclear cells, then the diagnosis of chronic inflammation requires a higher threshold, and this has been our approach. Some authors avoid the difficulties in diagnosing chronic inflammation at this site by defining "carditis" as infiltration by polymorphs.¹⁸ Using this criterion, equivalent to *active* chronic inflammation, we found "carditis" in only 45/226 cases (19.9%). Interestingly this is very close to the 18.4% prevalence of polymorph infiltration found by Der and colleagues¹⁶ yet these authors diagnosed chronic inflammation in all cases where cardia mucosa was present. In our series, acute and chronic inflammation at the cardia were significantly greater in *H. pylori* infected compared with non-infected patients but inflammation bore no significant association to clinical group. These findings run contrary to recent studies that emphasise the role of GORD^{3,9} but are entirely analogous to those reported by others.^{8,19,20}

With regard to the subtypes of IM found in cardia mucosa, we found a trend towards an association between incomplete IM and a high BRI, *H. pylori* negativity, absence of acute inflammation, and female sex but the groups were too small to yield significant associations. Previous work has examined IM subtypes at the cardia and their disease associations and found significant differences. For instance, Voutilainen and colleagues⁶ examined 235 patients with cardia IM and found that complete IM was associated with a pattern of gastritis characteristic of longstanding *H. pylori* infection while incomplete IM was associated with erosive oesophagitis, the lesion which predisposes to Barrett's oesophagus.

To our knowledge this is the first study that has demonstrated, albeit by indirect means, a link between bile reflux and cardia IM. This has potentially important implications for the development of cancer at the cardia. The connection between exposure to bile acids or their breakdown products and carcinogenesis at other sites is well known. The development of duodenal adenomas and carcinomas in patients with familial adenomatous polyposis and the predilection for adenocarcinoma to arise at the ampulla of Vater are thought to be bile related.²¹ The appearance of

dysplasia and adenocarcinoma in gastric remnants following partial gastrectomy, and the development of carcinoma in the stomachs of experimental animals subjected to enterogastric reflux, have also been attributed to exposure to bile and/or duodenopancreatic juice.²² Finally, there is evidence that faecal bile acids in conjunction with dehydrogenating bacteria are implicated in the development of sporadic colorectal cancer.²³ Thus bile acids have the potential not only to bring about IM at the cardia but also to generate the carcinogens which act on this altered mucosa to produce neoplasia.²⁴ Changes in pH and the composition of the refluxate after regurgitation through the stomach could alter its carcinogenic potential²⁵ and render it more damaging to the oesophagus.^{26, 27} Given the intrinsic instability of junctional sites and their predisposition to neoplasia,²⁸ it is certainly plausible that exposure of the cardia to duodenogastric reflux could contribute to the development of both IM and adenocarcinoma at this site. Nevertheless, our conclusions must remain speculative until confirmed by direct measurements of intragastric bile content.

ACKNOWLEDGEMENTS

We are grateful to Knoll Ltd for an unrestricted grant in support of Dr Neville during the conduct of this study. Thanks are due to Stephen Ryan and Peter Jackson for technical assistance, Tim Branch for pathology computer searches, and to Les Beck for slide retrieval. Paul Moayyedi is funded by a UK MRC Health Services Research Training Fellowship.

Authors' affiliations

M F Dixon, N P Mapstone, Academic Unit of Pathology, University of Leeds, UK

P M Neville, P Moayyedi, A T R Axon, Centre for Digestive Diseases, General Infirmary at Leeds, Leeds, UK

REFERENCES

- Owen DA. Stomach. In: Sternberg SS ed. *Histology for pathologists*, 2nd edn. Philadelphia: Raven Lippincott, 1997:481.
- Fenoglio-Preiser CM, Noffsinger AE, Stemmerman GN, et al. *Gastrointestinal pathology. An atlas and text*, 2nd edn. Philadelphia: Raven Lippincott, 1999:133.
- Chandrasoma PT, Lokuheetty DM, Demeester TR, et al. Definition of histopathologic changes in gastroesophageal reflux disease. *Am J Surg Pathol* 2000;**24**:344–51.
- Carton E, Caldwell MTP, McDonald G, et al. Specialized intestinal metaplasia in patients with gastro-oesophageal reflux disease. *Br J Surg* 2000;**87**:116–21.
- Spechler SJ, Zeroogian JM, Antonioli DA, et al. Prevalence of metaplasia at the gastro-oesophageal junction. *Lancet* 1994;**344**:1533–6.
- Voutilainen M, Farkkila M, Juhola M, et al. Complete and incomplete intestinal metaplasia at the oesophagogastric junction: prevalences and associations with endoscopic erosive oesophagitis and gastritis. *Gut* 1999;**45**:644–8.
- Ruol A, Parenti A, Zaninotto G, et al. Intestinal metaplasia is the probable precursor of adenocarcinoma in Barrett oesophagus and adenocarcinoma of the gastric cardia. *Cancer* 2000;**88**:2320–8.
- Goldblum JR, Vicari JJ, Falk GW, et al. Inflammation and intestinal metaplasia of the gastric cardia: The role of gastroesophageal reflux and *H. pylori* infection. *Gastroenterology* 1998;**114**:633–9.
- Oberg S, Peters JH, Demeester TR, et al. Inflammation and specialized intestinal metaplasia of cardiac mucosa is a manifestation of gastroesophageal reflux disease. *Ann Surg* 1997;**226**:522–30.
- Marshall EEK, Anggiansah A, Owen WJ. Bile in the oesophagus: clinical relevance and ambulatory detection. *Br J Surg* 1997;**84**:21–8.
- Sobala GM, O'Connor HJ, Dewar EP, et al. Bile reflux and intestinal metaplasia in gastric mucosa. *J Clin Pathol* 1993;**46**:235–40.
- Dixon MF, Genta RM, Yardley JH, et al. Classification and grading of gastritis: The updated Sydney system. *Am J Surg Pathol* 1996;**20**:1161–81.
- Shah KA, Deacon AJ, Dunscombe P, et al. Intestinal metaplasia subtyping: evaluation of Gomori's aldehyde fuchsin for routine diagnostic use. *Histopathology* 1997;**31**:277–83.
- Dixon MF, Neville PM, Mapstone NP, et al. Bile reflux gastritis and Barrett's oesophagus: further evidence of a role for duodenogastro-oesophageal reflux? *Gut* 2001;**49**:359–63.
- Rotterdam H. Pathology of the gastric cardia. *Verh Dtsch Ges Pathol* 1999;**83**:37–42.
- Der R, Tsao-Wei DD, Demeester T, et al. Carditis: A manifestation of gastroesophageal reflux disease. *Am J Surg Pathol* 2001;**25**:245–52.
- Weinstein WM, Bogoch ER, Bowes KL. The normal human esophageal mucosa: a histological reappraisal. *Gastroenterology* 1975;**68**:40–4.
- El-Zimaity HMT, Verghese VJ, Ramchatesingh J, et al. The gastric cardia in gastro-oesophageal disease. *J Clin Pathol* 2000;**53**:619–25.
- Chen YY, Antonioli DA, Spechler SJ, et al. Gastroesophageal reflux disease versus *Helicobacter pylori* infection as the cause of gastric carditis. *Mod Pathol* 1998;**11**:950–6.
- El-Serag HB, Sonnenberg A, Jamal MM, et al. Characteristics of intestinal metaplasia in the gastric cardia. *Am J Gastroenterol* 1999;**94**:622–7.
- Spigelman AD, Williams CB, Grenowska M, et al. Gastro-duodenal adenomas in polyposis—Is bile the missing link? *Gut* 1989;**30**:A710–11.
- Miwa K, Hattori T, Miyazaki I. Duodenogastric reflux and foregut carcinogenesis. *Cancer* 1995;**75**:1426–32.
- Nagengast FM. Bile acids and colon carcinogenesis. *Scand J Gastroenterol* 1988;**23**(suppl 154):76–81.
- Busby WF, Shuker DEG, Charnley G, et al. Carcinogenicity in rats of the nitrosated bile acid conjugates N-nitrosoglycocholic acid and N-nitrosotaurocholic acid. *Cancer Res* 1985;**45**:1367–71.
- Hofmann AF, Mysels KJ. Bile acid solubility and precipitation in vitro and in vivo: the role of conjugation, pH and Ca²⁺ ions. *J Lipid Res* 1992;**33**:617–26.
- Kauer WKH, Peters JH, DeMeester TR, et al. Mixed reflux of gastric and duodenal juices is more harmful to the esophagus than gastric juice alone. The need for surgical therapy re-emphasized. *Ann Surg* 1995;**222**:525–33.
- Nehra D, Howell P, Williams CP, et al. Toxic bile acids in gastro-oesophageal reflux disease: influence of gastric acidity. *Gut* 1999;**44**:598–602.
- van Zanten SJOV, Dixon MF, Lee A. The gastric transitional zones: Neglected links between gastroduodenal pathology and *Helicobacter* ecology. *Gastroenterology* 1999;**116**:1217–29.