LEADING ARTICLE

Liver tolerance mediated by antigen presenting cells: fact or fiction?

A H Lau, A de Creus, L Lu, A W Thomson

Gut 2003;52:1075–1078

Possible mechanisms by which liver antigen presenting cells (APC) may facilitate tolerance induction are discussed. Tolerance may be facilitated by a distinctive combination of factors, linked to the unique anatomical and microenvironmental features of the liver.

It is generally accepted that following organ transplantation, rejection mediated by alloreactive T cells will occur if donor and recipient are mismatched for so-called transplantation antigens (Ag), especially those encoded by the major histocompatibility complex (MHC). These rules do not always apply to liver transplantation as hepatic allografts may be accepted “spontaneously” in inbred mice, rats, and outbred pigs without the need for, or following withdrawal of, immunosuppression, even across full MHC barriers. Moreover, spontaneous acceptance of experimental liver grafts is accompanied by donor specific tolerance of other transplanted organs, such as the kidney or heart. The concept of liver “tolerogenicity” in humans is supported by evidence that hepatic allografts have a protective effect on the survival of other organ transplants performed at the same time¹ and that, unlike with other organs, MHC matching is not a prerequisite for successful transplant outcome. Furthermore, safe and lasting withdrawal of a proportion of stable liver transplant recipients from all anti-rejection therapy has been achieved under physician directed protocols.^{2,3}

The tolerogenic properties of the liver may have evolved in response to the need for immunological unresponsiveness to orally acquired Ag presented by Ag presenting cells (APC) and may explain the persistence of some viral infections within the liver as well as the survival advantage of liver allografts. Mechanisms proposed to account for the tolerogenic properties of the liver have not yet been validated convincingly, and indeed its inherent tolerogenicity may be due to a combination of several biological properties that distinguish the liver from other parenchymal organs. Thus in addition to its unique structure and vascular architecture that facilitates interaction between liver parenchymal and non-parenchymal cells and circulating immune cells, the liver also exhibits extensive regenerative and haematopoietic potential. An association between the long term persistence of donor derived haematopoietic cells and induction of tolerance has been demonstrated in liver graft recipients.⁴ Moreover, it has been speculated that donor derived dendritic cells (DC) generated from

precursors present in the liver may play a key role in the subversion of T cell mediated immunity and the long term survival of liver allografts.⁵

The diverse constituency of hepatic professional and non-professional APC, that include liver sinusoidal endothelial cells (LSEC), liver macrophages (Kupffer cells; KC), hepatocytes, and DC, may play key roles in regulating immune responses and facilitating tolerance induction. The unique architecture of the liver allows T cells to interact with resident liver APC, making these latter cells, particularly the uniquely well equipped DC, ideal candidates for directing immune responses towards immunity or tolerance. We speculate that within the liver microenvironment, production of anti-inflammatory cytokines and other molecules including growth factors, such as interleukin (IL)-10, transforming growth factor (TGF)- β , prostaglandin (PG)E₂, and granulocyte macrophage-colony stimulating factor results in the modulation of APC differentiation, trafficking, and function in both health and disease. Much has recently been discussed about the potential of DC to regulate immune responses and to promote tolerance induction. The aforementioned soluble factors may each play a role in “fashioning” intrahepatic tolerogenic DC and other APC.

Due to their capacity to express death ligands, DC and other APC may also contribute to the extensive apoptosis of systemically activated T cells within the liver, a phenomenon that may underlie inhibition of anti-donor responses following hepatic transplantation. The number of T cells that undergo apoptosis within the early inflammatory infiltrate of murine liver allografts is increased significantly compared with grafts that are rejected acutely.⁶ Both activation induced cell death and passive cell death may be involved in T cell apoptosis that predisposes to tolerance induction. Although CD8⁺ T cell apoptosis has been shown to be Fas-FasL (CD95-CD95L) pathway independent, alternative death receptor pathways have not been excluded. Apoptosing T cells release TGF- β and IL-10,⁷ further contributing to the anti-inflammatory microenvironment of the liver. Immune reactivity may also be

See end of article for authors' affiliations

Correspondence to:
Dr A Thomson,
Departments of Surgery
and Immunology,
University of Pittsburgh
Medical Center, W1544
Biomedical Science Tower,
200 Lothrop St, Pittsburgh,
PA 15213, USA;
thomsonaw@msx.upmc.edu

Accepted for publication
13 May 2003

Abbreviations: Ag, antigen; MHC, major histocompatibility complex; APC, antigen presenting cells; DC, dendritic cells; LSEC, liver sinusoidal endothelial cells; KC, Kupffer cells; IL, interleukin; TGF, transforming growth factor; PG, prostaglandin; NK, natural killer; TNF, tumour necrosis factor; Th, T helper; Treg, regulatory T cells; IFN, interferon; CTLA4lg, cytotoxic T lymphocyte Ag 4 immunoglobulin.

affected by interactions between DC and natural killer (NK) cells, NK T cells, and $\gamma\delta$ T cells that are found in unusually high proportions within the liver. Hereon, we will highlight and discuss possible mechanisms by which liver APC may facilitate tolerance induction.

“Among the different cell populations thought to be involved in hepatic tolerance, LSEC are also important because they are highly efficient sessile APC with immunoregulatory functions”

As the liver surveys a massive blood supply, mainly first pass blood from the gut, it is, compared with other organs, constantly exposed to a vast quantity and diversity of Ag. Among the different cell populations thought to be involved in hepatic tolerance, LSEC are also important because they are highly efficient sessile APC⁸ with immunoregulatory functions. Unlike vascular endothelial cells from other organs, such as skin, gut, or lung, which cannot function as APC for CD4⁺ and CD8⁺ T cells unless pre-exposed to proinflammatory cytokines, LSEC can cross present exogenous Ag⁹ and induce Ag specific tolerance.¹⁰ The consequence of cross presentation of Ag by LSEC in vivo is induction of systemic immune tolerance through clonal deletion as well as induction of T cell anergy.¹⁰

LSEC express surface molecules necessary for efficient Ag uptake by receptor mediated endocytosis, interaction with leucocytes, and efficient Ag presentation to T cells. In addition, LSEC can express apoptosis inducing molecules (FasL, tumour necrosis factor (TNF) receptor apoptosis inducing ligand (TRAIL), and membrane TNF- α) that may contribute to intra-hepatic T cell death.¹¹ In vitro, mouse LSEC can prime T cells efficiently but the tendency is towards tolerance induction. Thus murine CD8⁺ T cells activated by LSEC are functionally inert, lack cytotoxic effector function, exhibit poor cytokine release, and eventually apoptose¹⁰ whereas CD4⁺ T cells differentiate into IL-4 secreting T helper (Th)2 cells or IL-10 secreting regulatory T cells (Treg).¹² Furthermore, in the presence of IL-10 or PGE₂, murine LSEC downregulate MHC class II and costimulatory molecule expression and decrease mannose receptor mediated Ag uptake, which results in diminished T cell activation.¹³ Human LSEC, on the other hand, do not normally express MHC class II or costimulatory molecules and it is reasonable to assume that T cells are not activated by these APC under non-inflammatory conditions.¹⁴ Not only are IL-10 and PGE₂ important in promoting the tolerogenic properties of LSEC, but they are also expressed constitutively by these cells and upregulated upon stress.^{13, 15} Thus modulation of T cell activation by LSEC is a likely reflection of the influence of IL-10 or PGE₂ on the Ag presenting capacity of these cells.

“Modulation of T cell activation by LSEC is a likely reflection of the influence of IL-10 or PGE₂ on the Ag presenting capacity of these cells”

KC, the resident liver macrophages, phagocytose microorganisms efficiently and initiate T cell responses. In addition, like DC, they may also play a role in the induction of tolerance via mechanisms such as cross presentation of Ag derived from ingested apoptotic cells¹⁶ to either resident DC^{17, 18} or DC traversing the liver. Classical macrophage activation is induced by interferon (IFN)- γ and by microbial triggers such as bacterial lipopolysaccharide, and results in secretion of proinflammatory cytokines (TNF, IL-1, IL-6, IL-12), nitric oxide, and upregulation of MHC class II molecules.¹⁹ However, KC may also be deactivated under the influence of factors within the liver microenvironment. Indeed, it has been shown for other types of macrophages, such as those derived from human blood mononuclear cells,²⁰ that anti-inflammatory cytokines

such as IL-10 and TGF- β inhibit their ability to secrete proinflammatory factors (IL-12, IFN- γ), downregulate MHC class II expression, induce PGE₂ production, and promote paracrine and autocrine secretion of IL-10 and TGF- β .¹⁹ Furthermore, KC appear to play a crucial role in tolerance induction induced by portal venous administration of soluble alloAg.²¹

“DC are the best equipped and, when mature, the most potent APC”

DC are the best equipped and, when mature, the most potent APC. There is now considerable evidence for their phenotypic heterogeneity and functional diversity. Their activities are modulated by a range of stimuli, including pathogen derived products and inflammatory mediators. Depending on Ag dose, their maturation state, and environmental signals, DC can direct either Th1- or Th2-type immune responses. They are rendered tolerogenic by exposure to immunosuppressive cytokines (IL-10 and TGF- β) that inhibit production of the Th1 driving cytokine IL-12 and suppress costimulatory molecule expression. Blocking of costimulatory molecules on the surface of DC^{22, 23} (for example, by cytotoxic T lymphocyte Ag 4 immunoglobulin (CTLA4Ig)) inhibits their T cell stimulatory ability and promotes death of alloreactive T cells.²⁴ Induction of Treg cells by immature DC is also a possible mechanism that may contribute to induction of tolerance. It has been shown that the presence of Treg cells within allografts is important in transplant tolerance.²⁵ Activation of Treg cells is believed to downregulate MHC class II and costimulatory molecules on DC, maintaining them in an immature state.²⁶ In addition, repetitive interaction of allogeneic T cells with immature DC leads to expression of CTLA4, IL-10 production, and induction of a Treg phenotype.²⁷ In humans, injection of immature autologous DC pulsed with antigenic peptide induces Ag specific tolerance in CD8⁺ T cells.²⁸

“An important factor contributing to a role of liver APC in the promotion of tolerance is the unique hepatic microenvironment that is rich in IL-10 and TGF- β ”

We contend that an important factor contributing to a role of liver APC in the promotion of tolerance is the unique hepatic microenvironment that is rich in IL-10 and TGF- β (synthesised by hepatocytes, LSEC, and other liver cell types, including T cells). In particular, this environment may sustain or promote an immature/tolerogenic phenotype in resident DC. Indeed, it has been shown that freshly isolated hepatic DC are poor stimulators of naïve allogeneic T cells and, when administered in vivo, promote IL-10 production selectively in secondary lymphoid tissue.²⁹ We postulate that potentially tolerogenic APC of donor or recipient origin (the latter trafficking into the graft) may play a key role in the acceptance of liver allografts by inducing Treg cells and/or T cell apoptosis. Indeed, the unique microenvironment of the liver may provide an ideal site for the immature (tolerogenic) DC-T cell inhibitory feedback loop described recently by Min and colleagues.³⁰

In addition to their maturation status, the nature of the DC subsets present in the liver may contribute to its potential tolerogenicity. Mouse liver contains several DC subsets, including myeloid (CD8 α), lymphoid related (CD8 α ⁺), and the recently described plasmacytoid (B220⁺) DC.^{31, 32} CD8 α ⁺ DC have been implicated in the induction of central and peripheral tolerance. They express FasL, induce apoptosis in alloactivated CD4⁺ T cells, and inhibit IL-2 production by CD8⁺ T cells.^{33, 34} In addition, they prolong heart allograft survival independent of their maturation status.³⁵ Another subset of DC-like cells that are generated by the culture of hepatic non-parenchymal cells

with IL-3 and CD40L may be involved in promoting T cell apoptosis. These DEC205⁺CD11c⁺B220⁺CD19⁻ cells have been shown to first activate and then to induce T cell apoptosis.³⁶

A possible role of plasmacytoid DC, identified recently in murine liver (Lau AH *et al*, unpublished data), in immune regulation has not yet been addressed experimentally. Previous data suggest that these DC may have a regulatory function in their immature state in which they can promote CD4⁺ T cell anergy.³⁷ Furthermore, human plasmacytoid DC appear to inhibit T cell proliferation by secretion of the tryptophan catabolising enzyme indoleamine 2,3-dioxygenase³⁸ and CD40 ligand activated human plasmacytoid DC generate IL-10 producing CD8 Treg cells.³⁹ These data suggest a possible role for liver plasmacytoid DC in liver tolerance.³⁹

A further consideration is the influence of the liver microenvironment on trafficking of APC. BM derived DC cultured in the presence of IL-10 upregulate expression of message for the chemokine receptor CCR5 while downregulating CCR7 mRNA. This corresponds with impaired homing ability of the DC to secondary lymphoid tissue.^{40, 41} These data suggest that, in the presence of IL-10, liver DC could have reduced or defective ability to migrate to secondary lymphoid tissue. Such impaired migration could undermine induction of anti-donor responsiveness due to a reduced ability of the DC to reach T cell areas of the draining lymphoid tissue.

Following liver transplantation, massive Ag release occurs as a result of physical and ischaemic tissue injury, hepatocyte production of donor MHC class I Ag,⁴² and the migration of donor haematopoietic cells into the host. This extensive Ag release likely results in the binding of soluble donor Ag by host T cells—a fact that may block their recognition of target cells, suppress anti-donor reactivity, and promote tolerance induction.

“Tolerance may be facilitated by a distinctive combination of factors, linked to the unique anatomical features of the liver”

Mechanisms by which the liver may foster immunological tolerance under normal steady state conditions and in the context of liver transplantation can be inferred from its unique constituency of APC and their presence in a cytokine microenvironment conducive to their tolerogenic function and potential to subvert T cell responses. Tolerance may be facilitated by a distinctive combination of factors, linked to the unique anatomical features of the liver. Conventional and experimental immunosuppressive agents that inhibit both APC and T cell activation and function are likely to contribute significantly to hepatic tolerogenicity.

ACKNOWLEDGEMENT

The authors' work is supported by National Institutes of Health grants DK49745, AI41011, and AI/DK51698.

Authors' affiliations

A H Lau, A W Thomson, Department of Surgery and Department of Immunology, Thomas E Starzl Transplantation Institute, University of Pittsburgh Medical Center, Pittsburgh, PA 15213, USA

A de Creus, L Lu, Department of Surgery, Thomas E Starzl Transplantation Institute, University of Pittsburgh Medical Center, Pittsburgh, PA 15213, USA

REFERENCES

- 1 **Rasmussen A**, Davies H, Jamieson N, *et al*. Combined transplantation of liver and kidney from the same donor protects the kidney from rejection and improves kidney graft survival. *Transplantation* 1995;**59**:919–21.

- 2 **Ramos H**, Reyes J, Abu-Elmagd K, *et al*. Weaning of immunosuppression of in long-term liver transplant recipients. *Transplantation* 1995;**59**:212–17.
- 3 **Mazariegos G**, Reyes J, Marino I, *et al*. Weaning of immunosuppression in liver transplant recipients. *Transplantation* 1997;**63**:243–39.
- 4 **Lu L**, Rudert WA, Qian S, *et al*. Growth of donor-derived dendritic cells from the bone marrow of murine liver allograft recipients in response to granulocyte/macrophage colony-stimulating factor. *J Exp Med* 1995;**182**:379–87.
- 5 **Thomson AW**, Lu L. Are dendritic cells the key to liver transplant tolerance? *Immunol Today* 1999;**20**:27–32.
- 6 **Sharland A**, Yan Y, Wang C, *et al*. Evidence that apoptosis of activated T cells occurs in spontaneous tolerance of liver allografts and is blocked by manipulations which break tolerance. *Transplantation* 1999;**68**:1736–45.
- 7 **Silleit H**, Cruickshank S, Southgate J, *et al*. Transforming growth factor- β promotes 'death by neglect' in post-activated human T cells. *Immunology* 2001;**102**:310–16.
- 8 **Knolle PA**, Gerken G. Local control of the immune response in the liver. *Immunol Rev* 2000;**174**:21–34.
- 9 **Scoazec J**, Feldmann G. In situ immunophenotyping study of endothelial cells of the human hepatic sinusoid: results and functional implications. *Hepatology* 1991;**14**:789–97.
- 10 **Limmer A**, Ohl J, Kurts C, *et al*. Efficient presentation of exogenous antigen by liver endothelial cells to CD8⁺ T cells results in antigen-specific T-cell tolerance. *Nat Med* 2000;**6**:1348–54.
- 11 **Muschen M**, Warskulat U, Douillard P, *et al*. Regulation of CD95 (APO-1/Fas) receptor and ligand expression by lipopolysaccharide and dexamethasone in parenchymal and nonparenchymal rat liver cells. *Hepatology* 1998;**27**:200–8.
- 12 **Knolle PA**, Schmitt E, Jin S, *et al*. Induction of cytokine production in naive CD4⁺ T cells by antigen-presenting murine liver sinusoidal endothelial cells but failure to induce differentiation towards Th1 cells. *Gastroenterology* 1999;**116**:1428–40.
- 13 **Knolle PA**, Uhrig S, Hegenbarth S, *et al*. IL-10 down-regulates T cell activation by antigen-presenting liver sinusoidal endothelial cells through decreased antigen uptake via the mannose receptor and lowered surface expression of accessory molecules. *Clin Exp Immunol* 1989;**114**:427–33.
- 14 **Leifeld L**, Trautwein C, Dumoulin F, *et al*. Enhanced expression of CD80 (B7-1), CD86 (B7-2), and CD40 and their ligands CD28 and CD154 in fulminant hepatic failure. *Am J Pathol* 1999;**154**:1711–20.
- 15 **Lau A**, Thomson AW. Dendritic cells and immune regulation in the liver. *Gut* 2003;**52**:307–14.
- 16 **Ruzittu M**, Carla E, Montinari M, *et al*. Modulation of cell surface expression of liver carbohydrate receptors during in vivo induction of apoptosis with lead nitrate. *Cell Tissue Res* 1999;**298**:105–12.
- 17 **Albert M**, Pearce S, Francisco L, *et al*. Immature dendritic cells phagocytose apoptotic cells via α 5 β 1 and CD36, and cross-present antigens to cytotoxic T lymphocytes. *J Exp Med* 1998;**188**:1359–68.
- 18 **Uwatoku R**, Suematsu M, Ezaki T, *et al*. Kupffer cell-mediated recruitment of rat dendritic cells to the liver: roles of N-acetylgalactosamine-specific sugar receptors. *Gastroenterology* 2001;**121**:1460–72.
- 19 **Gordon S**. Alternative activation of macrophages. *Nat Rev Immunol* 2002;**3**:23–35.
- 20 **Fadok VA**, Bratton DL, Konowal A, *et al*. Macrophages that have ingested apoptotic cells in vitro inhibit proinflammatory cytokine production through autocrine/paracrine mechanisms involving TGF- β , PGE₂, and PAF. *J Clin Invest* 1998;**101**:890–8.
- 21 **Kamei T**, Callery M, Flye M. Kupffer cell blockade prevents induction of portal venous tolerance in rat cardiac allograft transplantation. *J Surg Res* 1990;**48**:393–6.
- 22 **Bartlett A**, McCall J, Ameratunga R, *et al*. Costimulatory blockade prevents early rejection, promotes lymphocyte apoptosis, and inhibits the upregulation of intragraft interleukin-6 in an orthotopic liver transplant model in the rat. *Liver Transpl* 2002;**8**:458–68.
- 23 **Li W**, Lu L, Wang L, *et al*. Costimulation blockade promotes the apoptotic death of graft-infiltrating T cells and prolongs survival of hepatic allografts from Flt3L-treated donors. *Transplantation* 2001;**72**:1423–32.
- 24 **Lu L**, Qian S, Hershberger P, *et al*. Fas Ligand (CD95L) and B7 expression on dendritic cells provide counter-regulatory signals for T cell survival and proliferation. *J Immunol* 1997;**158**:5676–84.
- 25 **Graca L**, Cobbold S, Waldmann H. Identification of regulatory T-cells in tolerated allografts. *J Exp Med* 2002;**195**:1641–6.
- 26 **Gimmi C**, Freeman G, Gribben J, *et al*. Human T-cell clonal anergy is induced by antigen presentation in the absence of B7 costimulation. *Proc Natl Acad Sci U S A* 1993;**90**:6586–90.
- 27 **Jonuleit H**, Schmitt E, Schuler G, *et al*. Induction of interleukin 10-producing, nonproliferating CD4⁺ T cells with regulatory properties by repetitive stimulation with allogeneic immature human dendritic cells. *J Exp Med* 2000;**192**:1213–22.
- 28 **Dhodapkar M**, Steinman RM, Krasovsky J, *et al*. Antigen-specific inhibition of effector T cell function in humans after injection of immature dendritic cells. *J Exp Med* 2001;**193**:233–8.
- 29 **Khanna A**, Morelli AE, Zhong C, *et al*. Effects of liver-derived dendritic cell progenitors on Th1- and Th2-like cytokine responses in vitro and in vivo. *J Immunol* 2000;**164**:1346–54.
- 30 **Min W-P**, Zhou D, Ichim T, *et al*. Inhibitory feedback loop between tolerogenic dendritic cells and regulatory T cells in transplant tolerance. *J Immunol* 2003;**170**:1304–12.

- 31 Lian Z-X, Okada T, He X-S, *et al.* Heterogeneity of dendritic cells in the mouse liver: Identification and characterization of four distinct populations. *J Immunol* 2003;**170**:2323–30.
- 32 Thomson AW, O'Connell PJ, Steptoe RJ, *et al.* Immunobiology of liver dendritic cells. *Immunol Cell Bio* 2002;**80**:65–73.
- 33 Kronin V, Winkel K, Suss G, *et al.* A subclass of dendritic cells regulates the response of naive CD8⁺ T cells by limiting their IL-2 production. *J Immunol* 1996;**157**:3819–27.
- 34 Suss G, Shortman K. A subclass of dendritic cells kills CD4⁺ T cells via Fas/Fas- ligand-induced apoptosis. *J Exp Med* 1996;**183**:1789–96.
- 35 O'Connell PJ, Li W, Wang Z, *et al.* Immature and mature CD8 α^+ dendritic cells prolong the survival of vascularized heart allografts. *J Immunol* 2002;**168**:143–54.
- 36 Lu L, Bonham CA, Liang X, *et al.* Liver-derived DEC205⁺B220⁺CD19⁺ dendritic cells regulate T cell responses. *J Immunol* 2001;**166**:7042–52.
- 37 Kuwana M, Kaburaki J, Wright T, *et al.* Induction of antigen-specific human CD4⁺ T cell anergy by peripheral blood DC2 precursors. *Eur J Immunol* 2001;**31**:2547–57.
- 38 Munn D, Sharma M, Lee J, *et al.* Potential regulatory function of human dendritic cells expressing indoleamine 2,3-dioxygenase. *Science* 2002;**297**:1867–70.
- 39 Gilliet M, Liu Y-J. Generation of human CD8 T regulatory cells by CD40 ligand- activated plasmacytoid dendritic cells. *J Exp Med* 2002;**195**:695–704.
- 40 D'Amico G, Frascaroli G, Bianchi G, *et al.* Uncoupling of inflammatory chemokine receptors by IL-10: generation of functional decoys. *Nat Immunol* 2000;**1**:387–91.
- 41 Takayama T, Morelli AE, Onai N, *et al.* Mammalian and viral IL-10 enhance C-C chemokine receptor 5 but down-regulate C-C chemokine receptor 7 expression by myeloid dendritic cells: impact on chemotactic responses and in vivo homing ability. *J Immunol* 2001;**166**:7136–43.
- 42 Sriwatanawongsa V, Davies H, Calne RY. The essential roles of parenchymal tissues and passenger leukocytes in the tolerance induced by liver grafting in rats. *Nat Med* 1995;**1**:428–32.

Clinical Evidence—Call for contributors

Clinical Evidence is a regularly updated evidence based journal available worldwide both as a paper version and on the internet. *Clinical Evidence* needs to recruit a number of new contributors. Contributors are health care professionals or epidemiologists with experience in evidence based medicine and the ability to write in a concise and structured way.

Currently, we are interested in finding contributors with an interest in the following clinical areas:

Altitude sickness; Autism; Basal cell carcinoma; Breast feeding; Burns; Carbon monoxide poisoning; Cervical cancer; Chronic renal failure; Cystic fibrosis; Ectopic pregnancy; Emphysema; Grief/bereavement; Halitosis; Hodgkins disease; Infectious mononucleosis (glandular fever); Jet lag; Kidney stones; Malignant melanoma (metastatic); Mesothelioma; Myeloma; Ovarian cyst; Pancreatitis (acute); Pancreatitis (chronic); Polycystic ovaries; Polymyalgia rheumatica; Post-partum haemorrhage; Pulmonary embolism; Recurrent miscarriage; Repetitive strain injury; Scoliosis; Seasonal affective disorder; Squint; Systemic lupus erythematosus; Testicular cancer; Uterine prolapse; Varicocele; Viral meningitis; Vitiligo

However, we are always looking for others, so do not let this list discourage you.

Being a contributor involves:

- Appraising the results of literature searches (performed by our Information Specialists) to identify high quality evidence for inclusion in the journal.
- Writing to a highly structured template (about 2000–3000 words), using evidence from selected studies, within 6–8 weeks of receiving the literature search results.
- Working with *Clinical Evidence* Editors to ensure that the text meets rigorous epidemiological and style standards.
- Updating the text every eight months to incorporate new evidence.
- Expanding the topic to include new questions once every 12–18 months.

If you would like to become a contributor for *Clinical Evidence* or require more information about what this involves please send your contact details and a copy of your CV, clearly stating the clinical area you are interested in, to Claire Folkes (cfolkes@bmjgroup.com).

Call for peer reviewers

Clinical Evidence also needs to recruit a number of new peer reviewers specifically with an interest in the clinical areas stated above, and also others related to general practice. Peer reviewers are health care professionals or epidemiologists with experience in evidence based medicine. As a peer reviewer you would be asked for your views on the clinical relevance, validity and accessibility of specific topics within the journal, and their usefulness to the intended audience (international generalists and health care professionals, possibly with limited statistical knowledge). Topics are usually 2000–3000 words in length and we would ask you to review between 2–5 topics per year. The peer review process takes place throughout the year, and our turnaround time for each review is ideally 10–14 days.

If you are interested in becoming a peer reviewer for *Clinical Evidence*, please complete the peer review questionnaire at www.clinicalevidence.com or contact Claire Folkes (cfolkes@bmjgroup.com).