INTESTINE

Depletion of intestinal resident macrophages prevents ischaemia reperfusion injury in gut

Y Chen, V C H Lui, N V Rooijen, P K H Tam

Gut 2004;53:1772-1780. doi: 10.1136/gut.2003.034868

See end of article for authors' affiliations

Correspondence to:
Dr P K H Tam, Division of
Paediatric Surgery,
Department of Surgery,
University of Hong Kong
Medical Centre, Queen
Mary Hospital, Pokfulam,
Hong Kong SAR, China;
paultam@hkucc.hku.hk

Revised version received
2 May 2004
Accepted for publication
6 May 2004

The intestinal mucosa, which lines the lumen of the intestine, acts as a defence barrier against pathogen invasion. Ischaemia reperfusion (IR) causes damage to the intestinal mucosa, producing increased gut permeability and bacterial translocation which can result in systemic inflammatory response syndrome and multiple organ failure. IR injury associated mortality remains high despite advances in patient care for IR injury.¹ The cellular and molecular events involved in IR injury are complex, involving the production of reactive oxygen species (ROS), reactive nitrogen species, and polymorphonuclear neutrophils (PMN) infiltration. This is accompanied by the release of other moderators such as nitric oxide, platelet activation factor, and phospholipase A2.² Recruitment of PMN into reperfused sites after ischaemia is critical for IR injury in the heart, lung, and liver.³⁻⁵ Anti-PMN antiserum treatment offers only limited protection against IR injury in the heart.⁶ This suggests that other inflammatory cells are likely to be involved. For example, mast cells have also been suggested to play a role in IR injury,⁷ and anti-complement agent K-76COONa, which prevents mast cell degranulation, has been shown to ameliorate gut IR injury. However, K-76COONa also inhibits other inflammatory cells such as macrophages⁸ and lymphocytes.⁹ Identifying the additional inflammatory cells that contribute to gut IR injury remains a priority for the development of an effective treatment strategy directed towards specific cellular targets.

Macrophages are inflammatory cells, which on activation by bacterial lipopolysaccharide destroy bacteria by the release

Background and aims: The cellular and molecular events involved in ischaemia reperfusion (IR) injury are complex and not fully understood. Previous studies have implicated polymorphonuclear neutrophils (PMN) as major inflammatory cells in IR injury. However, anti-PMN antiserum treatment offers only limited protection, indicating that other inflammatory cells are involved. We have therefore investigated the contribution of resident macrophages in IR injury using an IR gut injury model.

Methods: DA rats were divided into sham operation and IR groups. The superior mesenteric artery was clamped for 30, 45, or 60 minutes (ischaemia) followed by 60 minutes of reperfusion. IR injuries were evaluated by histological staining. Expression of early growth response factor 1 (Egr-1), myeloperoxidase (MPO), and proinflammatory cytokines was analysed by immunohistochemistry, reverse transcription-polymerase chain reaction, and western blotting analysis. The specific role of macrophages in IR gut injury was also evaluated in resident macrophage depleted rats.

Results: Mucosal sloughing and villi destruction were seen in 45/60 minute and 60/60 minute IR guts. PMN infiltration at the damaged mucosal area was undetectable in 45/60 minute and 60/60 minute IR guts. PMN were localised around the capillaries at the base of the crypts in 60/60 minute IR gut. Obvious Egr-1 immunostaining was localised in resident macrophages at the damaged villi before histological appearance of mucosal damage. Furthermore, resident macrophages at the damaged site expressed MPO. Protein levels of the proinflammatory cytokines RANTES and MCP-1 were increased in IR gut. Depletion of resident macrophages by dichloromethylene bisphosphonate significantly reduced mucosal damage in rat guts after IR.

Conclusion: Our findings indicate that resident macrophages play a role in early mucosal damage in IR gut injury. Therefore, macrophages should be treated as a prime target for therapeutic intervention for IR damage.

of cytokines, ROS, induction of myeloperoxidase (MPO) activity, and phagocytosis of bacteria.¹⁰ Acute inflammation induces PMN infiltration and increases MPO activity in situ, causing local tissue damage.¹¹ Elevated MPO activity occurs in an IR injured gut and localises to the ischaemic site shortly after reperfusion and before the appearance of tissue damage.¹² Activated macrophages exhibit high levels of MPO activity.^{13, 14} Activation of macrophages mediates the apoptosis of stromal, parenchymal, and other immune cells.¹⁵ Chloramines, which are catalytic products of MPO, can cause growth arrest and apoptosis in gastric mucosal cells.^{16, 17} However, it is still unclear whether the increased MPO activity at the IR site is produced as a result of infiltrated PMN, activated macrophages in situ, or both.¹⁸ Mucosa damage is an early histological landmark of IR injury to the gut¹⁹ which manifests functionally as increased gut permeability. Macrophages mainly locate at the mucosa and smooth muscle layers of the intestine, and are termed either mucosal or muscularis resident macrophages. Hierholzer *et al* have shown that muscularis resident macrophages release

Abbreviations: IR, ischaemia reperfusion; PMN, polymorphonuclear neutrophils; Egr-1, early growth response factor 1; MPO, myeloperoxidase; RT-PCR, reverse transcription-polymerase chain reaction; ROS, reactive oxygen species; ICAM-1, intercellular adhesion molecule 1; MCP-1, monocyte chemoattractant protein 1; SMA, superior mesenteric artery; Cl₂MBP, dichloromethylene bisphosphonate; PBS, phosphate buffered saline; iNOS, inducible nitric oxide synthase; LFA-1, lymphocyte function associated antigen 1

the proinflammatory cytokine interleukin 6 and directly contribute to decreased gut motility in the IR injured gut.²⁰ However, the role of mucosal resident macrophages in intestinal IR injury has not been investigated.

Early growth response factor 1 gene (*Egr-1*) encodes a transcription factor and controls the development of the central and peripheral nervous systems.²¹ In vitro studies have identified downstream target genes of *Egr-1*. These include genes encoding tumour necrosis factor α , intercellular adhesion molecule 1 (ICAM-1), CD44, platelet derived growth factor A/B chain, transforming growth factor β , and macrophage colony stimulating factor.²² Expression levels of *Egr-1* are upregulated in the heart,²³ lung,²² and kidney²⁵ after IR challenge. These findings indicate that *Egr-1* is involved in IR injury in these organs. The involvement of *Egr-1* in IR injury is substantiated by findings that lung injury is less severe and associated with improved survival after IR in *Egr-1* knockout mice.²² Production of the proinflammatory cytokine monocyte chemoattractant protein 1 (MCP-1) and adhesion molecule ICAM-1 in IR injured lungs is lower in *Egr-1* knockout mice. This probably accounts for the reduced PMN infiltration and mild lung damage seen in lung IR injury in *Egr-1* knockout mice.

We hypothesise that resident macrophages play a key role in the initiation of intestinal IR injury. To test our hypothesis, we investigated the cellular and molecular events involved in early gut mucosal damage from IR. We showed in a rat IR gut injury model: (i) the onset of mucosal damage before PMN infiltration; (ii) activation of mucosa resident macrophages in the IR injured gut; (iii) expression of high levels of MPO and *Egr-1* in activated resident macrophages; and (iv) depletion of intestinal resident macrophages, which prevented IR injury to the gut in rats.

MATERIALS AND METHODS

Rat IR gut model

Six to eight week old DA rats (~250 g) were divided into sham operation and IR groups. Rats were anaesthetised by intraperitoneal injection (45 mg/kg body weight) of pentobarbital sodium solution (Abbott Lab, Illinois, USA). The superior mesenteric artery (SMA) was occluded by clamping for 30 (n = 6), 45 (n = 6), or 60 minutes (n = 6). A change in colour of the occluded intestine from red to pale immediately after clamping indicated successful induction of ischaemia. After induction of ischaemia, clamps were released and guts were reperused for 60 minutes. In the sham operation groups, SMA was separated but no occlusion was applied for 30+60 minutes (n = 6), 45+60 minutes (n = 6), or 60+60 minutes (n = 6). The operation site was covered with cotton gauze soaked with saline. Segments of the jejunum (8–10 cm) of the IR and sham operation rats were collected for RNA, protein, and immunohistochemistry. The study was approved by the ethics committee of the University of Hong Kong.

Macrophages depletion

Liposome encapsulated dichloromethylene bisphosphonate (Cl₂MBP liposome) or phosphate buffered saline (PBS, control liposome)²⁷ was provided by Roche Diagnostics (Mannheim, Germany) and encapsulated in liposomes in Dr Rooijen's laboratory. The Cl₂MBP or control liposome (4 ml) was injected into the intraperitoneal cavity three days before IR. The liver and jejunum were removed and depletion of macrophages in these organs was verified by immunostaining for resident macrophage. Rats were injected with Cl₂MBP liposome (n = 4) or control liposome (n = 4), and subjected to 45 minutes of ischaemia and 60 minutes of reperfusion.

Immunohistochemistry and immunofluorescence staining

Gut specimens were fixed in 4% paraformaldehyde/PBS (pH 7.2) at 4°C for 18 hours, dehydrated in alcohol, cleared in xylene, and embedded in paraffin. Sections (4 µm thickness) of jejunum were mounted onto TESPA (3-aminopropyl-triethoxy-silane; Sigma, St Louis, Missouri, USA) coated microscope glass. Antigen unmasking was performed by heating the sections in 10 mM of citrate buffer (pH 6.2) for 10 minutes at 95°C. Sections were incubated in 3% (v/v) H₂O₂ in methanol to quench endogenous peroxidase activity. Tissue sections were blocked with Dako protein block (Dako A/S, Glostrup, Denmark) and incubated with *Egr-1* (Santa Cruz Biotechnology, California, USA; 1:800), ED1 (Serotec; 1:100), ED2 (Serotec, Kidlington, Oxford, UK; 1:200) ICAM-1 (Santa Cruz Biotechnology; 1:100), or rat PMN (Accurate, New York, USA; 1:10 000) antibodies in 50 mM Tris buffer (pH 7.4) containing 1% bovine serum albumin (USB Corporation, Ohio, USA) at 37°C for 40 minutes or 4°C for 16 hours. ED1 antibody recognises a cytoplasmic antigen in monocytes and infiltrating macrophages, and is used as a pan macrophage marker.²⁸ ED2 antibody, which recognises a membrane bound antigen similar to human CD163, has been commonly used as a specific marker for resident macrophages.²⁸ Signal amplification and detection were performed with the Dako EnVision System, HRP. Sections were counterstained with haematoxylin, dehydrated in ethanol, cleared in xylene, and mounted with DPX mounting medium (BDH Laboratory Supplies, Poole, Dorset, UK). Immunofluorescence staining was performed by incubating FITC and TRITC conjugated secondary antibodies on sections and mounting them in DAPI containing Vectashield mounting medium (Vector Laboratories, California, USA). Immunofluorescent stained sections were analysed with a fluorescence microscope (Nikon Eclipse E600). Three jejunum sections of each rat in different treatment groups were analysed for histology and immunohistochemistry. Rats of the same group showed similar histology and immunostaining pattern. Representative pictures of haematoxylin/eosin or immunostained sections of each group were shown for comparison.

Quantitative analysis of macrophages and PMN in gut

Numbers of ED2 and ED1 immunopositive cells were counted from 10 randomly selected high power (400×) fields of jejunum sections. The ratio of ED2 to ED1 immunopositive cells was tabulated under different treatments. *Egr-1* immunopositive cells were counted and separated into two groups: cells showing cytoplasmic *Egr-1* staining and cells showing nuclear *Egr-1* staining. A minimum of three sections of each rat was examined and 10 randomly selected high power (400×) fields of each gut section were examined. Two observers counted the sections separately, and the results were averaged and tabulated.

Table 1 Primers used for the polymerase chain reactions

Primer	Sequence
<i>Egr-1</i> : sense	5'-CCG GTT GCC TCC CAT CAC CTA-3'
<i>Egr-1</i> : antisense	5'-GAG CGA AGG CTG CTG GGT ACG-3'
MPO: sense	5' CCG CAG GAC AAG TAT CGA A 3'
MPO: antisense	5' ATC TGG TTG CGA ATG GTA ATG 3'
β -actin: sense	5'-TGG AAT CCT GTG GCA TCC ATG AAA C-3'
β -actin: antisense	5'-ACC ATC AAG GCA CAG CAA CTC-3'

Egr-1, early growth response factor 1; MPO, myeloperoxidase.

Quantitative analysis of PMN in gut

Numbers of PMN (immunopositive for anti-PMN antibody) were counted from 10 randomly selected high power (400 $\times$) fields of jejunum sections, and three sections of each rat were examined. Two observers counted the sections separately, and the results were averaged and tabulated.

Quantitative analysis of mucosal damage

Sloughing of villi tips² was used as a histological criterion of IR induced mucosal damage in CL₂MBP or control liposome treated rats. Five randomly selected low power (100 $\times$) fields of gut mucosa sections were examined, and the numbers of damaged villi and the total number of villi were recorded. Three sections from each rat were examined. Two observers counted the sections separately, and the results were averaged and tabulated.

Reverse transcription-polymerase chain reaction (RT-PCR)

Expression of *Egr-1* and β -actin was analysed by RT-PCR. Full thickness gut wall was used for total RNA isolation using Trizol (Invitrogen, Carlsbad, California, USA). First strand cDNA was synthesised from 1 μ g of total RNA using oligo d(T)₁₂₋₁₈ as a reverse primer and SuperScript First-strand synthesis System (Invitrogen). PCR was performed in 25 μ l

of standard buffer containing primers (1 μ M each; table 1), cDNA (3 μ l), 0.25 units of AmpliTaq Gold (Applied Biosystems, California, USA), MgCl₂ (1.5 mM), and dNTP (0.2 mM each). After initial denaturation (94°C, three minutes), PCR was performed for 30 cycles for *Egr-1*, *MPO*, and β -actin as follows: 30 seconds at 94°C; one minute at 60°C; and one minute at 72°C for each cycle. A final extension for 10 minutes at 72°C was added. PCR products were analysed by electrophoresis in 1% agarose gel. Intensities of PCR bands were quantified and relative expression levels of *Egr-1* to β -actin were determined using Scion Image software (Scion Co, Massachusetts, USA).

Western blotting analysis

Total proteins, and cytoplasmic and nuclear protein fractions were prepared as previously described.³⁰ Equal amounts of protein (50 μ g or 20 μ g in the case of nuclear protein) of each sample were separated in 12.5% sodium dodecyl sulphate polyacrylamide gels before being electrotransferred onto a PVDF membrane (Amersham Bioscience, Hong Kong, China) in a transfer buffer (48 mM Tris, 39 mM glycine, 20% methanol, pH 9.2). The membrane was blocked in PBST (pH 7.2, 0.1% Tween 20) containing 10% (w/v) skimmed milk for one hour before incubation with primary antibodies in PBST against Egr-1 (Geneka Biotechnology, Quebec,

Figure 1 Ischaemia reperfusion (IP) injury of gut occurs before polymorphonuclear neutrophil (PMN) infiltration. Sections of jejunum of different IR groups were stained with haematoxylin and eosin. Clear mucosal damage was observed in the 45/60 minute (45'/60') IR gut (B) but not in the 30/60 minute (30'/60') IR gut (A). Sections of sham controls and IR groups were immunostained with anti-PMN antibody (C-E). Few PMN (arrows) were localised in the 45/60 minute IR jejunum (C). PMNs were detected near and inside the blood capillaries at the base of the crypt of the 60/60 minute IR jejunum (D; inset). Heavy PMN infiltration was detected in the 45/180 minute IR jejunum (E).

Figure 2 Resident macrophages express myeloperoxidase (MPO) and inducible nitric oxide synthase (iNOS) in ischaemia reperfusion (IR) gut. Sections of 45/60 minute (45'/60') IR jejunum were immunostained using anti-MPO and ED2 antibodies (A). ED2 immunoreactive resident macrophages were also immunoreactive for MPO. Adjacent sections were immunostained with anti-iNOS antibody and ED-2 antibody (B). ED2 immunopositive were also iNOS positive. Protein extracts of the sham operation and IR jejunum of 30/60 minute (30'/60') and 45/60 minute (45'/60') ischaemic groups were analysed by western blotting (C). Levels of MPO were increased in the 30 minute and 45 minute IR jejunum (IR) but a negligible level of MPO was detected in the sham operation jejunum (sham). MPO mRNA levels were similar between the sham operation and IR jejunum, as revealed by reverse transcription-polymerase chain reaction (D).

Canada; diluted 1:1000), RANTES, ICAM-1, MCP-1 (Santa Cruz Biotechnology; diluted 1:100), or actin (Santa Cruz Biotechnology; diluted 1:2000) for 16 hours at 4°C. After three PBST washings, blots were incubated with appropriate horseradish peroxidase conjugated secondary antibodies (Zymed, California, USA) in PBST for one hour at ambient temperature. After three PBST washings, signal detection was performed with an ECL kit (Amersham Bioscience).

Statistical analysis

Means (SEM) of the relative expression of *Egr-1* to β -actin, ratio of ED2 to ED1 immunopositive cells, and ratio of damaged villi to total villi were determined for each group. One way (and non-parametric) ANOVA, with Newman-Keuls multiple comparison test was used (GraphPAD Software, San Diego, California, USA) to evaluate the significance of the differences between different treatments, and a p value <0.05 was considered significant.

RESULTS

Localisation of PMN in the gut after IR injury

To define a time point for early intestinal damage after IR, histological analyses were performed on IR guts treated with different periods of ischaemia (10, 30, 45, 60, and 120 minutes) and reperfusion (1, 3, 6, and 15 hours). Histological analyses showed that reperfusion after SMA clamping caused IR induced gut damage characterised by mucosal damage,

sloughing of the mucosal epithelium, and villi destruction observed in the IR jejunum at 45/60 minutes (fig 1B). Mucosal damage was not evident in the jejunum of either the 30/60 minute IR group (fig 1A) or the sham operation group (data not shown). To localise the infiltrated PMN in the IR jejunum, immunohistochemistry using antirat PMN antibody was performed. Only a few PMN were detected in the damaged villi of the jejunum of the 45/60 minute (0.7 (0.16):0.65 (0.23); IR:sham) and 60/60 minute IR groups (1.25 (0.28):0.8 (0.2); IR:sham) (fig 1C, D). In the 60/60 minute IR gut, PMN were localised near and inside the blood capillaries at the base of the crypt (fig 1D). In contrast, extensive PMN infiltration was detected in damaged villi after prolonged reperfusion (three hours) (28.35 (2.65):0.5 (0.15); IR:sham; $p < 0.001$) (fig 1E).

MPO expression and localisation

Absence of infiltrated PMN in damaged villi of the 45/60 minute IR jejunum prompted us to investigate the role of resident macrophages in mucosal damage. We performed immunofluorescent staining on sections of 45/60 minute IR jejunum using anti-MPO and ED2 antibodies (fig 2A). Expression of MPO overlapped with that of ED2. ED2 immunopositive resident macrophages also stained for inducible nitric oxide synthase (iNOS) and lymphocyte function associated antigen 1 (LFA-1) (fig 2B; data not shown). Expression levels of MPO in the IR jejunum and the

Figure 3 Upregulation of early growth response factor 1 (*Egr-1*), RANTES, monocyte chemoattractant protein 1 (MCP-1), and intercellular adhesion molecule 1 (ICAM-1) in ischaemia reperfusion (IR) gut. Reverse transcription-polymerase chain reaction (RT-PCR) was performed to analyse the relative expression of *Egr-1* in the IR (IR) and sham operation (sham) gut of the 30/60 minute (30'/60') IR and 45/60 minute (45'/60') IR groups (A). Weak expression of *Egr-1* mRNA was detected in control jejunum (control) and sham operation jejunum. A marked increase in *Egr-1* mRNA expression in the 30/60 minute IR jejunum and a moderate increase in *Egr-1* mRNA expression in the 45/60 minute IR jejunum were detected. Relative expression of *Egr-1* to β -actin was determined in each group and is shown next to the gel picture of RT-PCR. *Significant difference between the IR and sham jejunum. Protein levels of *Egr-1*, RANTES, ICAM-1, and MCP-1 in the jejunum of different IR groups were analysed by western blotting (B). Actin was used as a loading control for each lane. Elevated levels of *Egr-1*, RANTES, and MCP-1 were detected in the 45/60 minute IR jejunum. Immunoreactivity of ICAM-1 was detected in villi (C), but only expressed in the 45/60 minute IR jejunum in endothelial cells (arrows) of blood capillaries.

sham operation jejunum were analysed using western blotting. Levels of MPO were increased markedly in the 30/60 minute and 45/60 minute IR jejunum while the amount of MPO in the sham operation jejunum was negligible (fig 2C). Similar intensities of actin were observed in all lanes, indicating that equivalent amounts of protein extracts of each sample were analysed. RT-PCR analysis revealed similar levels of MPO mRNA in both sham controls and IR groups (fig 2D) suggesting that MPO induction in our IR model was regulated at the translation level.

Expression of mRNA and protein of *Egr-1* gene in IR gut

To investigate if upregulation of *Egr-1* of resident macrophages is crucial for gut IR injury, a kinetic study of *Egr-1* expression in the IR jejunum was performed. Relative expression levels of *Egr-1* mRNA to β -actin mRNA were determined in IR jejunum, sham operation jejunum, and control jejunum. The relative expression level of *Egr-1* mRNA increased by 3.9-fold in the 30/60 minute IR jejunum compared with sham operation jejunum ($Egr-1/\beta$ -actin 1.8 (0.19):0.46 (0.29); IR:sham; $p < 0.01$) and by threefold in the 45/60 minute IR jejunum compared with the sham jejunum ($Egr-1/\beta$ -actin 0.75 (0.23):0.25 (0.05); IR:sham; $p > 0.05$) (fig 3A). The increase in the relative level of *Egr-1* mRNA in the 30/60 minute IR jejunum over the sham jejunum reached statistical significance ($p < 0.01$). Levels of *Egr-1* protein and protein products of *Egr-1* downstream target genes in the IR jejunum were analysed by western blotting (fig 3B). A marked increase in *Egr-1*, RANTES, and MCP-1 was detected only in the 45/60 minute IR jejunum, and a slight increase in expression of ICAM-1 was also detected in the 45/60 minute IR jejunum. Immunohistochemistry showed expression of ICAM-1 at endothelial cells of the blood capillaries in the 45/60 minute IR jejunum (fig 3C).

Subcellular distribution of *Egr-1* in resident macrophages of IR jejunum

Transcription factor *Egr-1* is translocated into the nucleus to induce expression of its target genes. Subcellular distribution of *Egr-1* in the sham and IR jejunum was investigated by immunohistochemistry (fig 4). In the sham operation group, only a few cells at the mucosal epithelium were stained for *Egr-1* (fig 4A). In the 30/60 minute IR jejunum, clear *Egr-1* immunopositive cells were detected inside the villi, and *Egr-1* protein was mainly localised in the cytoplasm (fig 4B). A larger number of more intensely stained *Egr-1* immunopositive cells were localised in the villi of the 45/60 minute IR jejunum, and *Egr-1* protein was found mainly in the nucleus (fig 4C). Colocalisation studies revealed in 45/60 minute IR injury section that ED2 immunopositive macrophages were immunoreactive for *Egr-1* (fig 4D). Western blotting of the nuclear and cytoplasmic fractions of IR jejunum also revealed the presence of nuclear *Egr-1* in the 45/60 minute IR group (fig 4E).

Effect of resident macrophage depletion on IR injury

To determine if Cl_2 MBP is effective in the depletion of resident macrophages in the gut, Cl_2 MBP liposome and control liposome injected rats were sacrificed three days post-injection, and their gastrointestinal tracts processed for immunohistochemistry to localise the resident macrophages (ED2 immunopositive) and total macrophages (ED1 immunopositive) in the liver and gut. Resident macrophages were markedly decreased both in the liver and gut of Cl_2 MBP liposome treated rats (fig 5A–D). The percentage of resident macrophages to total macrophages decreased from 41.3% (4.9) in control liposome treated guts to 1.8% (0.8) in the Cl_2 MBP liposome treatment group (fig 5). Levels of expression of *Egr-1*, cytokines, and adhesion molecule ICAM-1 were analysed in the jejunum of control and Cl_2 MBP liposome

Figure 4 Resident macrophages expressed a high level of early growth response factor 1 (Egr-1) in the ischaemia reperfusion (IR) jejunum. In the sham operation jejunum, Egr-1 immunoreactivity was not detected in villi and only a few cells at the mucosal epithelium showed Egr-1 expression (A). A few Egr-1 immunopositive cells (arrows) were localised in the villi of the 30/60 minute (30'/60') IR jejunum (B), and the Egr-1 protein was in the cytoplasm (inset). A large number of Egr-1 immunopositive cells (arrows) were localised in the damaged villi of the 45/60 minute (45'/60') IR jejunum (C), and the Egr-1 protein was mainly in the nuclei (inset). Cells showing cytoplasmic Egr-1 staining (Cytoplasmic +) and cells showing nuclear Egr-1 staining (Nuclear +) in villi of different treatment groups were counted and tabulated as shown. An immunofluorescent colocalisation study using anti-Egr-1 antibody and ED2 antibody (D) in 45/60 minute IR injury gut indicated that ED2 immunopositive resident macrophages were also immunopositive for Egr-1 (arrowheads). Distribution of Egr-1 protein in the nuclear and cytoplasmic fractions of jejunum of 30/60 minute and 45/60 minute IR groups were investigated with western blotting. Egr-1 was detected in the nuclear fraction (Egr-1n) of 45/60 minute IR jejunum while Egr-1 was detected in cytoplasmic fractions (Egr-1c) in both groups (E). Actin was used as a loading control for each lane.

treatment groups. mRNA and protein levels of Egr-1 were reduced in the Cl₂MBP liposome treatment group (fig 5E, F). Levels of RANTES and MCP-1 were reduced after Cl₂MBP treatment (fig 5F). In contrast, only a slight reduction in ICAM-1 level was observed in the Cl₂MBP liposome treatment group (fig 5F). To examine if the depletion of gut resident macrophages can alleviate mucosal damage after IR, we performed histological analysis of the 45/60 minute IR jejunum of Cl₂MBP liposome and control liposome treated rats. Sloughing of mucosal epithelium and villi destruction were detected in the control jejunum (fig 6A). In contrast, the mucosa appeared intact and epithelial sloughing was rarely observed in the Cl₂MBP liposome treated jejunum (fig 6B). The percentage of damaged villi to total villi was reduced from 34.2% (2.5) in the control liposome treated jejunum to 5.2% (2.2) in the Cl₂MBP liposome treated jejunum.

DISCUSSION

In this study, we have reported that mucosal damage occurs before PMN infiltration in an IR injured gut in a rat model. IR induced mucosal damage was correlated with activation of gut resident macrophages and upregulation of MPO and

Egr-1 of resident macrophages. Depletion of gut resident macrophages by Cl₂MBP significantly reduced mucosal damage. The present study provides evidence suggesting that activated resident macrophages play a key role in early IR damage to the gut.

Sloughing of mucosal epithelium and ulceration of villi have been recognised as early histological landmarks of IR gut injury,² and these were observed in 45/60 minute IR gut in our model. However, only a few PMN were found at the damaged villi and at the base of the crypt although clear mucosal damage was evident at the IR jejunum. PMN were localised at the base of the crypt in the 60/60 minute IR gut. Massive PMN infiltration was detectable in the damaged area after three hours of reperfusion. This finding is in line with previous reports that PMN infiltration is detectable in damaged tissues 4–6 hours after IR.^{31–33} Taken together, these findings suggest that early mucosa damage in IR injury is not attributable to PMN. Interaction between ICAM-1 on endothelial cells and LFA-1 on PMN is a prerequisite for PMN infiltration.³⁴ ICAM-1 expression of endothelial cells was first observed at 45/60 minute IR gut, which suggested that IR induced mucosal damage began prior to PMN

Figure 5 Intraperitoneal injection of dichloromethylene bisphosphonate (Cl₂MBP) depleted resident macrophages from the gut and liver. Sections from the liver (A, C) and jejunum (B, D) of rats treated with control liposome or Cl₂MBP liposome were immunostained with ED2 antibody to localise resident macrophages. ED2 immunopositive resident macrophages (arrows) were localised in the liver (A) and jejunum (B) of control liposome treated rats but were undetectable in the liver (C) and jejunum (D) of Cl₂MBP liposome treated rats. The percentage of ED2 immunopositive resident macrophages versus ED1 immunopositive total macrophages in the control liposome treated and Cl₂MBP liposome treated jejunum was determined and tabulated as shown (number of views studied for each group = 30). Expression of early growth response factor 1 (Egr-1) mRNA in the control liposome treatment group and the Cl₂MBP treatment group was analysed by reverse transcription-polymerase chain reaction (E). Levels of Egr-1, cytokines, and intercellular adhesion molecule 1 (ICAM-1) in the jejunum of the control liposome treatment group and the Cl₂MBP treatment group were analysed by western blotting (F). Reduction of Egr-1, RANTES, and monocyte chemoattractant protein 1 (MCP-1) was observed in the Cl₂MBP treatment group. In contrast, only a slight reduction in the level of ICAM-1 was observed in the Cl₂MBP treatment group compared with that in the control liposome treatment group.

infiltration, and corroborates a report that anti-PMN anti-serum fails to prevent tissue damage after IR in rats.³³ Our results also suggest the importance of ischaemia in the priming of gut damage in which longer ischaemia results in more damage after reperfusion.

Elevated tissue MPO enzyme activity has been frequently used as an indication of PMN infiltration in IR injuries.¹⁸ However, monocytes and macrophages also express MPO, which indicates that elevated tissue MPO activity is not necessarily derived from infiltrated PMN. Furthermore, IR induced mucosal damage began prior to PMN infiltration, which indicated that elevated MPO activity in the early phase of IR gut injury was not attributed to PMNs. Immunostaining using an anti-MPO antibody and tissue macrophage marker (ED2) revealed that resident macrophages in IR guts expressed MPO enzyme. Activated macrophages *in vitro*¹⁴ have already been shown to express a high level of MPO activity. Indeed, resident macrophages in the IR jejunum

expressed activated macrophage markers LFA-1³⁵ and iNOS,³⁶ suggesting that resident macrophages were activated in the IR gut. Elevated expression of MPO in resident macrophages and increased tissue MPO levels were detected in the 30/60 minute IR jejunum, and mucosal damage was evident in the 45/60 minute IR jejunum. MPO catalyses the production of OCl⁻ and chloramines which have already been demonstrated to cause gastric mucosal cell apoptosis and cell sloughing.^{16, 17} Elevated MPO activity is associated with pulmonary injury, renal glomerular damage, and the initiation of atherosclerotic lesions.³⁷ We failed to detect mucosal damage in the jejunum after prolonged ischaemia (two hours; unpublished data). This suggests that the oxygen burst during reperfusion triggers the cascade of cellular activations and molecular events leading to IR damage.

To prove that resident macrophages were involved in gut IR injury, we compared IR damage in Cl₂MBP liposome and control liposome treated rats. Cl₂MBP liposome can

Figure 6 Ischaemia reperfusion (IR) induced mucosal damage was markedly reduced in dichloromethylene bisphosphonate (Cl₂MBP) liposome treated jejunum. Haematoxylin and eosin staining of sections of the 45/60 minute IR jejunum of control liposome treated rats (A) and Cl₂MBP liposome treated rats (B) are shown. The percentage of damaged villi versus total villi in the control liposome and Cl₂MBP liposome treated rats was determined and tabulated as shown.

Treatment	Damage villi / total villi
Control liposome	34.2% (2.5%)
Cl ₂ MBP-liposome	5.2% (2.2%)

specifically deplete resident macrophages in the gut without changing PMN activity.^{38 39} The percentage of resident macrophages (ED2 immunopositive) to total macrophages (ED1 immunopositive) in the gut decreased from 41.3% in control liposome treated guts to only 1.8% in the Cl₂MBP liposome treated gut. Only ED2 immunopositive resident macrophages were depleted in Cl₂MBP treated rats indicating that Cl₂MBP was specific in the depletion of resident macrophages. The degree of IR induced mucosal damage and villi destruction was markedly reduced in the Cl₂MBP liposome treated jejunum compared with the control jejunum. Mucosal damage was not detectable in resident macrophages depleted rats even after three hours of reperfusion, and very few PMN were localised at the villi. Our findings on IR injury to resident macrophage depleted guts suggest that resident macrophages contribute to the initiation of mucosal damage in IR.

Colocalisation analysis established that resident macrophages in the damaged area expressed Egr-1. Activation of resident macrophages induces mRNA and protein expression of *Egr-1* gene, and nuclear compartmentalisation of Egr-1 protein in resident macrophages. Involvement of Egr-1 in lung IR injury has been shown in mice. PMN infiltration and upregulation of cytokines, including RANTES and MCP-1, in mouse lungs after IR challenge are markedly reduced in *Egr-1* knockout mice and these mice show very mild IR damage in lung tissue.²² The reduction in Egr-1, RANTES, and MCP-1 after Cl₂MBP treatment indicated that macrophages produced these cytokines. However, studies of gut IR in *Egr-1* knockout mice could further clarify the Egr-1 function in the onset of gut IR damage. Our data suggest the PMN infiltration by Egr-1 upregulation is not directly implicated in the onset of gut IR injury. Reduction of PMN infiltration in Cl₂MBP treated guts after IR is likely to be an indirect effect

of reduced macrophage derived cytokines that induce PMN infiltration.⁴⁰

In summary, activated resident macrophages in IR gut express a high level of MPO, Egr-1, and proinflammatory cytokines. Elevated MPO activity catalyses the production of reactive free radicals during reperfusion, causing initial mucosal epithelial cell death. Accumulation of proinflammatory cytokines induces PMN infiltration, resulting in further tissue damage. Further exploration of potential therapeutic intervention for IR damage might therefore profitably be directed towards resident macrophages, as they have the capacity to prevent initial mucosal destruction and arrest subsequent tissue damage.

Authors' affiliations

Y Chen, V C H Lui, P K H Tam, Division of Paediatric Surgery, Department of Surgery, University of Hong Kong Medical Centre, Queen Mary Hospital, University of Hong Kong, China
N V Rooijen, Department of Cell Biology, Division of Histology, Faculty of Medicine, Free University, Amsterdam, the Netherlands

REFERENCES

1 **Jrvinen O**, Laurikka J, Salenius JP, *et al*. Acute intestinal ischaemia. A review of 214 cases. *Ann Chir Gynaecol* 1994;**83**:22-5.
2 **Kong SE**, Blennerhassett LR, Heel KA, *et al*. Ischaemia-reperfusion injury to the intestine. *Aust N Z J Surg* 1998;**68**:554-61.
3 **Jaeschke H**. Mechanisms of reperfusion injury after warm ischemia of the liver. *J Hepatobiliary Pancreat Surg* 1998;**5**:402-8.
4 **Jordan JE**, Zhao ZQ, Vinten-Johansen J. The role of neutrophils in myocardial ischemia-reperfusion injury. *Cardiovasc Res* 1999;**43**:860-78.
5 **Turnage RH**, Guice KS, Oldham KT. Pulmonary microvascular injury following intestinal reperfusion. *New Horiz* 1994;**2**:463-75.
6 **Kohtani T**, Abe Y, Sato M, *et al*. Protective effects of anti-neutrophil antibody against myocardial ischemia/reperfusion injury in rats. *Eur Surg Res* 2002;**34**:313-20.

- 7 Andoh A, Fujiyama Y, Araki Y, *et al*. Role of complement activation and mast cell degranulation in the pathogenesis of rapid intestinal ischemia/reperfusion injury in rats. *Digestion* 2001;**63**(Suppl 1):103–7.
- 8 Nagaoaka I, Kaneko H, Yamashita T. Inhibition of the accumulation of macrophages and the generation of macrophage chemotactic activity by dexamethasone in concanavalin A-induced peritonitis of mice. *Agents Actions* 1988;**25**:156–63.
- 9 Hudig D, Redelman D, Minning L, *et al*. Inhibition of human lymphocyte natural cytotoxicity and antibody-dependent cell-mediated cytotoxicity by K-76 COONa, a reagent that blocks complement activity. *J Immunol* 1984;**133**:408–14.
- 10 Coates NJ, McCall SR. Production of chemokines in vivo in response to microbial stimulation. *J Immunol* 2001;**166**:5176–82.
- 11 Askari AT, Brennan ML, Zhou X, *et al*. Myeloperoxidase and plasminogen activator inhibitor 1 play a central role in ventricular remodeling after myocardial infarction. *J Exp Med* 2003;**197**:615–24.
- 12 Zimmerman BJ, Grisham MB, Granger DN. Role of oxidants in ischemia/reperfusion-induced granulocyte infiltration. *Am J Physiol* 1990;**258**:G185–90.
- 13 Heinecke JW, Li W, Daehnke HLR, *et al*. Dityrosine, a specific marker of oxidation, is synthesized by the myeloperoxidase-hydrogen peroxide system of human neutrophils and macrophages. *J Biol Chem* 1993;**268**:4069–77.
- 14 Rodriguez MR, Rodriguez D, Russo M, *et al*. Macrophage activation includes high intracellular myeloperoxidase activity. *Biochem Biophys Res Commun* 2002;**292**:869–73.
- 15 Duffield JS. The inflammatory macrophage: a story of Jekyll and Hyde. *Clin Sci (Lond)* 2003;**104**:27–38.
- 16 Naito Y, Yoshikawa T, Fujii T, *et al*. Monochloramine-induced cell growth inhibition and apoptosis in a rat gastric mucosal cell line. *J Clin Gastroenterol* 1997;**25**(suppl 1):S179–85.
- 17 Suzuki H, Seto K, Mori M, *et al*. Monochloramine induced DNA fragmentation in gastric cell line MKN45. *Am J Physiol* 1998;**275**:G712–16.
- 18 De Greef KE, Ysebaert DK, Ghielli M, *et al*. Neutrophils and acute ischemia-reperfusion injury. *J Nephrol* 1998;**11**:110–22.
- 19 Chiu CJ, McArdle AH, Brown R, *et al*. Intestinal mucosal lesion in low-flow states. I. A morphological, hemodynamic, and metabolic reappraisal. *Arch Surg* 1970;**101**:478–83.
- 20 Hierholzer C, Kalff JC, Audolfsson G, *et al*. Molecular and functional contractile sequelae of rat intestinal ischemia/reperfusion injury. *Transplantation* 1999;**68**:1244–54.
- 21 Beckmann AM, Wilce PA. Egr transcription factors in the nervous system. *Neurochem Int* 1997;**31**:477–510.
- 22 Yan SF, Fujita T, Lu J, *et al*. Egr-1, a master switch coordinating upregulation of divergent gene families underlying ischemic stress. *Nat Med* 2000;**6**:1355–61.
- 23 Aeber H, Cornelius T, Ehr T, *et al*. Expression of immediate early genes after cardioplegic arrest and reperfusion. *Ann Thorac Surg* 1997;**63**:1669–75.
- 24 Brand T, Sharma HS, Fleischmann KE, *et al*. Proto-oncogene expression in porcine myocardium subjected to ischemia and reperfusion. *Circ Res* 1992;**71**:1351–60.
- 25 Bonventre JV, Sukhatme VP, Bamberger M, *et al*. Localization of the protein product of the immediate early growth response gene, Egr-1, in the kidney after ischemia and reperfusion. *Cell Regul* 1991;**2**:251–60.
- 26 Ouellette AJ, Malt RA, Sukhatme VP, *et al*. Expression of two “immediate early” genes, Egr-1 and c-fos, in response to renal ischemia and during compensatory renal hypertrophy in mice. *J Clin Invest* 1990;**85**:766–71.
- 27 Van Rooijen N, Sanders A. Liposome mediated depletion of macrophages: mechanism of action, preparation of liposomes and applications. *J Immunol Methods* 1994;**174**:83–93.
- 28 van den Berg TK, Dopp EA, Dijkstra CD. Rat macrophages: membrane glycoproteins in differentiation and function. *Immunol Rev* 2001;**184**:45–57.
- 29 Barbe E, Damoiseaux JG, Dopp EA, *et al*. Characterization and expression of the antigen present on resident rat macrophages recognized by monoclonal antibody ED2. *Immunobiology* 1990;**182**:88–99.
- 30 Deryckere F, Gannon F. A one-hour miniprep technique for extraction of DNA-binding proteins from animal tissues. *Biotechniques* 1994;**16**:405.
- 31 Koike K, Moore EE, Moore FA, *et al*. CD11b blockade prevents lung injury despite neutrophil priming after gut ischemia/reperfusion. *J Trauma* 1995;**39**:23–7.
- 32 Schoenberg MH, Poch B, Younes M, *et al*. Involvement of neutrophils in postischemic damage to the small intestine. *Gut* 1991;**32**:905–12.
- 33 Simpson R, Alon R, Kobzik L, *et al*. Neutrophil and nonneutrophil-mediated injury in intestinal ischemia-reperfusion. *Ann Surg* 1993;**218**:444–53.
- 34 Olanders K, Sun Z, Borjesson A, *et al*. The effect of intestinal ischemia and reperfusion injury on ICAM-1 expression, endothelial barrier function, neutrophil tissue influx, and protease inhibitor levels in rats. *Shock* 2002;**18**:86–92.
- 35 Kalff JC, Turler A, Schwarz NT, *et al*. Intra-abdominal activation of a local inflammatory response within the human muscularis externa during laparotomy. *Ann Surg* 2003;**237**:301–15.
- 36 Hori M, Kita M, Torihashi S, *et al*. Upregulation of iNOS by COX-2 in muscularis resident macrophage of rat intestine stimulated with LPS. *Am J Physiol Gastrointest Liver Physiol* 2001;**280**:G930–8.
- 37 Klebanoff SJ. Myeloperoxidase. *Proc Assoc Am Physicians* 1999;**111**:383–9.
- 38 Fryer J, Grant D, Jiang J, *et al*. Influence of macrophage depletion on bacterial translocation and rejection in small bowel transplantation. *Transplantation* 1996;**62**:553–9.
- 39 Schmidt-Weber CB, Rittig M, Buchner E, *et al*. Apoptotic cell death in activated monocytes following incorporation of clodronate-liposomes. *J Leukoc Biol* 1996;**60**:230–44.
- 40 Ebe Y, Hasegawa G, Takatsuka H, *et al*. The role of Kupffer cells and regulation of neutrophil migration into the liver by macrophage inflammatory protein-2 in primary listeriosis in mice. *Pathol Int* 1999;**49**:519–32.

EDITOR'S QUIZ: GI SNAPSHOT

Answer

From question on page 1771

Terminal ileoscopy revealed multiple red spots and raised cherry red bumps 3–5 mm in diameter, with a mosaic pattern of the mucosa around their base. Biopsies revealed oedema in the mucosa and arteriolar dilations that sometimes contained fibrin thrombi and were compatible with hypertensive portal ileopathy. Videocapsule endoscopy revealed diffuse involvement of the small intestine, with increasingly severe vascular lesions from the proximal jejunum up to the distal ileum where these particular raised red bumps were exclusively located. All of the lesions were attributed to portal hypertension enteropathy and considered to be the cause of the recurrent gastrointestinal bleeding. The patient was given 80 mg/day propranolol and six months later he no longer exhibited either melena or a relapsing decrease in haemoglobin.

This case exhibited a novel endoscopic appearance of portal hypertensive enteropathy characterised by raised red bumps in the ileum, combined with particular histological features similar to those of gastric antral vascular ectasia.

doi: 10.1136/gut.2003.034611