CASE REPORT

Iron chelation therapy in aceruloplasminaemia: study of a patient with a novel missense mutation

R Mariani, C Arosio, S Pelucchi, M Grisoli, A Piga, P Trombini, A Piperno

Gut 2004;53:756–758. doi: 10.1136/gut.2003.030429

We describe a novel missense mutation of ceruloplasmin in a patient with aceruloplasminaemia causing the replacement of a neutral amino acid (phenylalanine) with a polar one (serine) at position 198, probably leading to abnormal folding and secretion of the protein. The patient showed mild microcytic anaemia, mild hepatic iron overload, and marked brain iron overload. Six months of therapy with deferiprone was ineffective in removing iron from the tissues. Deferoxamine was more efficient in removing excess iron from the liver but aggravated the disease related anaemia. After more than one year of chelation treatment, the brain magnetic resonance imaging signal did not change. Overall, these findings indicate that treatment of iron overload in aceruloplasminaemia is a difficult challenge and that new iron chelators, more efficient in crossing the blood-brain barrier, are needed.

Ceruloplasmin (Cp) has an essential role in normal iron homeostasis favouring cellular iron release and iron incorporation into apo-transferrin.^{1 2} Aceruloplasminaemia is a rare autosomal recessive disorder due to mutations of the Cp gene characterised by iron overload in the brain, pancreas, retina, and liver due to cellular iron retention and increased iron absorption.³ In adulthood, patients develop ataxia and dementia, diabetes, and retinal degeneration.^{4 5} Mild microcytic anaemia, low serum iron, and increased serum ferritin are typical features and are related to the defective cellular iron release into plasma.^{3 6 7} Plasma may be helpful but long term use has not been challenged and the risk of infections should be considered.⁴ Phlebotomy is not helpful⁸ and may worsen the disease related anaemia. Different outcomes have been reported using deferoxamine^{7 9} whereas no data are available for the oral iron chelator deferiprone that can be an attractive therapeutic candidate due its stability, lipophilicity, and ability to penetrate the blood-brain barrier.¹⁰

We report the case of a young woman with aceruloplasminaemia due to a novel missense mutation of the Cp gene in whom we also evaluated the efficacy of both deferiprone and deferoxamine on iron removal and their effect on iron and haematological indices.

CASE REPORT

A 40 year old Italian woman was investigated for chronic asthenia and mild microcytic anaemia. Haemoglobin was 10.6 g/dl and mean corpuscular volume 74 fl, serum iron was 27 µg/l, transferrin saturation 10%, and serum ferritin level 471 µg/l in the absence of inflammatory alterations (C reactive protein 0.3). No alteration in glucose or lipid metabolism was found. Cp was undetectable on immunoturbidimetric assay (Tina-quant; Roche, Mannheim, Germany) and at western blot analysis using human

anticeruloplasmin antibody (DakoCytomation, Milan, Italy). Ferroxidase activity was not measured. General, neurological, and fundus oculi examinations were normal.

Imaging studies

Magnetic resonance imaging (MRI) of the proband's brain showed marked hypointensity in the dentate and red nuclei, thalamus, pulvinar, neostriatum, and putamen. Gradient echo images also showed a thin hypointense rim along most of the cerebellar and cerebral cortex. Magnetic susceptibility by superconductive quantum interference device (SQUID) analysis, performed according to Brittenham and colleagues,¹¹ showed a moderate hepatic iron overload (fig 1B).

Genetic investigation and family study

There was no history of consanguinity in the family although the grandparents originated from the same village in Southern Italy. Only the proband's younger brother and offsprings were available for clinical and laboratory analysis, and genetic study. Direct sequencing of the whole Cp gene, including exon 1 to 20 and intron-exon boundaries, was carried out by ABI Prism 3100 Avant DNA sequencer (PE Applied Biosystems, Foster City, California, USA). Sequences were compared with GenBank Accession Nos D45028-D45045 and HGMD No M13699. Four nucleotide changes were detected at the homozygous state in the proband: two intronic T→C transversions (IVS2 +20 and IVS15 -12), previously described in healthy subjects;¹² a never described G→A transversion (AAG→AAA) at nucleotide 2682 in exon 16 that does not cause an amino acid change (K894K) and that was also present in homozygosity in the other unaffected relatives; and a T→C transversion (TTT→TCT) at nucleotide 593 in exon 4 causing a phenylalanine to serine replacement in the protein (F198S). The F198S mutation was confirmed in the proband by restricted fragment length polymorphism (RFLP) analysis using Hinf I digestion and was not found in 50 healthy individuals. The Cp gene sequence and RFLP analysis were performed in all available relatives. The proband's family pedigree, together with the results of the molecular study, biochemical tests, and SQUID measurements are shown in fig 1A and 1B. A mild hepatic iron overload was observed in the proband's brother, daughter, and son at SQUID (fig 1B). MRI was normal in the proband's offspring whereas a mild T2 weighted hypointensity in the pulvinar and posterior and lateral putamen was found in the brother.

Iron chelation therapy

The patient's written informed consent and local ethics committee approval were obtained. Deferiprone (75 mg/kg/day) was administered for six months but was not effective

Abbreviations: Cp, ceruloplasmin; MRI, magnetic resonance imaging; SQUID, superconductive quantum interference device; RFLP, restricted fragment length polymorphism

Figure 1 (A) Family pedigree. (B) Age, iron, and biochemical data of the proband and relatives. SQUID, superconductive quantum interference device.

(fig 2). Deferoxamine (20 mg/kg/day for five days a week) was administered by subcutaneous bolus injections for the next eight months (fig 2). Urinary iron excretion levels and haemoglobin, mean corpuscular volume, serum iron, and serum ferritin behaviour during the treatment periods are shown in fig 2A and B. A second brain MRI, performed after more than one year of chelation treatment, was unchanged.

DISCUSSION

We describe the case of a young woman with aceruloplasmin-
aemia, homozygous for a new missense Cp gene mutation
(F198S). The phenylalanine 198 residue is located just before
Val¹⁹⁹ that forms together with other non-polar residues
(Leu⁷⁴, Ile⁷⁷, Ile¹⁵⁹, and Phe²⁶⁷) a tightly packed hydrophobic
pocket critical for correct folding of human Cp.¹³ A proline
residue at position 177 projects into that pocket and a
mutation changing proline to arginine at this position
(P177R), causing retention of the mutant human Cp in the
endoplasmic reticulum and aceruloplasminaemia, was
recently described.¹³ We suggest that replacement of a neutral
amino acid (phenylalanine) with a polar one (serine) at
position 198 leads to incorrect folding of the protein and
impaired cellular trafficking of Cp, as demonstrated for the
P177R mutant.¹³ The proband had a slight microcytic
anaemia with low serum iron and a moderate hepatic iron
overload not in the range capable of inducing hepatic
damage. In contrast, brain MRI showed marked iron over-
load in the basal nuclei, and mild iron overload in the
cerebellar and cerebral cortex.³ Absence of neurological

abnormalities in the patient is probably explained by her
young age as most patients with aceruloplasminaemia
develop neurological symptoms in their sixties,^{7 8 12} indicating
that brain iron overload requires several years before cellular
damage becomes clinically manifest. The three heterozygous
relatives showed slightly increased hepatic iron stores by
SQUID and the oldest one showed a reduction in the brain
MRI signal suggesting mild iron accumulation in the basal
nuclei. Heterozygosity for aceruloplasminaemia is generally
considered unable to affect iron homeostasis but recent
reports noted neurological abnormalities in elderly individ-
uals with a single Cp mutation showing brain iron
deposition at MRI and autopsy.^{14 15} Follow up studies are
needed to understand whether iron overload increases with
time and leads to clinical manifestations in heterozygous
individuals.

At the time of the study the patient had no major sign or
symptom of the disease but the amount of iron accumulation
in the brain makes the development of neurological altera-
tions likely in the future. For this reason we challenged the
efficacy of the two iron chelators available for human
therapy. Deferiprone was not effective in removing iron from
tissues, as demonstrated by the very low urinary induced iron
excretion and lack of a reduction in serum ferritin after five
months of therapy. The patient was highly compliant with all
diagnostic and therapeutic procedures, and hence we suggest
that deferiprone failure may be explained by individual
variability in deferiprone response and/or by the lower
efficacy of deferiprone compared with deferoxamine.¹⁰

Figure 2 (A) Urinary iron excretion (U-Fe) by deferiprone and deferoxamine. (B) Serum iron (s-Fe), serum ferritin (s-Ferr), haemoglobin (Hb), and mean corpuscular volume (MCV) during iron chelation therapy.

Deferoxamine induced iron excretion was higher than that induced by deferiprone. Serum ferritin halved during the first four months of deferoxamine treatment but a concomitant decrease in haemoglobin and serum iron level was observed, suggesting that deferoxamine sequestered iron available for erythropoiesis. The brain MRI signal did not change in our patient after more than one year of iron chelation treatment although it is possible that minor changes were not detectable as MRI is a semiquantitative method of brain iron measurement. Loréal and colleagues⁷ reported normalisation of serum ferritin and hepatic iron concentration in a woman with aceruloplasminaemia after one year of deferoxamine plus vitamin C treatment together with aggravation of anaemia leading to discontinuation of treatment. Brain iron accumulation and neurological manifestations did not improve. In contrast, Miyajima and colleagues⁹ showed slight improvement in neurological symptoms, decrease in brain iron stores by MRI, and no adverse effects in a 63 year old woman with aceruloplasminaemia after 10 months of a less intensive deferoxamine treatment that induced a partial decrease in hepatic iron overload and serum ferritin levels.

In conclusion, treatment of iron overload in aceruloplasminaemia remains a difficult challenge. Deferoxamine seems to be more efficient than deferiprone in removing excess iron from the liver but it should be administered cautiously to avoid aggravation of the disease related anaemia. Both deferoxamine and deferiprone seem unable to remove sufficient amounts of iron from the brain although it cannot

be excluded that they can prevent further iron accumulation and progression of neurological symptoms. Development of new iron chelators (capable of crossing the blood-brain barrier) and of more sensitive methods (capable of quantifying brain iron) could improve treatment and monitoring of therapeutic efficacy in brain iron overload disorders.

ACKNOWLEDGEMENTS

We thank Filomena Longo, MD (Thalassemia Centre, Department of Paediatric Sciences, University of Turin), for providing us with the SQUID analysis. We also thank Donatella Barisani, MD (Department of Experimental, Environmental Medicine and Medical Biotechnology, University of Milano-Bicocca) for performing the western blot analysis.

Authors' affiliations

R Mariani, C Arosio, S Pelucchi, P Trombini, A Piperno, Clinica Medica, Azienda Ospedaliera San Gerardo, Università Milano-Bicocca, Monza, Italy

M Grisoli, Istituto Nazionale Neurologico "C Besta", Milano, Italy

A Piga, Dipartimento di Scienze Pediatriche, Azienda Sanitaria Ospedaliera S Anna, Università Torino, Italy

Correspondence to: Dr A Piperno, Clinica Medica, Azienda Ospedaliera San Gerardo, Via Donizetti 106, 20052, Monza, Italy; apiperno@tiscali.it

Accepted for publication 23 December 2003

REFERENCES

- Mukhopadhyay CK, Attieh ZK, Fox PL. Role of ceruloplasmin in cellular iron uptake. *Science* 1998;**279**:714-17.
- Harris ZL, Durlay AP, Man TK, et al. Targeted gene disruption reveals an essential role for ceruloplasmin in cellular iron efflux. *Proc Natl Acad Sci U S A* 1999;**96**:10812-17.
- Niitts T, Gitlin JD. The copper-iron connection: hereditary aceruloplasminemia. *Semin Hematol* 2002;**39**:282-9.
- Yonekawa M, Okabe T, Asamoto Y, et al. A case of hereditary ceruloplasmin deficiency with iron deposition in the brain associated with chorea, dementia, diabetes mellitus and retinal pigmentation: administration of fresh-frozen human plasma. *Eur Neurol* 1999;**42**:157-62.
- Morita H, Ikeda S, Yamamoto K, et al. Hereditary ceruloplasmin deficiency with hemosiderosis: a clinicopathological study of a Japanese family. *Ann Neurol* 1995;**37**:646-56.
- Bosio S, De Gobbi M, Roetto A, et al. Anemia and iron overload due to compound heterozygosity for novel ceruloplasmin mutations. *Blood* 2002;**100**:2246-8.
- Loréal O, Turlin B, Pigeon C, et al. Aceruloplasminemia: new clinical, pathophysiological and therapeutic insights. *J Hepatol* 2002;**36**:851-6.
- Hellman NE, Schafer M, Gehrke S, et al. Hepatic iron overload in aceruloplasminemia. *Gut* 2000;**47**:858-60.
- Miyajima H, Takahashi Y, Kamats T, et al. Use of desferrioxamine in the treatment of aceruloplasminemia. *Ann Neurol* 1997;**41**:404-7.
- Beutler EA, Hoffbrand VA, Cook JD. Iron deficiency and overload. *Hematology (Am Soc Hematol Educ Program)*, 2003:40-61.
- Brittenham GM, Farrell DE, Harris JW, et al. Magnetic-susceptibility measurement of human iron stores. *N Engl J Med* 1982;**307**:1671-5.
- Lee PL, Gelbart T, West C, et al. A study of genes that may modulate the expression of hereditary hemochromatosis: transferrin receptor-1, ferroportin, ceruloplasmin, ferritin light and heavy chain, iron regulatory proteins (IRP)-1 and -2, and hepcidin. *Blood Cells Mol Dis* 2001;**27**:783-802.
- Hellman NE, Kono S, Miyajima H, et al. Biochemical analysis of a missense mutation in aceruloplasminemia. *J Biol Chem* 2002;**277**:1375-80.
- Daimon M, Morial S, Susa A, et al. Hypoaceruloplasminemia with heteroallelic ceruloplasmin gene mutation: MRI of the brain. *Neuroradiology* 1999;**41**:185-7.
- Miyajima H, Kono S, Takahashi Y, et al. Cerebellar ataxia associated with heteroallelic ceruloplasmin gene mutation. *Neurology* 2001;**57**:2205-10.