STEATOSIS IN CHRONIC HEPATITIS C: WHY DOES IT REALLY MATTER?

RECENT ADVANCES IN CLINICAL PRACTICE

T Asselah, L Rubbia-Brandt, P Marcellin, F Negro

123

Gut 2006;55:123–130. doi: 10.1136/gut.2005.069757

SUMMARY

Hepatic steatosis is a common histological feature of chronic hepatitis C. Various factors are associated with hepatic steatosis, including obesity, high alcohol consumption, diabetes type II, and hyperlipidaemia. These factors may contribute to steatosis in patients with chronic hepatitis C. In humans, hepatitis C virus (HCV) genotype 3 is more commonly associated with steatosis. In vitro studies and the transgenic mouse model have suggested that the HCV core protein (genotype 1) can induce lipid accumulation within hepatocytes.

However, what is the relevance of steatosis in chronic hepatitis C? It seems that in certain populations, steatosis may be associated with fibrosis progression and this may be genotype specific. The mechanisms underlying this association are unknown; neither is it clear whether this holds true for all patients or only a subgroup. Indeed, after antiviral treatment, virus related steatosis disappears whereas the host associated steatosis remains unaffected.

This review describes and discusses the basic and clinical aspects of the relationship between steatosis and progression of fibrosis, and response to treatment in patients with chronic hepatitis C.

INTRODUCTION

HCV is a major cause of chronic liver disease with about 170 million people infected worldwide. The severity of disease varies widely from asymptomatic chronic infection to cirrhosis and hepatocellular carcinoma.

Non-alcoholic fatty liver disease (NAFLD) represents a spectrum from simple steatosis at one end to severe inflammation with extensive fibrosis or cirrhosis at the other. Hepatic steatosis without inflammation is thought to have a good prognosis but non-alcoholic steatohepatitis (NASH) can progress to cirrhosis in a significant proportion of cases.

Hepatic steatosis is also a common histological feature of chronic hepatitis C but may be independently associated with obesity, high alcohol consumption, type II diabetes, and hyperlipidaemia and, when these occur, may contribute to steatosis in patients with chronic hepatitis C. In this review we will focus on the relevance of steatosis in patients with chronic hepatitis C with respect to disease outcome.

FREQUENCY OF STEATOSIS IN CHRONIC HEPATITIS C PATIENTS IN COMPARISON WITH THE GENERAL POPULATION

Hepatic steatosis can develop secondary to obesity, diabetes mellitus, alcohol abuse, protein malnutrition, total parenteral nutrition, acute starvation, drug therapy, carbohydrate overload,^{1–7} and chronic hepatitis C infection. An excellent review of lipid metabolism has been published elsewhere (fig 1).⁵ Estimates based on imaging and autopsy studies suggest that approximately 20–30% of adults in the USA and other Western countries have excess fat accumulation in the liver^{1–2}; approximately 10% of these individuals (that is, 2–3% of adults) are estimated to have NASH. Wanless and Lentz found steatosis in 70% of obese and 35% of lean patients in a consecutive autopsy study.⁸ The prevalence of steatosis in patients with abnormal liver tests (in which known causes of liver disease have been excluded) is 64%; of these 50% had steatosis and 50% had steatohepatitis.⁹

In chronic hepatitis C patients, the prevalence of steatosis ranges from 40% to 86% (mean ~55%) (table 1).^{10–23} The majority of patients with steatosis (78%) have mild steatosis affecting less than 30% of hepatocytes. Thus steatosis occurs more frequently in patients with chronic hepatitis C (55%) than in the general population (20–30%) of adults in the Western world.²

FACTORS ASSOCIATED WITH STEATOSIS IN CHRONIC HEPATITIS C PATIENTS

Patients with HCV infection may have coexisting obesity, diabetes, alcohol abuse, etc., which contribute to the development of fatty liver (box 1). In many studies a link between steatosis and

See end of article for authors' affiliations

Correspondence to:
Dr P Marcellin, Service
d'Hépatologie and INSERM
CRB3, Hôpital Beaujon, Clichy
92 110, France;
patrick.marcellin@
bjn.ap-hop-paris.fr

Figure 1 Pathophysiology of steatosis. Hepatic fat accumulation occurs when the fatty acid input exceeds the amount that can be either secreted as very low density lipoprotein (VLDL) or oxidised as energy source (output). The main causes of fatty acid accumulation in the liver are numbered below. (1, 2) Increased hepatic uptake of fatty acids. Fatty acids are mobilised from peripheral adipocytes under the action of hormone sensitive lipoprotein lipase. Insulin resistance leads to increased lipolysis with accumulation of circulating fatty acids; this is then followed by increased passive uptake in hepatocytes. (3) Increased glycolysis. Insulin resistance/hyperinsulinaemia stimulates glycolysis with synthesis of fatty acids. (4) Decreased removal of fatty acids due to impaired mitochondrial β -oxidation. (5) Decreased export of fatty acids secreted from the hepatocyte bound to apolipoprotein B as VLDL.

a high body mass index (BMI) has been reported (table 2). Adinolfi *et al* reported that overall, steatosis was not significantly associated with BMI, but when analysis was done by single genotype, a significant association was shown between high visceral fat distribution and grade of steatosis in HCV genotype 1 patients (but not in genotype 3).¹⁴ This led to the idea that in patients with HCV infection there is a

“metabolic fat” (especially in patients with HCV genotype 1) and a “viral fat” (especially in patients with genotype 3). Epidemiological studies have suggested a link between type 2 diabetes and chronic HCV infection.^{24–28} However, the presence of confounding factors such as obesity, aging, or cirrhosis precludes the establishment of a definite relationship between these two conditions.

Table 1 Frequency and distribution of steatosis in patients with chronic hepatitis C

Author, year	Patients (n)	Patients with steatosis (n (%))			Distribution of steatosis (n (%))	
		Overall	Genotype 3	Other	Mild (<30%)	Marked (>30%)
Mihm 1997 ¹⁰	85	73 (86)	ND	ND	60 (82)	13 (18)
Czaja 1998 ¹¹	60	31 (52)	ND	ND	31 (100)	0 (0)
Hourigan 1999 ¹²	148	91 (61)	14/17 (78)	14/23 (61)	61 (67)	30 (33)
Rubbia-Brandt 2000 ¹³	70	28 (40)	16/24 (67)	12/46 (26)	18 (65)	10 (35)
Adinolfi 2001 ¹⁴	180	86 (48)	20/26 (77)	66/154 (43)	44 (51)	42 (49)
Serfaty 2001 ¹⁵	100	ND	ND	ND	88 (88)	12 (12)
Monto 2002 ¹⁶	297	171 (58)	ND	ND	146 (85)	25 (15)
Westin 2002 ¹⁷	98	41 (42)	22/25 (88)	11/45 (24)	25 (61)	16 (39)
Hui 2002 ¹⁸	124	90 (73)	33/43 (77)	50/71 (70)	55 (61)	35 (39)
Castéra 2003 ¹⁹	96	51 (54)	15/20 (75)	36/76 (47)	42 (82)	9 (18)
Poynard 2003 ²⁰	1428	935 (65)	175/210 (83)	760/1218 (62)	836 (89)	99 (11)
Asselah 2003 ²¹	290	135 (46)	36/58 (63)	97/232 (42)	91 (68)	44 (32)
Rubbia-Brandt 2004 ²²	755	315 (42)	109/178 (61)	206/577 (36)	206 (65)	109 (35)
Patton 2004 ²³	574	277 (48)	61/84 (91)	216/490 (44)	187 (68)	90 (32)
Total	4305	2324 (55)	501/685 (73)	1468/2932 (50)	1890 (78)	531 (22)

ND, not done.

Box 1 Factors associated with steatosis in chronic hepatitis C

- ▶ Viral factor
 - HCV genotype 3
- ▶ Host factors
 - Alcohol consumption
 - Overweight
 - Hyperlipidaemia
 - Diabetes
 - Insulin resistance
- ▶ Drug therapy
 - Corticosteroids, amiodarone, methotrexate, etc....

STEATOSIS, HCV GENOTYPE 3, AND VIRAL REPLICATION

Several studies have observed a significant association between HCV genotype 3 infection and the presence of steatosis (table 1). In our study, including 290 chronic hepatitis C patients, steatosis, in multivariate analysis, was associated with HCV genotype 3 infection, higher BMI, and a higher grade of necroinflammation.²¹ Thus steatosis is more frequent in HCV genotype 3 than in HCV genotype 1 infected patients. Steatosis is present in 73% of patients infected with genotype 3 and in 50% of patients infected with genotypes other than 3 (table 1).^{10–23} The mechanisms underlying this genotype specific steatosis are unknown. We have recently shown that HCV genotype 3 is associated with higher quasispecies heterogeneity than genotype 1.²⁹ Serum levels of apolipoprotein B and cholesterol are reduced in patients in whom steatosis responds to antiviral therapy.¹⁵ Thus the disappearance of steatosis correlates with normalisation of apolipoprotein B and cholesterol levels. Hypocholesterolaemia in patients with chronic hepatitis C (especially genotype 3) has been reported by others.^{20–30} These observations suggest that HCV may interfere with secretion of very low density lipoprotein (VLDL), which is corrected by antiviral therapy.

A significant correlation was observed between steatosis and high titres of intrahepatic negative strand HCV RNA in patients infected with HCV genotype 3, indicating that steatosis may be secondary to a viral cytopathic effect.¹³ In two studies in patients with HCV genotype 3, an association between grade of steatosis and serum HCV RNA levels was observed.^{14–20} However, other studies have not confirmed this.³¹

Experimental models: demonstration that HCV can induce steatosis

In vitro studies and the transgenic mouse model have both suggested that the HCV core protein is sufficient to induce lipid accumulation in hepatocytes. In cell culture, at least two HCV proteins, core and NS5A, are suspected to interact with the cell machinery involved in lipid metabolism.^{32–33} However, the role of NS5A (if any) seems to be confined to cells coexpressing the HCV core. These proteins interact with apolipoproteins A1 and A2, which are likely involved in triglyceride accumulation and storage in the hepatocytes. Both the core and NS5A proteins are localised on the surface of lipid droplets. Overexpression of core protein further stimulates the formation of lipid droplets.

In transgenic mice it has been reported that HCV core protein inhibits microsomal triglyceride transfer protein (MTP) activity.³⁴ As this is a rate limiting enzyme playing a key role in the VLDL assembly, the direct and likely consequence of its inactivation is accumulation of unsecreted triglycerides, hence steatosis.³⁴ Direct interaction with MTP is unlikely as it would require secretion of the core protein into the endoplasmic reticulum lumen which has not been reported. However, MTP inhibition may still be indirect. One disadvantage of the mouse model is that mice made transgenic with the HCV core protein have normal apolipoprotein B levels in serum.¹⁵ A potential indirect effect of the HCV core on the VLDL secretory pathway may be brought about by the non-specific effect of reactive oxygen species (ROS).³⁵ It has been shown that the HCV core protein causes mitochondrial injury, leading to oxidative stress. These effects were prevented by a mitochondrial electron transport inhibitor.³⁵ Thus it seems that, at least in this model, the core protein of HCV may cause oxidative injury as a result of its localisation and direct toxic effect on mitochondria. Thus HCV transgenic mice have increased sensitivity to oxidative stress, and increased hepatic lipid peroxidation occurs in response to carbon tetrachloride.³⁶ Increased production of ROS may cause peroxidation of membrane lipids and structural proteins (such as those involved in trafficking and secretion).

Thus there is evidence that HCV proteins can cause steatosis in the absence of an immune response. However, all of these models used constructs derived from genotype 1 isolates. For this reason, we have established an in vitro model of expression of the HCV core protein of several different types (from 1 to 5), and our preliminary data show

Table 2 Characteristics associated with the presence of steatosis in patients with chronic hepatitis C

Author, year	Patients (n)	Characteristics associated with steatosis
Mihm 1997 ¹⁰	85	Genotype 3
Czaja 1998 ¹¹	60	BMI, cholesterol, triglycerides
Hourigan 1999 ¹²	148	BMI
Rubbia-Brandt 2000 ¹³	70	Genotype 3, liver HCV RNA (in genotype 3)
Adinolfi 2001 ¹⁴	180	BMI (in genotype 1), genotype 3
Serfaty 2001 ¹⁵	100	Hypobetalipoproteinaemia, genotype 3
Monto 2002 ¹⁶	297	BMI, genotype 3
Westin 2002 ¹⁷	98	BMI, alcohol, genotype 3
Hui 2002 ¹⁸	124	BMI, genotype 3, cholesterol, triglyceride
Castera 2003 ¹⁹	96	BMI, genotype 3
Poynard 2003 ²⁰	1428	Age > 40 y, BMI, triglycerides, genotype 3
Asselah 2003 ²¹	290	BMI, genotype 3
Rubbia-Brandt 2004 ²²	755	BMI, age, alcohol, genotype 3,
Patton 2004 ²³	574	BMI, genotype 3, age, duration of infection, serum HCV RNA (genotype 3)

BMI, body mass index.

that the core proteins from both genotypes 1b and 3a share the capability of inducing accumulation of triglycerides in Huh-7 hepatoma cells, although genotype 3a is much more efficient.³⁷ These results raise the issue of which factors, other than the HCV core protein, may induce steatosis observed in humans. In particular, the role of other viral proteins should be properly investigated.

DOES STEATOSIS AFFECT THE PROGRESSION OF THE DISEASE?

Limitations of published studies

Published data on the influence of steatosis in chronic hepatitis C have several limitations. Firstly, only limited studies are available, almost all being cross sectional or retrospective. Histological criteria for fatty liver disease may be different and the populations studied may have different risk factors for steatosis. For example, in a study from the

USA, mean BMI was 28.5 kg/m² and 32% of patients were obese,¹⁶ whereas in our study (France), mean BMI was 24 kg/m² and only 8% of patients were obese.²¹ Moreover, the HCV genotype distribution differs from region to region. For example, genotype 3 is more frequent in Europe (24%) than in the USA (14%).^{16, 22}

Experimental models: steatosis and fibrosis development

In animal models, such as the leptin deficient ob/ob mouse or the leptin receptor deficient mouse (db/db) and rat (fa/fa), there is very marked hepatic steatosis but little steatohepatitis or fibrosis.³⁸ Development of steatohepatitis in these models depends on additional factors, such as endotoxin exposure, acute liver injury (for example, ischaemia-reperfusion), alcohol, excess dietary polyunsaturated fatty acids, or aging. A unifying hypothesis envisages that all causes capable

Figure 2 Pathogenesis of liver fibrosis in chronic hepatitis C. Proposed mechanisms leading to steatosis and fibrosis in hepatitis C virus (HCV) infection. Steatosis may be caused directly by HCV core protein expression (with or without the cooperation of other viral proteins) but other mechanisms are likely, such as the hyperglycaemic/hyperinsulinaemic state, the hallmark of insulin resistance (where HCV may play an additional direct role through activation of cytokines), and production of reactive oxygen species (ROS) (again, possibly induced by HCV and by insulin resistance). Furthermore, steatosis renders hepatocytes more insulin resistant and, on the other hand, provides the fuel for amplifying the production of ROS. Fatty accumulation, however, is not fibrogenic per se. The production of ROS may provide a pathogenetic link between steatosis and fibrosis. ROS induce cytokines formation by hepatocytes (tumour necrosis factor α (TNF- α), transforming growth factor β (TGF- β), interleukin 8 (IL-8), and FasLigand (FasL)). TGF- β activates collagen synthesis by stellate cells while IL-8 is a potent chemoattractant for human neutrophils. ROS trigger lipid peroxidation, which releases malondialdehyde (MDA) and 4-hydroxynonenal (HNE). MDA and HNE also increase collagen synthesis by stellate cells while HNE is a potent chemoattractant for human neutrophils. Additional fibrogenic stimuli are represented by the fibrogenic inflammatory cytokines (released as part of both innate and adaptive immune responses to HCV infection and antigen expression) and by the direct profibrogenic effect of hyperinsulinaemia.

of changing the redox equilibrium of the hepatocyte may result in liver inflammation and fibrogenesis activation. For example, ROS may promote hepatic stellate cell activation and collagen fibre deposition.³⁹ Lipid peroxidation products, such as 4-hydroxy-2,3-nonenal, may elicit activation and nuclear translocation of c-Jun N-terminal kinases, upregulate c-Jun and increase AP-1 binding, all of which may lead to procollagen type I overexpression.⁴⁰ Furthermore, these phenomena are efficiently prevented by antioxidants.³⁹

Direct involvement of HCV in the development of insulin resistance

A prominent mechanism linking steatosis and fibrogenesis is insulin resistance. The molecular cause of insulin resistance, a major factor in the pathogenesis of type 2 diabetes, is unknown. Several papers have suggested an association between chronic hepatitis C and type 2 diabetes.^{24–28} This is relevant as prolonged hyperglycaemia results in several metabolic changes that are of interest in liver fibrogenesis and recent studies show indeed a close correlation between the degree of insulin resistance and extent of fibrosis.^{41–42}

Recently, it was shown that glucose tolerance was impaired in a mouse model transgenic for the HCV core gene, with plasma glucose levels being higher at all time points, including in the fasting state, although the difference was not statistically significant.⁴³ Transgenic mice exhibited marked insulin resistance, as revealed by the insulin tolerance test, as well as significantly higher basal serum insulin levels. Feeding with a high fat diet led to the development of overt diabetes in transgenic mice but not in control mice. Moreover, there were increased levels of tumour necrosis factor α (TNF- α) in the liver of HCV core gene transgenic mice. High levels of TNF- α have also been observed in human chronic hepatitis C patients.⁴⁴ TNF- α has been shown to induce insulin resistance in experimental animals and cultured cells.^{45–46} Inhibition of tyrosine phosphorylation of insulin receptor substrate 1 and 2 may be one of the mechanisms by which a high level of TNF- α causes insulin resistance.^{46–48} Thus TNF- α was considered to be a predominant cause of insulin resistance in transgenic mice. Administration of an anti-TNF- α antibody restores insulin sensitivity.⁴³ These results provide direct experimental evidence for the contribution of HCV in the development of insulin resistance. There are experimental arguments for a direct role of insulin in fibrosis progression in HCV infection.

Firstly, insulin receptors were identified on hepatic stellate cells isolated from liver sections from patients with chronic HCV infection.⁴⁹ Secondly, in hepatic stellate cells, connective tissue growth factor mRNA and protein were significantly increased following incubation with insulin.⁵⁰

Diabetes per se may lead to increased production of ROS,⁵¹ and secondary upregulation of antioxidant genes, glycosylation of low density lipoprotein and other proteins, as well as the formation of advanced glycation end products (AGE).⁵² Expression of the AGE specific receptor (RAGE) in the liver is restricted to stellate cells, and is upregulated at the time of activation and transition to the myofibroblasts phenotype.⁵³ In genetically obese rodents, characterised by severe steatosis, insulin resistance is promoted by increased activity of IkappaB kinase- β , the activator of nuclear factor κ B (NF κ B).⁵⁴ The latter is a transcription factor that plays a key role in expression of several proinflammation genes. Activation of NF κ B may occur via oxidative stress mediated signalling involving the p38 mitogen activated protein kinase (MAPK).⁵⁵ The link between insulin resistance and steatosis is complex, and the exact sequence of events is unclear, although several hypotheses have been made.⁵⁶ Again, a potential link between severity of steatosis and fibrosis may be secondary to increased formation of reactive oxygen species by steatotic hepatocytes (fig 2).

Human studies: steatosis and fibrosis progression

Factors influencing fibrosis progression in chronic hepatitis C are poorly understood (box 2).⁵⁷ There is some controversy with regard to the influence of steatosis on the progression of fibrosis. Among 14 studies, 10 found an association between the presence of steatosis and a higher degree of fibrosis (table 3). However, in most of these studies this association was of low statistical significance. Furthermore, the histological scores used were different, multivariate analysis was not always performed, and the study populations were different regarding risk factors for steatosis. Thus one possible explanation accounting for the discrepancies is that steatosis may be a cofactor of accelerated fibrosis progression as a function of its pathogenesis (that is, viral v metabolic).

The fact that obesity has been related to the risk of cirrhosis is an argument for a role of metabolic steatosis in the progression of fibrosis. In several studies, among alcoholics and patients with HCV infection, obesity favoured fibrosis.^{58–62} Fibrosis can indeed develop in overweight patients that are

Table 3 Relationship between the presence of steatosis and histological grade and stage of chronic hepatitis C

Author, year	Patients (n)	Histology	
		Inflammation (p)	Fibrosis (p)
Mihm 1997 ¹⁰	85	Yes (ND)	Yes (ND)
Czaja 1998 ¹¹	60	No (ND)	No (ND)
Hourigan 1999 ¹²	148	Yes (ND)	Yes (<0.03)
Rubbia-Brandt 2000 ¹³	70	Yes (<0.001)	Yes (0.02)
Adinolfi 2001 ¹⁴	180	Yes (<0.007)	Yes (<0.001)
Serfaty 2001 ¹⁵	100	No (ND)	Yes (0.008)
Monto 2002 ¹⁶	297	No (0.52)	No (0.26)
Westin 2002 ¹⁷	98	Yes (ND)	Yes (ND)
Hui 2002 ¹⁸	124	No (ND)	No (ND)
Castera 2003 ¹⁹	96	No (ND)	Yes (0.03)
Poynard 2003 ²⁰	1428	No (0.09)	Yes (0.007)
Asselah 2003 ²¹	290	Yes (<0.001)	No (0.35)
Rubbia-Brandt 2004 ²²	755	No (ND)	Yes (<0.001, only in genotype 3)
Patton 2004 ²³	574	No (ND)	Yes (<0.01, only in genotype 1)

ND, not done.

Box 2 Factors associated with more rapid fibrosis progression in chronic hepatitis C

- ▶ Associated
 - Male sex
 - Duration of infection
 - Age at infection (older)
 - Long term excessive alcohol consumption
 - Long term immunosuppression (organ transplantation, bone marrow transplantation)
 - HIV coinfection
 - Overweight
 - High initial fibrosis stage
 - High initial necroinflammation grade
- ▶ Possibly associated
 - Steatosis
 - Insulin resistance
 - High HCV heterogeneity
 - HCV genotype 3
 - Cigarette consumption

devoid of any other causes of liver disease.⁶³ However, in this study, the degree of steatosis was not associated with fibrosis, suggesting that a high BMI could be associated with fibrosis through mechanisms other than steatosis.

At least four studies did not find a link between the presence of steatosis and increased fibrosis (table 3). In our study, steatosis, in multivariate analysis, was associated with HCV genotype 3 infection, higher BMI, and high grade of necroinflammation, but was not associated with high degree of fibrosis.²¹ However, in a recent European study with 755 consecutive chronic hepatitis C patients, steatosis was independently associated with fibrosis, but multivariate analysis on patients divided according to viral genotype (that is, 3 *v* non-3) showed that steatosis was associated with fibrosis only in genotype 3 infected patients.²² It seems that “non-viral steatosis” could be as benign as NAFLD whereas viral steatosis (genotype 3 induced steatosis) may accelerate fibrosis progression. In another recent study from the USA,²³ the opposite was true (that is, some evidence was presented that steatosis may affect fibrosis progression only among patients with genotype 1).

Hepatitis C virus, insulin resistance, and fibrosis

In one interesting study, Hui and colleagues⁴¹ observed that patients with chronic hepatitis C with mild fibrosis (stage 0 or 1) had higher levels of insulin, C peptide, and HOMA-IR (all $p \leq 0.01$) compared with matched healthy controls. In the 250 hepatitis C virus patients (fibrosis stages 0–4), independent predictors of HOMA-IR included higher BMI ($p < 0.001$), previous failed antiviral treatment ($p < 0.001$), portal inflammatory grade ($p < 0.001$), and genotype 3 infection ($p = 0.01$). Genotype 3 had significantly lower HOMA-IR than other genotypes (which were comparable when adjusted for the effects of the remaining independent predictors). HOMA-IR was an independent predictor for the degree of fibrosis ($p < 0.001$) and the rate of fibrosis progression ($p = 0.03$). HCV may induce insulin resistance irrespective of the severity of liver disease, and this effect seems to be genotype specific. Furthermore, their findings support the hypothesis that insulin resistance may contribute to fibrosis progression in chronic HCV infection. Interestingly, insulin resistance may contribute to fibrosis progression in chronic HCV infection independent of the grade of steatosis. Surprisingly, in this

study, HCV genotype 3 had significantly lower insulin resistance than other genotypes.

In another recent study,⁶⁴ marked steatosis was associated with HCV genotype 3 infection and high BMI. Overweight patients had increased circulating insulin compared with lean patients. More advanced fibrosis was associated with obesity and insulin concentrations. Interestingly, in obese patients with chronic HCV infection, circulating insulin levels increased at an early stage of fibrosis (an argument for a causal role). We should remember that in patients with NAFLD, increasing insulin resistance is associated with more severe disease and diabetes is a risk factor for disease progression.⁶⁵ Therefore, in overweight patients with HCV infection, increasing circulating insulin levels may be one factor responsible for the association between increasing BMI and fibrosis.

In a retrospective survey, high serum glucose was found to be associated with advanced fibrosis and higher fibrosis progression rate in chronic hepatitis C patients, and was independent of age at infection or duration of infection.⁶⁶ The profibrogenic impact of glucose intolerance was higher than obesity. High serum glucose has also been independently associated with fibrosis in alcoholic liver disease.⁶⁷

STEATOSIS AND NECROINFLAMMATION

Several studies found a link between necroinflammation and steatosis (table 3). Necroinflammatory activity in chronic hepatitis C may fluctuate over time.⁶⁸

There are several mechanisms which may account for the relationship between steatosis and necroinflammation. In vitro studies have shown that the HCV core protein could lead to oxidative stress.³⁵ Moreover, HCV is associated with increased production of cytokines⁴⁴ that enhance inflammation and lead to increased lipid peroxidation. Furthermore, some patients may have NASH in addition to having chronic hepatitis C. Steatosis could be the consequence of more severe cell injury and necroinflammation rather than the direct cause of worsening fibrosis.

HOW TO RECONCILE ALL OF THESE DATA?

Taking into account all of the published data, it seems that there is an association between steatosis and fibrosis. But does this mean that steatosis is the cause of fibrosis progression? Steatosis could be a marker of necroinflammation and therefore a marker of fibrosis progression as several studies found a significant association between necroinflammation and high stage of fibrosis.^{16 21 69 70} Steatosis could also be a marker of insulin resistance. In patients with HCV infection, insulin resistance may be one factor responsible for both steatosis in one way and increasing fibrosis in another. Interestingly, we reported recently the rate and risk of cirrhosis in 96 patients infected by HCV during the bone marrow transplantation period. At a median follow up of 15.7 years, 15 patients developed biopsy proven cirrhosis leading to a cumulative incidence of cirrhosis of 24% at 20 years. By multivariate analysis, HCV genotype 3 was associated with risk of cirrhosis.⁷¹

DOES TREATMENT IMPROVE STEATOSIS?

We initially reported a patient with HCV genotype 3 infection, recurrent hepatitis after liver transplantation, and massive steatosis, in whom steatosis disappeared when HCV replication was inhibited by treatment and recurred when replication relapsed after treatment withdrawal.^{72 73} This

observation was confirmed by other studies, thus supporting a cytopathic effect of HCV genotype 3. In the first study, Kumar and colleagues⁷⁴ reported that in patients with HCV genotype 1 (n = 28) there was no change in steatosis after treatment, irrespectively of response; however in those infected with genotype 3 (n = 34), a sustained virological response (SVR) resulted in significantly reduced steatosis (p < 0.001) but no change in steatosis among those without an SVR. In another study involving 1428 naive patients, the presence of steatosis was associated with a lower SVR (p < 0.001)²⁰ but in patients with an SVR, steatosis was most improved in those with HCV genotype 3. In genotype 3 responders, baseline low serum cholesterol was corrected by treatment (p < 0.001). These results have been confirmed by others.²³

Interestingly, in a retrospective analysis of 174 patients with chronic hepatitis C, obesity (and not steatosis) was a negative predictor of SVR.⁷⁵ It could be that obesity causes steatosis and that each independently diminishes SVR. Obesity decreases interferon bioavailability and impairs the immune response to HCV. The mechanisms underlying the influence of steatosis on SVR remain to be determined. An intriguing aspect that arises from these studies is the fact that steatosis seems to negatively influence SVR, as suggested by earlier reports.^{76–77} This phenomenon seems to be limited to patients with a non-HCV related type of steatosis as the presence of even severe steatosis in patients with genotype 3 was not necessarily associated with a decreased rate of SVR.²⁰

It is interesting to note that interferon α and insulin share some factors in signalling transduction, such as p38 MAPK⁷⁸; whether insulin resistance translates into a relative resistance to interferon α remains a matter of speculation.

CONCLUSION

The relationship between HCV and steatosis in terms of fibrosis and disease progression is unclear. HCV genotype 3 is clearly associated with steatosis ("viral steatosis"). It seems that in some patients, steatosis may be associated with fibrosis progression. The mechanisms underlying this association are unknown. The challenge for the clinician in care of patients with chronic HCV infection and steatosis is to be able to differentiate between those with pure steatosis alone and a benign course from those with progressive disease (the same challenge in separating NASH from NAFLD).

There remain many unanswered questions and there is a need for large prospective studies on hepatitis C patients, using multivariate analysis, which take into account confounding factors for steatosis, insulin resistance, fibrosis, and response to treatment. A meta-analysis of individual patient data (the HCV MAID Study) is ongoing to investigate the relationship between steatosis, inflammation, and fibrosis in chronic hepatitis C. Moreover, we need to develop experimental models using different HCV genotypes in order to understand the mechanisms leading to steatosis.

ACKNOWLEDGEMENT

Experimental unpublished work by F Negro is supported by the Swiss National Science Foundation grant Nos 3200-63549.00 and 3200BD-103727.

Authors' affiliations

T Asselah, P Marcellin, Service d'Hépatologie and INSERM CRB3, Hôpital Beaujon, Clichy, France
 L Rubbia-Brandt, Service de Pathologie Clinique, Hôpital Universitaire, Genève, Switzerland

F Negro, Services de Pathologie Clinique, de Gastroentérologie et d'Hépatologie, Hôpital Universitaire, Genève, Switzerland

REFERENCES

- Sanyal AJ. AGA technical review on nonalcoholic fatty liver disease. *Gastroenterology* 2002;123:1705–25.
- Clark JM, Brancati FL, Diehl AM. Nonalcoholic fatty liver disease. *Gastroenterology* 2002;122:1649–57.
- Neuschwander Tetri BA, Caldwell SH. Nonalcoholic steatohepatitis: summary of an AASLD single topic conference. *Hepatology* 2003;37:1202–19.
- Chitturi S, Farrell GC. Etiopathogenesis of nonalcoholic steatohepatitis. *Semin Liver Dis* 2001;21:27–41.
- Kwiterovich PO Jr. The metabolic pathways of high-density lipoprotein, low-density lipoprotein, and triglycerides: a current review. *Am J Cardiol* 2000;86:5–10L.
- Marsh JB. Lipoprotein metabolism in obesity and diabetes: insights from stable isotope kinetic studies in humans. *Nutr Rev* 2003;61:363–75.
- Ludwig J, Viggiano TR, McGill DB, et al. Nonalcoholic steatohepatitis: Mayo Clinic experiences with a hitherto unnamed disease. *Mayo Clin Proc* 1980;55:434–8.
- Wanless IR, Lentz JS. Fatty liver hepatitis (steatohepatitis) and obesity: an autopsy study with analysis of risk factors. *Hepatology* 1990;12:1106–10.
- Skelly MM, James PD, Ryder SD. Findings on liver biopsy to investigate abnormal liver function tests in the absence of diagnostic serology. *J Hepatol* 2001;35:195–9.
- Mihm S, Fayyazi A, Hartmann H, et al. Analysis of histopathological manifestations of chronic hepatitis C virus infection with respect to virus genotype. *Hepatology* 1997;25:735–9.
- Czaja AJ, Carpenter HA, Santrach PJ, et al. Host- and disease-specific factors affecting steatosis in chronic hepatitis C. *J Hepatol* 1998;29:198–206.
- Hourigan LF, Macdonald GA, Purdie D, et al. Fibrosis in chronic hepatitis C correlates significantly with body mass index and steatosis. *Hepatology* 1999;29:1215–19.
- Rubbia-Brandt L, Quadri R, Abid K, et al. Hepatocyte steatosis is a cytopathic effect of hepatitis C virus genotype 3. *J Hepatol* 2000;33:106–15.
- Adinolfi LE, Gambardella M, Andreana A, et al. Steatosis accelerates the progression of liver damage of chronic hepatitis C patients and correlates with specific HCV genotype and visceral obesity. *Hepatology* 2001;33:1358–64.
- Serfaty L, Andreani T, Giral P, et al. Virus induced hypobetalipoproteinemia: a possible mechanism for steatosis in chronic hepatitis C. *J Hepatol* 2001;34:428–34.
- Monto A, Alonzo J, Watson JJ, et al. Steatosis in chronic hepatitis C: relative contributions of obesity, diabetes mellitus, and alcohol. *Hepatology* 2002;36:729–36.
- Westin J, Nordlinder H, Lagging M, et al. Steatosis accelerates fibrosis development over time in hepatitis C virus genotype 3 infected patients. *J Hepatol* 2002;37:837–42.
- Hui JM, Kench J, Farrell GC, et al. Genotype-specific mechanisms for hepatic steatosis in chronic hepatitis C infection. *J Gastroenterol Hepatol* 2002;17:873–81.
- Castéra L, Hézode C, Roudot-Thoraval F, et al. Worsening of steatosis is an independent factor of fibrosis progression in untreated patients with chronic hepatitis C and paired liver biopsies. *Gut* 2003;52:288–92.
- Poynard T, Ratziu V, McHutchison J, et al. Effect of treatment with peginterferon or interferon alpha-2b and ribavirin on steatosis in patients infected with hepatitis C. *Hepatology* 2003;38:75–85.
- Asselah T, Boyer N, Guimont MC, et al. Liver fibrosis is not associated with steatosis but with necroinflammation in French patients with chronic hepatitis C. *Gut* 2003;52:1638–43.
- Rubbia-Brandt L, Fabris P, Paganin S, et al. Steatosis affects chronic hepatitis C progression in a genotype-specific way. *Gut* 2004;53:406–12.
- Patton HM, Patel K, Behling C, et al. The impact of steatosis on disease progression and early sustained treatment response in chronic hepatitis C patients. *J Hepatol* 2004;40:484–90.
- Allison ME, Wreghitt T, Palmer CR, et al. Evidence for a link between hepatitis C virus infection and diabetes mellitus in a cirrhotic population. *J Hepatol* 1994;21:1135–9.
- Caronia S, Taylor K, Pagliaro L, et al. Further evidence for an association between non-insulin-dependent diabetes mellitus and chronic hepatitis C virus infection. *Hepatology* 1999;30:1059–63.
- Mason AL, Lau JY, Hoang N, et al. Association of diabetes mellitus and chronic hepatitis C virus infection. *Hepatology* 1999;29:328–33.
- Petit JM, Bour JB, Galland-Jos C, et al. Risk factors for diabetes mellitus and early insulin resistance in chronic hepatitis C. *J Hepatol* 2001;35:279–83.
- Mehta SH, Brancati FL, Strathdee SA, et al. Hepatitis C virus infection and incident type 2 diabetes. *Hepatology* 2003;38:50–56.
- Asselah T, Martinot M, Cazals-Hatem D, et al. Hypervariable region 1 quasiespecies in hepatitis C virus genotypes 1b and 3 infected patients with normal and abnormal alanine aminotransferase levels. *J Viral Hepat* 2002;9:29–35.
- Hofer H, Bankl HC, Wrba F, et al. Hepatocellular fat accumulation and low serum cholesterol in patients infected with HCV-3a. *Am J Gastroenterol* 2002;97:2880–5.
- McGuinness PH, Bishop GA, Painter DM, et al. Intrahepatic hepatitis C RNA levels do not correlate with degree of liver injury in patients with chronic hepatitis C. *Hepatology* 1996;23:676–87.

- 32 Barba G, Harper F, Harada T, *et al.* Hepatitis C virus core protein shows a cytoplasmic localization and associates to cellular lipid storage droplets. *Proc Natl Acad Sci U S A* 1997;**94**:1200–5.
- 33 Shi ST, Polyak SJ, Tu H, *et al.* Hepatitis C virus NS5A colocalizes with the core protein on lipid droplets and interacts with apolipoproteins. *Virology* 2002;**292**:198–210.
- 34 Perlemuter G, Sabile A, Letteron P, *et al.* Hepatitis C virus core protein inhibits microsomal triglyceride transfer protein activity and very low density lipoprotein secretion: a model of viral-related steatosis. *FASEB J* 2002;**16**:185–94.
- 35 Okuda M, Li K, Beard MR, *et al.* Mitochondrial injury, oxidative stress, and antioxidant gene expression are induced by hepatitis C virus core protein. *Gastroenterology* 2002;**122**:366–75.
- 36 Lerat H, Honda M, Beard MR, *et al.* Steatosis and liver cancer in transgenic mice expressing the structural and nonstructural proteins of hepatitis C virus. *Gastroenterology* 2002;**122**:352–65.
- 37 Abid K, Pazienza V, de Gottardi A, *et al.* An in vitro model of hepatitis C virus genotype 3a-associated triglycerides accumulation. *J Hepatol* 2005;**42**:744–51.
- 38 Koteish A, Diehl AM. Animal models of steatosis. *Semin Liver Dis* 2001;**21**:89–104.
- 39 Parola M, Robino G. Oxidative stress-related molecules and liver fibrosis. *J Hepatol* 2001;**35**:297–306.
- 40 Parola M, Robino G, Marra F, *et al.* HNE interacts directly with JNK isoforms in human hepatic stellate cells. *J Clin Invest* 1998;**102**:1942–50.
- 41 Hui JM, Sud A, Farrell GC, *et al.* Insulin resistance is associated with chronic hepatitis C and virus infection fibrosis progression. *Gastroenterology* 2003;**125**:1695–705.
- 42 McCaughan GW, George J. Fibrosis progression in chronic hepatitis C virus infection. *Gut* 2004;**53**:318–21.
- 43 Shintani Y, Fujie H, Miyoshi H, *et al.* Hepatitis C virus infection and diabetes: direct involvement of the virus in the development of insulin resistance. *Gastroenterology* 2004;**126**:840–8.
- 44 Neuman MG, Benhamou JP, Malkiewicz IM, *et al.* Kinetics of serum cytokines reflect changes in the severity of chronic hepatitis C presenting minimal fibrosis. *J Viral Hepat* 2002;**9**:134–40.
- 45 Halse R, Pearson SL, McCormack JG, *et al.* Effects of tumor necrosis factor- α on insulin action in cultured human muscle cells. *Diabetes* 2001;**50**:1102–9.
- 46 Uysal KT, Wiesbrock SM, Marino MW, *et al.* Protection from obesity-induced insulin resistance in mice lacking TNF- α function. *Nature* 1997;**389**:610–14.
- 47 Ozes ON, Akca H, Mayo LD, *et al.* A phosphatidylinositol 3-kinase/Akt/mTOR pathway mediates and PTEN antagonizes tumor necrosis factor inhibition of insulin signaling through insulin receptor substrate-1. *Proc Natl Acad Sci U S A* 2001;**98**:4640–5.
- 48 Aytug S, Reich D, Sapiro LE, *et al.* Impaired IRS-1/PI3-kinase signaling in patients with HCV: a mechanism for increased prevalence of type 2 diabetes. *Hepatology* 2003;**38**:1384–92.
- 49 Paradis V, Dargere D, Vidaud M, *et al.* Expression of connective tissue growth factor in experimental rat and human liver fibrosis. *Hepatology* 1999;**30**:968–76.
- 50 Svegliati-Baroni G, Ridolfi F, Di Sario A, *et al.* Insulin and insulin-like growth factor-1 stimulate proliferation and type I collagen accumulation by human hepatic stellate cells: differential effects on signal transduction pathways. *Hepatology* 1999;**29**:1743–51.
- 51 Marfella R, Quagliari L, Nappo F, *et al.* Acute hyperglycemia induces an oxidative stress in healthy subjects. *J Clin Invest* 2001;**108**:635–6.
- 52 Wolff SP, Jiang ZY, Hunt JV. Protein glycation and oxidative stress in diabetes mellitus and ageing. *Free Radic Biol Med* 1991;**10**:339–52.
- 53 Fehrenbach H, Weiskirchen R, Kasper M, *et al.* Up-regulated expression of the receptor for advanced glycation end products in cultured rat hepatic stellate cells during transdifferentiation to myofibroblasts. *Hepatology* 2001;**34**:943–52.
- 54 Yuan M, Konstantopoulos N, Lee J, *et al.* Reversal of obesity- and diet-induced insulin resistance with salicylates or targeted disruption of Ikk β . *Science* 2001;**293**:1673–7.
- 55 Igarashi M, Wakasaki H, Takahara N, *et al.* Glucose or diabetes activates p38 mitogen-activated protein kinase via different pathways. *J Clin Invest* 1999;**103**:185–95.
- 56 Clark JM, Diehl AM. Hepatic steatosis and type 2 diabetes mellitus. *Curr Diabetes Rep* 2002;**2**:210–15.
- 57 Marcellin P, Asselah T, Boyer N. Fibrosis and disease progression in hepatitis C. *Hepatology* 2002;**36**:47–56.
- 58 Teli MR, Day CP, Burt AD, *et al.* Determinants of progression to cirrhosis or fibrosis in pure alcoholic fatty liver. *Lancet* 1995;**346**:987–90.
- 59 Ithurriaga H, Bunout D, Hirsch S, *et al.* Overweight as a risk factor or a predictive sign of histological liver damage in alcoholics. *Am J Clin Nutr* 1988;**47**:235–8.
- 60 Ortiz V, Berenguer M, Rayón JM, *et al.* Contribution of obesity to hepatitis C-related fibrosis progression. *Am J Gastroenterol* 1997;**9**:2408–14.
- 61 Friedenber F, Pungpapong S, Zaeri N, *et al.* The impact of diabetes and obesity on liver histology in patients with hepatitis C. *Diabetes Obes Metab* 2003;**5**:150–5.
- 62 Hickman IJ, Clouston AD, Macdonald GA, *et al.* Effect of weight reduction on liver histology and biochemistry in patients with chronic hepatitis C. *Gut* 2002;**51**:89–94.
- 63 Ratziu V, Giral P, Charlotte F, *et al.* Liver fibrosis in overweight patients. *Gastroenterology* 2000;**118**:1117–23.
- 64 Hickman IJ, Powell EE, Prins JB, *et al.* In overweight patients with chronic hepatitis C, circulating insulin is associated with hepatic fibrosis: implications for therapy. *J Hepatol* 2003;**39**:1042–8.
- 65 Dixon JB, Bathal PS, O'Brien PE. Nonalcoholic fatty liver disease: predictors of nonalcoholic steatohepatitis and liver fibrosis in the severely obese. *Gastroenterology* 2001;**121**:91–100.
- 66 Ratziu V, Munteanu M, Charlotte F, *et al.* LIDO Study Group. Fibrogenic impact of high serum glucose in chronic hepatitis C. *J Hepatol* 2003;**39**:1049–55.
- 67 Raynard B, Balian A, Fallik D, *et al.* Risk factors of fibrosis in alcohol-induced liver disease. *Hepatology* 2002;**35**:635–8.
- 68 Boyer N, Marcellin P. Pathogenesis, diagnosis and management of hepatitis C. *J Hepatol* 2000;**32**(suppl 1):98–112.
- 69 Ghany MG, Kleiner DE, *et al.* Progression of fibrosis in chronic hepatitis C. *Gastroenterology* 2003;**124**:97–104.
- 70 Ryder SD, Irving WL, Jones DA, *et al.* Trent Hepatitis C Study Group. Progression of hepatic fibrosis in patients with hepatitis C: a prospective repeat liver biopsy study. *Gut* 2004;**53**:451–5.
- 71 Peffault de Latour R, Levy V, Asselah T, *et al.* Long-term outcome of hepatitis C infection after bone marrow transplantation. *Blood* 2004;**103**:1618–24.
- 72 Rubbia-Brandt L, Giostra E, Mentha G, *et al.* Expression of liver steatosis in hepatitis C virus infection and pattern of response to alpha interferon. *J Hepatol* 2001;**35**:307.
- 73 Rubbia-Brandt L, Leandro G, Spahr L, *et al.* Liver steatosis in chronic hepatitis C: a morphological sign suggesting infection with HCV genotype 3. *Histopathology* 2001;**39**:119–24.
- 74 Kumar D, Farrell GC, Fung C, *et al.* Hepatitis C virus genotype 3 is cytopathic to hepatocytes: reversal of hepatic steatosis after sustained therapeutic response. *Hepatology* 2002;**36**:1266–72.
- 75 Bressler BL, Guindi M, Tomlinson G, *et al.* High body mass index is an independent risk factor for nonresponse to antiviral treatment in chronic hepatitis C. *Hepatology* 2003;**38**:639–44.
- 76 Bjoro K, Bell H, Hellum KB, *et al.* Effect of combined interferon-alpha induction therapy and ribavirin on chronic hepatitis C virus infection: a randomized multicentre study. *Scand J Gastroenterol* 2002;**37**:226–32.
- 77 Akuta N, Suzuki F, Tsubota A, *et al.* Efficacy of interferon monotherapy to 394 consecutive naive cases infected with hepatitis C virus genotype 2a in Japan: therapy efficacy as consequence of tripartite interaction of viral, host and interferon treatment-related factors. *J Hepatol* 2002;**37**:831–6.
- 78 Platanias LC. The p38 mitogen-activated protein kinase pathway and its role in interferon signaling. *Pharmacol Ther* 2003;**98**:129–42.