The association between gastro-oesophageal reflux disease and asthma: a systematic review

B D Havemann, C A Henderson, H B El-Serag

This paper is freely available online under the BMJ Journals unlocked scheme, see <http://gut.bmj.com/info/unlocked.dtl>

Gut 2007;56:1654–1664. doi: 10.1136/gut.2007.122465

See end of article for authors' affiliations

Correspondence to: H B El-Serag, Gastroenterology and Health Services Research Sections, Michael E DeBakey VA Medical Center and Baylor College of Medicine, Houston, Texas, USA; hasheme@bcm.tmc.edu

Revised 12 July 2007
Accepted 15 July 2007
Published Online First
6 August 2007

Background and aim: Gastro-oesophageal reflux disease (GORD) has been linked to a number of extra-oesophageal symptoms and disorders, primarily in the respiratory tract. This systematic review aimed to provide an estimate of the strength and direction of the association between GORD and asthma.

Methods: Studies that assessed the prevalence or incidence of GORD in individuals with asthma, or of asthma in individuals with GORD, were identified in Medline and EMBASE via a systematic search strategy.

Results: Twenty-eight studies met the selection criteria. The sample size weighted average prevalence of GORD symptoms in asthma patients was 59.2%, whereas in controls it was 38.1%. In patients with asthma, the average prevalence of abnormal oesophageal pH, oesophagitis and hiatal hernia was 50.9%, 37.3% and 51.2%, respectively. The average prevalence of asthma in individuals with GORD was 4.6%, whereas in controls it was 3.9%. Pooling the odds ratios gave an overall ratio of 5.5 (95% CI 1.9–15.8) for studies reporting the prevalence of GORD symptoms in individuals with asthma, and 2.3 (95% CI 1.8–2.8) for those studies measuring the prevalence of asthma in GORD. One longitudinal study showed a significant association between a diagnosis of asthma and a subsequent diagnosis of GORD (relative risk 1.5; 95% CI 1.2–1.8), whereas the two studies that assessed whether GORD precedes asthma gave inconsistent results. The severity–response relationship was examined in only nine studies, with inconsistent findings.

Conclusions: This systematic review indicates that there is a significant association between GORD and asthma, but a paucity of data on the direction of causality.

Gastro-oesophageal reflux disease (GORD) develops when the reflux of stomach contents into the oesophagus causes chronic troublesome symptoms or complications.¹ The most recognisable symptoms of GORD are heartburn and acid regurgitation, but the reflux of noxious material may have wider-reaching effects. In addition to the well-established oesophageal complications associated with the disease,² GORD is believed to lead to extra-oesophageal symptoms and complications, primarily in the respiratory tract.³ An association between GORD and asthma has been accepted for many years, and has been the focus of numerous studies and reviews.^{4–5} Asthma could arise as a result of acid reflux via two possible mechanisms: damage to the pulmonary tree after direct exposure to acid refluxate (reflux theory); or through bronchial constriction as a result of the stimulation of vagal nerve endings in the oesophagus (reflex theory).⁶ In addition, cough and increased respiratory effort may exacerbate GORD by bringing about an increased pressure gradient across the lower oesophageal sphincter.⁷ This could have particular relevance in patients with hiatus hernia, as gastro-oesophageal junction competence is compromised by hiatus hernia during intra-abdominal pressure increases.⁸

The aim of this systematic review is to provide a realistic estimate of the strength and direction of the association between GORD and asthma in adults. Despite the large number of publications examining the clinical and epidemiological nature of this association, ambiguity remains. For example, estimates of the prevalence of GORD in individuals with asthma vary from 30% to 90%.⁹ A particular challenge is that the prevalence of GORD has been measured in a number of different ways in the literature. First, symptom frequency and/or severity have been used as a measure of disease. This is a patient-focused method that can be used in large population-based surveys, but a definitive symptom

cutoff point for disease has not yet been established. At least weekly heartburn and/or acid regurgitation is known to impair quality of life,¹⁰ and this definition has been used in a recent systematic review,¹¹ which reported that 10–20% of the population in the western world have GORD. Oesophageal pH monitoring is a more objective way of measuring abnormal acid reflux, but its diagnostic accuracy is modest.^{12–13} Endoscopy is an objective way of examining for the presence of oesophagitis, but it cannot distinguish microscopic changes in the oesophageal mucosa that may underlie symptoms in some individuals. Erosive oesophagitis is present in approximately 20–40% of individuals with GORD.^{14–16}

We have therefore chosen to review all of these different methodologies to gain a realistic picture of the association between the two diseases. We examined studies that assess the prevalence or incidence of GORD in individuals with asthma, and the prevalence or incidence of asthma in individuals with GORD. We have employed an epidemiological framework for causality that assesses the strength of association, the consistency of association, the temporal association between GORD and asthma, and finally, the severity–response association between the two diseases.

METHODS

Search strategy

Studies published between 1966 and October 2006 were identified in Medline and EMBASE using the following combinations of search terms: 'asthma and reflux' and 'asthma and (reflux or GER or oesophagitis or hiatal hernia) and (risk or odds or incidence or prevalence)'. There was no language restriction imposed on the search. Articles that potentially

Abbreviations: ATS, American Thoracic Society; GORD, gastro-oesophageal reflux disease

assessed the prevalence or incidence of reflux symptoms, abnormal oesophageal acid exposure, oesophagitis, hiatal hernia or Barrett's oesophagus in adults with asthma, or the prevalence of asthma among adults with reflux symptoms or abnormal acid exposure were selected first based on the title, and then based on the abstract. Translations of relevant non-English language studies were obtained. Two independent investigators conducted the search and data abstraction.

Study selection

We sought to ensure as far as possible that the true prevalence of asthma was recorded. Studies conducted in a primary or secondary care setting were required to define asthma in accordance with American Thoracic Society (ATS) guidelines.¹⁷ Patients were therefore required to have the following: a previous diagnosis of asthma with a history of discrete attacks of wheezing, coughing or dyspnoea, and either an increase in the forced expiratory volume in one second (FEV₁) of 20% from baseline after bronchodilator administration, or a decrease in FEV₁ of 20% after methacholine bronchoprovocation.¹⁷ For population-based surveys or studies that included large administrative datasets, the definition of asthma did not need to meet the ATS guidelines. Studies describing the prevalence of reflux symptoms were required to give a description of the symptoms, including their severity and/or frequency. Studies that monitored oesophageal pH were excluded if the monitoring was performed for less than 24 hours. Studies were also excluded if the study cohorts were composed entirely of asthmatic patients referred for suspected GORD, or if the population source was not defined. Studies were excluded if they had a sample size of less than 50.

Tabulation of results

The full papers from the studies selected based on the content of their abstracts were analysed. A standardised abstraction form, constructed a priori, was used. The following data were collected: sampling frame, study design, sample size, control groups (if any), definition of asthma, definition of reflux symptoms, parameters for interpreting 24 hour pH study results, endoscopic findings, number of patients on medications for asthma and/or GORD, severity of asthma or GORD, and temporal relationships between the development of these conditions.

Analysis

We determined overall prevalence estimates by pooling values from studies meeting the selection criteria and calculating average values weighted by sample size. For the studies reporting reflux symptoms, the average prevalence was calculated both with and without the studies reporting less frequent than weekly heartburn and/or acid regurgitation. Unadjusted odds ratios were pooled from studies that had included a comparison group to give overall estimates of the association between GORD and asthma. Heterogeneity was calculated using the I² test. I² is the percentage of total variation across studies caused by heterogeneity.¹⁸ Severity-response and temporal relationships were also identified and presented. Publication bias was examined by constructing funnel plots of the prevalence values from the included studies, which were tested for asymmetry using Macaskill's test and the test proposed by Peters *et al.*¹⁹

RESULTS

In total, 65 relevant studies were identified, and 28 of these met our inclusion and exclusion criteria. The progression of studies through the search and selection process is illustrated in fig 1, and the number of studies in each subject area is shown in fig 2. Funnel plots indicate the absence of publication bias or a small

study effect among the studies reporting the prevalence of GORD in asthma (Macaskill's $p = 0.2461$, modified Macaskill's $p = 0.80$; fig 3) and the presence of a possible small study effect among the studies of asthma in GORD (Macaskill's $p = 0.002$, Peters' $p = 0.28$; fig 4). One study reported both the incidence of GORD in patients with asthma, and the incidence of asthma in patients with GORD. Several studies reported prevalence estimates for reflux symptoms, abnormal oesophageal pH and endoscopic findings, making the total number of prevalence estimates higher than the total number of studies. Only studies in adults were selected, but a minority of the studies also included some children,^{20–21} and in some studies a lower age limit was not reported.^{22–27} In most of the included studies, the ratio of men to women was reasonably even (32–62% men). In one study, only 12% of the study population was male,²⁸ and in several studies over 90% of the population was male.^{29–33}

Symptoms of GORD in patients with asthma

We identified a total of 22 studies that reported the presence of symptoms of GORD in patients with asthma. Eight studies satisfied our criteria, and are detailed in table 1.^{20–22 28 29 34–36} Fourteen studies were excluded from the analysis, as described in table 2.^{23 24 37–48} Among the included studies, one was based on a large primary care administrative database and seven were performed in secondary care settings (table 1). Most of the studies were cross-sectional ($n = 7$), whereas the database study was a longitudinal cohort study with nested case-control analysis. The pooled sample-size weighted average prevalence of GORD in asthma from the seven cross-sectional studies was 59.2%. The Montreal definition of GORD recommends that moderate heartburn and/or regurgitation at least weekly should be used as a cutoff point for disease in epidemiological studies,¹ and this has been employed in a recent systematic review.¹¹ When we only included those studies that reported the prevalence of at least weekly heartburn and/or acid regurgitation ($n = 5$), the average prevalence of GORD was 58.4%. Three studies reported the prevalence of GORD in controls, with an average prevalence of 38.1%. Pooling the odds ratios from these studies gave an overall odds ratio of 5.45 (95% CI 1.89–15.76). At least 90% of patients with asthma used bronchodilators in the three studies in which this was reported.^{22 29 35} In the cohort study based on a large UK primary care database, the current use of oral or inhaled steroids was associated with a non-significant increased risk of GORD.²⁰

Two studies of symptoms of GORD in patients with asthma warrant particular attention. By far the largest study was the cohort study by Ruigomez and colleagues,²⁰ which measured the occurrence of a new diagnosis of GORD in UK primary care patients. The authors found a significantly higher incidence rate for GORD (eight cases per 1000 person-years; 95% CI 7.0–9.1) in those with a previous diagnosis of asthma than in controls (4.4 cases per 1000 person-years; 95% CI 3.9–5.0), indicating that patients with asthma were 1.8 times more likely to develop GORD than those without asthma.²⁰ The second study of particular interest aimed to avoid bias by using a strictly consecutive recruitment protocol, excluding any patients who were referred to the study by other clinicians because of gastrointestinal symptoms.²⁹ That study found a significantly higher prevalence of symptoms of GORD in patients with asthma compared with controls without asthma (OR 2.4; 95% CI 1.6–3.6).²⁹ Asthma severity was directly related to the age of onset of reflux symptoms.²⁹

Studies that monitored oesophageal pH in patients with asthma

We identified 32 studies in which oesophageal pH monitoring was performed in patients with asthma. Nine studies were

Table 1 Included studies reporting the prevalence of reflux symptoms in individuals with asthma

Reference	Country	Study design	Patient recruitment	Population source	Definition of reflux symptoms	Method of data collection	Prevalence or incidence of reflux symptoms in patients with asthma (%)	Prevalence of reflux symptoms in controls (%)	Odds ratio/relative risk (95% CI)	
									Unadjusted	Adjusted
Field <i>et al</i> 1996 ²⁷	Canada	Cross-sectional	Consecutive	Secondary care, asthma clinic	Heartburn in past week	Questionnaire	49*/109 (45.0%)	7*/68 (10.3%)†	OR 7.1 (3.0 to 17.0)‡	–
Carmona-Sanchez <i>et al</i> 1999 ²⁸	Mexico	Cross-sectional	Consecutive	Secondary care, asthma clinic	Heartburn twice weekly for 3 months	Questionnaire	30/60 (50.0%)	–	–	–
Compte <i>et al</i> 2000 ²⁹	Spain	Cross-sectional	Consecutive	Secondary care, asthma clinic	Any heartburn, acid regurgitation or dysphagia in past 6 weeks	Questionnaire	40/81 (49.4%)	–	–	–
Gatto <i>et al</i> 2000 ³¹	Italy	Cross-sectional	Consecutive	Secondary care, asthma clinic	Heartburn and/or acid regurgitation at least twice weekly	Questionnaire	51/100 (51.0%)	–	–	–
Sontag <i>et al</i> 2004 ³²	USA	Cross-sectional	Consecutive	Secondary care, asthma clinic	Heartburn twice weekly for 1 year	Personal interview	185‡/261 (71.0%)	111‡/218 (51.0%)§	OR 2.4 (1.6 to 3.6)	–
Kiljander and Laitinen, 2004 ³⁵	Finland	Cross-sectional	Random sample of consecutive patients	Secondary care	Heartburn weekly	Self-administered questionnaire	47/90 (52.2%)	–	–	–
Ruigomez <i>et al</i> 2005 ³⁰	UK	Cohort study with nested case-control analysis	Random sample from GPRD	Population-based administrative database	GORD diagnosed by physician	Database review	Incidence 8 per 1000 person-years (7.0–9.1) (219/9712)	Incidence 4.4 per 1000 person-years (3.9–5.0) (241/19334)	–	RR 1.5 (1.2 to 1.8)‡
Shimizu <i>et al</i> 2006 ³³	Japan	Cross-sectional	Not given	Secondary care, asthma clinic	QUEST score of at least 4**	Questionnaire	54/78 (69.2%)	27/150 (18.0%)††	OR 10.3 (5.4 to 19.4)‡	–

CI, Confidence interval; GPRD, General Practice Research Database; OR, odds ratio; QUEST, questionnaire for the diagnosis of reflux disease; RR, relative risk.

*Exact values provided by the author.

†Controls were patients attending a family practice.

‡Calculated based on values given in publication.

§Controls were general medical clinic outpatients without pulmonary disease.

|| Controls were age and sex-matched primary care patients without a diagnosis of asthma or gastro-oesophageal reflux disease (GORD) at baseline.

#Adjusted for age, sex, smoking, previous morbidity and healthcare utilisation.

**QUEST score derived from seven questions about regurgitation and stomach and chest discomfort.

††Controls were outpatients or patients admitted to the hospital because of diseases other than GORD or asthma.

included in the analysis (table 3),^{23–26 28 30 34 35 49} all of which were cross-sectional and had a secondary care setting. The pooled sample-size weighted average prevalence of abnormal oesophageal acid exposure in asthma patients was 50.9%. Only one study included a control group, but that study did not report the prevalence of abnormal acid exposure among the controls.³⁰ Without any measure of the prevalence of abnormal acid exposure among controls, such as patients seen in a clinic other than an asthma clinic, it is not clear from the studies whether the rate of abnormal oesophageal acid exposure is higher than expected among patients with asthma or not. In total, 23 studies were excluded (table 4).^{37 39 41 43 50–68} All but one²⁸ of the nine included studies reported medication use, and bronchodilators were widely used.

The US study by Sontag and co-workers³⁰ found the highest prevalence of abnormal pH over 24 hours, at 81.8%, despite

excluding patients who had a referral for GORD. That study considered more criteria in their analysis than other studies, examining the frequency of reflux episodes, acid contact times, and oesophageal acid clearance times measured using three different methods. Controls had significantly fewer reflux episodes ($p=0.0001$), shorter total acid contact time ($p<0.0001$) and shorter oesophageal clearance times ($p=0.0001$) than patients with asthma. When reported, there were no significant differences in medication use between patients with asthma who had normal and abnormal oesophageal pH.^{26 30 34 49}

Five studies also reported, as sub-analyses, the prevalence of abnormal oesophageal pH particularly in asthma patients without the typical symptoms of GORD of heartburn and/or acid regurgitation, giving an overall prevalence of 10–50%.^{23 24 26 35 49} In one study,²³ significant predictors of abnormal

Table 2 Excluded studies reporting the prevalence of reflux symptoms among individuals with asthma, and reasons for exclusion

Reference	Sample size <50	Population source not reported	Patients referred for suspected GORD	Reflux symptoms and/or frequency not defined	Respiratory symptoms or diagnosis not specific for asthma	ATS criteria for asthma diagnosis not satisfied in secondary care setting	Prevalence of reflux symptoms in patients with asthma (%)
Perrin-Fayolle <i>et al</i> 1980 ³⁷				✓			126/150 (84.0)
Luo 1989 ³⁸						✓	34/55 (61.8)
Harding <i>et al</i> 1999 ³⁹			✓				164/199 (82.4)
Nakase <i>et al</i> 1999 ⁴⁰				✓			43/72 (59.7)
Harding <i>et al</i> 2000 ⁴¹						✓	140/220 (63.6)
Mogica Martinez <i>et al</i> 2001 ⁴²			✓	✓		✓	71/100 (71.0)
Heaney <i>et al</i> 2003 ⁴³				✓			29/73 (39.7)
Al-Asoom <i>et al</i> 2003 ²³				✓			28/50 (56.0)
Liou <i>et al</i> 2003 ⁴⁴				✓			40/149 (26.8)
Bochenska-Marciniak and Gorski, 2004 ⁴⁵				✓			131/208 (63.0)
Carlo <i>et al</i> 2005 ⁴⁶	✓					✓	16/34 (47.1)
Gopal <i>et al</i> 2005 ⁴⁷					✓	✓	52/70 (74.3)
Leggett <i>et al</i> 2005 ²⁴				✓			39/52 (75.0)
Chunlertrith <i>et al</i> 2005 ⁴⁸				✓			86/151 (57.0)

ATS, American Thoracic Society; GORD, gastro-oesophageal reflux disease.

Figure 1 Literature search strategy. GER, gastrooesophageal reflux.

pH were nocturnal asthma symptoms (OR 7.7; 95% CI 1.8–32.7) and hoarseness (OR 6.6; 95% CI 1.8–24.1). The frequency of symptoms was not, however, described by the authors. Another study specifically reported the frequency of night-time asthma symptoms, but found no significant difference between patients with normal and abnormal oesophageal pH.⁴⁹

Studies reporting the results of endoscopy in patients with asthma

We identified 18 studies in which endoscopy was performed in patients with asthma. Six studies satisfied our inclusion criteria (table 5),^{28 31 32 36 40 69} and 12 studies were excluded from the analysis (table 6).^{37 38 42 46 61–63 66 68 70–72} All of the included studies were cross-sectional and based in secondary care. The prevalence of erosive oesophagitis ranged from 27.8% to 47.4%, giving a pooled sample-size weighted average prevalence of 37.3%. The reported prevalence of hiatal hernia among patients with asthma ranged from 37.1% to 61.7%, giving a pooled sample-size weighted average prevalence of 51.2%. Only two

Figure 2 Organisation of articles retrieved from literature searches. GORD, gastro-oesophageal reflux disease.

studies included a control group, and in both studies there was a significant positive association between asthma and erosive oesophagitis or hiatal hernia (table 5).^{28 36}

The level of bronchodilator use was not given in three of the studies.^{28 36 40} In the study by Sontag *et al.*,³² 75% of patients used bronchodilators, and there was no significant difference between the prevalence of oesophagitis in those taking and not taking this medication. Bronchodilator use was similarly high (72%) in the study performed by Avidan and colleagues.³¹ In another study,⁶⁹ although the level of bronchodilator use was not reported directly, the authors did state that drug consumption did not differ between those patients with oesophageal dysfunction (including hiatal hernia, oesophagitis, dysmotility or low lower oesophageal sphincter pressure) and those without. That study also found that frequent wheezing and cough was significantly more common among patients with oesophageal dysfunction than those without, but there were no significant differences in spirometric measurements between the two groups.

Asthma in individuals with GORD

We identified a total of 15 studies that evaluated the presence of asthma in adults with GORD. Eleven studies met our inclusion criteria (table 7)^{20 27 33 73–80} and four studies were excluded (table 8).^{81–84} Of the included studies, nine were cross-sectional and two were cohort studies. Seven studies were general population surveys, three took their data from large administrative databases and one was based in secondary care. Nine studies reported the prevalence of asthma in individuals with GORD, giving an average prevalence of 4.6%. The average prevalence in controls was 3.9%, reported in seven of the studies. When only those studies that reported the prevalence of at least weekly heartburn and/or acid regurgitation were included (n = 4), the average prevalence increased to 12.3%, largely because of the exclusion of a very large database study (n = 101 366), which reported the lowest prevalence of asthma in GORD (4.3%).³³ Overall, seven cross-sectional studies included a control group (table 7). Pooling the unadjusted odds ratios using a random effects model gave an overall odds ratio of 2.27 (95% CI 1.814–2.834; fig 5). The calculated I² was 85%, however, indicating considerable heterogeneity. Six of

Table 3 Included studies in which 24 hour oesophageal pH monitoring in patients with asthma was performed

Reference	Country	Study design	Patient recruitment	Population source	Prevalence of abnormal oesophageal acid exposure in patients with asthma (%)
Sontag <i>et al</i> 1990 ³⁰	USA	Cross-sectional	Consecutive	Secondary care, asthma clinic	85*/104 (81.8)
Suzuki <i>et al</i> 1997 ²⁵	Japan	Cross-sectional	Not reported	Secondary care	42/58 (72.4)
Vincent <i>et al</i> 1997 ²⁶	France	Cross-sectional	Consecutive	Secondary care, asthma clinic	30/94 (31.9)
Kiljander <i>et al</i> 1999 ⁴⁹	Finland	Cross-sectional	Not reported	Secondary care, asthma clinic	57/107 (53.3)
Carmona-Sanchez <i>et al</i> 1999 ²⁸	Mexico	Cross-sectional	Consecutive	Secondary care, asthma clinic	45/60 (75.0)
Compte <i>et al</i> 2000 ³⁴	Spain	Cross-sectional	Consecutive	Secondary care, asthma clinic	12/81 (14.8)
Al-Asoom <i>et al</i> 2003 ²³	Saudi Arabia	Cross-sectional	Consecutive	Secondary care, asthma clinic	22/50 (44.0)
Kiljander and Laitinen 2004 ³⁵	Finland	Cross-sectional	Random sample of consecutive patients	Secondary care	32/90 (35.6)
Leggett <i>et al</i> 2005 ²⁴	UK	Cross-sectional	Not reported	Secondary care, asthma clinic	29/52 (55.8)

*Calculated based on values given in publication.

those studies with controls were population based and were thus considered to be of high generalisability. When only those studies were included in the analysis, the pooled odds ratio was 2.68 (95% CI 1.82–3.96) and the I^2 test gave a value of 81%. Only one study reported medication use, finding no association between the current use of gastrointestinal drugs and the occurrence of asthma.²⁰

There were two cohort analyses that reported incidence estimates for asthma in individuals with GORD, or a complication associated with GORD. One of these, which was a follow-up study from the third US National Health and Nutrition Examination Survey (NHANES III), found that the incidence rate of hospitalisation as a result of asthma in patients who had had previous hospitalisation for hiatal hernia or oesophagitis was 2.6 cases per 1000 person-years, whereas in controls without hiatal hernia or oesophagitis it was 1.0 cases per 1000 person-years.⁷⁴ The other study used a UK primary care administrative database, and found an incidence rate of a new diagnosis of asthma among patients with an existing diagnosis of GORD of 6.0 (95% CI 4.9–7.3) per 1000 person-years.²⁰ In patients without a previous diagnosis of GORD, the incidence rate of asthma was significantly lower at 3.8 cases (95% CI 3.1–4.6) per 1000 person-years.

Figure 3 Funnel plot showing the prevalence of gastro-oesophageal reflux disease in individuals with asthma against sample size. GORD, gastro-oesophageal reflux disease.

Severity–response relationship between GORD and asthma

Only three of the studies evaluating the presence of symptoms of GORD in individuals with asthma considered whether the severity of asthma had an impact on the presence, severity or frequency of GORD symptoms. One study from Italy found that a greater proportion of patients with severe asthma experienced at least twice weekly heartburn and/or acid regurgitation than those with mild or moderate symptoms ($p < 0.03$).²¹ Reflux symptoms were present in 30% of patients with mild asthma, 46% of those with moderate asthma and 70% of those with severe asthma.²¹ In a cross-sectional study from Spain,³⁴ a composite score for GORD was calculated based on the percentage of time that pH was less than four in upright and supine positions and in total, the number of reflux episodes in total, the number of reflux episodes longer than five minutes and the duration of the longest reflux episode. The value of this composite score was similar in patients with mild asthma (median 8.8; range 8.0–22.1), moderate asthma (median 9.5; range 7.9–144.5) and severe asthma (median 10.5; range 8.0–66.6).³⁴

Vincent and colleagues²⁶ found that in patients with GORD, there was a very strong association between the provocative dose of methacholine causing a 20% fall in FEV₁ and the number of oesophageal reflux episodes ($p < 0.001$). There were, however, no other correlations between lung function (FEV₁ or mean expiratory flow) and GORD criteria (percentage of 24-hour period with oesophageal pH < 4, number of reflux episodes or lower oesophageal sphincter pressure).

Only two endoscopy studies examined the association between the severity of asthma and the severity of endoscopic findings. Overall, they indicated that more severe asthma is associated with an increased risk of GORD. In one study from Japan,⁴⁰ patients with intermittent, mildly persistent or moderately persistent asthma had a lower mean endoscopic grade of oesophagitis than patients with severe asthma ($p < 0.05$). In the other study,³⁶ also from Japan, patients with mild asthma were most frequently classified as having no apparent mucosal changes, those with moderate asthma most frequently had minimal changes, and those with severe asthma most frequently had oesophagitis of Los Angeles grade A (mucosal break ≤ 5 mm).

In the studies that examined the presence of asthma in individuals with GORD, the severity of GORD defined by the frequency of reflux symptoms was associated with a higher prevalence of asthma in three studies (fig 6).^{73 78 80} Another study found that there was a higher prevalence of asthma

Table 4 Excluded studies in which oesophageal pH monitoring was performed in patients with asthma, and the reasons for exclusion

Reference	Sample size <50	Population source not reported	Patients referred for suspected GORD	pH Monitoring for <24 hours	ATS criteria for asthma diagnosis not satisfied	Prevalence of abnormal oesophageal acid exposure in patients with asthma (%)
Perrin-Fayolle <i>et al</i> 1980 ³⁷					✓	138/150 (92.0)
Perpina <i>et al</i> 1983 ⁵⁰	✓	✓		✓		9/26 (34.6)
Ducolone <i>et al</i> 1987 ⁵¹	✓	✓		✓		11/21 (52.4)
Ekstrom and Tibbling 1988 ⁵²	✓		✓			18/37 (48.6)
Nagel <i>et al</i> 1988 ⁵³	✓				✓	15/44 (34.1)
Guidicelli <i>et al</i> 1990 ⁵⁴					✓	76/121 (62.8)
DeMeester <i>et al</i> 1990 ⁵⁵					✓	54/77 (70.1)
Larrain <i>et al</i> 1991 ⁵⁶				✓	✓	94/142 (66.2)
Herve <i>et al</i> 1993 ⁵⁷		✓				41/53 (77.4)
Gastal <i>et al</i> 1994 ⁵⁸	✓		✓			12/25 (48.0)
Schnatz <i>et al</i> 1996 ⁵⁹			✓		✓	42/54 (77.8)
Suh <i>et al</i> 1997 ⁶⁰	✓				✓	11/42 (26.2)
Campo <i>et al</i> 1997 ⁶¹	✓		✓			14/17 (82.4)
Dal Negro <i>et al</i> 1999 ⁶²	✓					30/37 (81.1)
Harding <i>et al</i> 1999 ³⁹			✓			128/199 (64.3)
Garcia-Compean <i>et al</i> 2000 ⁶³					✓	34/57 (59.6)
Harding <i>et al</i> 2000 ⁴¹	✓		✓			16/26 (61.5)
Heaney <i>et al</i> 2003 ⁴³					✓	31/54 (57.4)
Oldigs 2004 ⁶⁴			✓		✓	39/63 (61.9)
Paleev <i>et al</i> 2005 ⁶⁵	✓				✓	17/37 (45.9)
Hsu <i>et al</i> 2005 ⁶⁶		✓	✓		✓	29/56 (51.8)
Sami <i>et al</i> 2006 ⁶⁷			✓		✓	25/50 (50.0)
Wong <i>et al</i> 2006 ⁶⁸	✓					17/30 (56.7)

ATS, American Thoracic Society; GORD, gastro-oesophageal reflux disease.

among patients who had GORD and erosive oesophagitis (169/2114, 5.2%) than those who had GORD without erosive oesophagitis (127/2065, 4.3%).⁷⁶

Temporal sequence relationship between GORD and asthma

The temporal relationship between GORD and asthma was investigated in two studies.^{20 74} One of those studies,²⁰ which used the UK General Practice Research Database, found a clear association between the presence of a new diagnosis of asthma and the subsequent development of GORD in a multivariate analysis adjusted for age, sex, smoking, previous morbidity and healthcare utilisation (RR 1.5; 95% CI 1.2–1.8). That study also assessed the likelihood of GORD preceding asthma, showing a

non-significantly increased risk of a new diagnosis of asthma among patients with GORD compared with those in the control cohort without GORD (RR 1.2; 95% CI 0.9–1.6).²⁰ The longitudinal US study that used the NHANES III data⁷⁴ showed an increased likelihood of hospitalisation as a result of asthma in individuals who had previously been hospitalised with oesophagitis or hiatal hernia (RR 2.1; 95% CI 1.1–4.2).

DISCUSSION

The findings of our systematic review support a significant association between GORD and asthma. The pooled prevalence values indicate that the prevalence of symptoms of GORD among individuals with asthma is substantially higher (1.6-fold) than in controls. Similarly, although to a lesser degree, the

Table 5 Included studies reporting endoscopic oesophageal findings in patients with asthma

Reference	Country	Study design	Patient recruitment	Sampling frame	Prevalence of hiatal hernia in patients with asthma	Prevalence of erosive oesophagitis in patients with asthma	Prevalence of Barrett's oesophagus in patients with asthma	Prevalence of erosive oesophagitis in controls	Prevalence of hiatal hernia in controls	Unadjusted odds ratio/relative risk (95% CI)
Kjellen <i>et al</i> 1981 ⁴⁶	Sweden	Cross-sectional	Consecutive	Secondary care, asthma clinic	36/97 (37.1%)	–	–	–	–	–
Sontag <i>et al</i> 1992 ³²	USA	Cross-sectional	Consecutive	Secondary care, asthma clinic	97/186 (52.2%)	73/186 (39.2%)	24/186 (12.9%)	–	–	–
Carmona-Sanchez <i>et al</i> 1999 ²³	Mexico	Cross-sectional	Consecutive	Secondary care, asthma clinic	37/60 (61.7%)	–	–	–	61/180 (33.9%)*	RR 3.13 (1.64 to 6.01)
Avidan <i>et al</i> 2001 ³¹	USA	Cross-sectional	Consecutive	Secondary care, asthma clinic	71/128 (55.5%)	43/128 (33.6%)	–	–	–	–
Nakase <i>et al</i> 1999 ⁴⁰	Japan	Cross-sectional	Consecutive	Secondary care	–	20†/72 (27.8%)	–	–	–	–
Shimizu <i>et al</i> 2006 ²⁴	Japan	Cross-sectional	Consecutive	Secondary care, asthma clinic	–	37/78 (47.4%)	–	6/150 (4.0%)‡	–	OR 21.7 (8.5 to 54.9)†

CI, Confidence interval; OR, odds ratio; RR, relative risk.

*Controls were patients with dyspeptic symptoms, including heartburn and/or acid regurgitation, but without oesophagitis or previous gastric surgery.

†Calculated based on values given in publication.

‡Controls were outpatients who had visited the hospital for a routine health examination.

Figure 4 Funnel plot showing the prevalence of asthma in individuals with gastro-oesophageal reflux disease against sample size. GORD, gastro-oesophageal reflux disease.

average prevalence of asthma in individuals with GORD is also higher than in controls (1.2-fold). The average prevalence of reflux symptoms in adults with asthma was 59%, and the prevalence of erosive oesophagitis was 37%. The average prevalence of GORD diagnosed by pH monitoring was 51%. These values in individuals with asthma are substantially higher than those reported in the general population (10–20% for GORD symptoms, 7–16% for erosive oesophagitis^{11 14 15 85}). None of the studies reporting the prevalence of GORD in asthma were population based, however, and some of this

difference may be because the subjects came from selected primary and secondary care populations.

Although there appears to be a strong association between GORD and asthma, most of the studies included in our analysis were cross-sectional or case-control in design, and therefore could not give a clear indication of the temporal sequence of these conditions, an important criterion for causal associations. The temporal sequence between GORD and asthma was explored in only two studies. The single study that assessed whether primary care patients with asthma were at an increased risk of subsequently developing GORD found a significantly increased incidence of GORD among those patients compared with controls. That and another study reported the likelihood of GORD preceding asthma, with inconsistent findings. Similarly, the severity-response relationship, another criterion for a causal association, was reported in a minority of studies. Results were again inconsistent, but tended towards a positive correlation when the increasing severity of GORD (based on either increasing symptom frequency or the increasing severity of oesophagitis) was associated with an increase in the prevalence of asthma. In several studies, increasingly severe asthma was associated with an increased prevalence of symptoms of GORD or severity of GORD. As a result, the available evidence does not yet clearly indicate whether GORD precedes asthma, or asthma triggers GORD. The recently published Montreal definition of GORD concludes that GORD can be an “aggravating cofactor” in asthma.¹

In addition to statistical association and temporal and severity-response relationships, the controlled introduction or

Figure 5 A forest plot of odds ratios obtained from seven cross-sectional studies that examined the prevalence of asthma among patients with GORD. The point estimate and 95% CI for the pooled odds ratio (represented by the diamond) is 2.26 (1.813–2.834).

Table 6 Excluded studies reporting endoscopic oesophageal findings in patients with asthma, and reasons for exclusion							
Reference	Sample size <50	Population source not reported	Patients referred for suspected GORD	ATS criteria not satisfied	Prevalence of oesophagitis in patients with asthma (%)	Prevalence of hiatal hernia in patients with asthma (%)	Prevalence of Barrett's oesophagus in patients with asthma (%)
Mays 1976 ⁷⁰	✓			✓	13/28 (46.4)	18/28 (64.3)	
Diez Gomez et al 1979 ⁷¹				✓	23/82 (28.0)		
Perrin-Fayolle et al 1980 ³⁷				✓		32/150 (21.3)	
Luo 1989 ³⁸				✓	35/55 (63.6)*		
Campo et al 1997 ⁶¹	✓		✓		11/13 (84.6)		
Dal Negro et al 1999 ⁶²	✓					10/37 (27.0)	
Garcia-Compean et al 2000 ⁶³				✓	6/57 (10.5)	18/57 (31.6)	3/57 (5.3)
Mogica Martinez et al 2001 ⁴²			✓	✓		33/100 (33.0)	
Ben-Noun 2001 ⁷²			✓	✓		19†/141 (13.50)	
Carlo et al 2005 ⁴⁶	✓				7/27 (3.7)		
Hsu et al 2005 ⁶⁶		✓	✓			6/56 (10.7)	
Wong et al 2006 ⁶⁸	✓				10/30 (33.3)		

ATS: American Thoracic Society; GORD: gastro-oesophageal reflux disease.
*Prevalence of oesophagitis, hiatal hernia or Barrett's oesophagus in patients with asthma.

Table 7 Included studies reporting the prevalence or incidence of asthma in individuals with symptoms of gastro-oesophageal reflux disease or evidence of oesophagitis

Reference	Country	Study design	Patient recruitment	Sampling frame	Definition of GORD	Definition of asthma	Method of data collection	Prevalence or incidence of asthma in individuals with GORD (%)	Prevalence or incidence of asthma in controls	OR/RR (95% CI)	Adjusted
El-Serafi and Sonnenberg 1997 ²³	USA	Cross-sectional	All patients in Veterans Affairs system	Administrative database	Physician-diagnosed oesophagitis or stricture	Physician diagnosis	Database review	4314/101 366 (4.3%)	2602/101 366 (2.6%)*	OR 1.5 (1.4 to 1.6)	-
Locke <i>et al</i> 1997 ²³	USA	Cross-sectional	Random	General population	Heartburn and/or acid regurgitation at least weekly	Self-reported	Questionnaire	351/303 (11.6%)	511/642 (7.9%)†	OR 1.5 (1.0 to 2.4)†	OR 1.0 (0.6 to 1.3)§
Ruhl <i>et al</i> 2001 ²⁴	USA	Cohort	Random	General population	Physician-diagnosed oesophagitis or hiatal hernia	Physician diagnosis	Database review	Incidence 2.6 cases per 1000 py (10/537)	Incidence 1.0 per 1000 py (167/6391)‡	-	RR 2.1 (1.1 to 4.2)§
Gislason <i>et al</i> 2002 ²⁵	Iceland, Belgium, Sweden	Cross-sectional	Random	General population	Nocturnal heartburn or belching at least weekly	Self-reported physician diagnosis	Questionnaire	91/101 (8.9%)	84/2096 (4.0%)**	OR 2.3 (1.1 to 4.8)†	OR 2.2 (1.0 to 4.7)††
Jaspersen <i>et al</i> 2003 ²⁶	Germany, Austria, Switzerland	Cross-sectional	Not reported	Patients referred for endoscopy	Erosive oesophagitis on endoscopy	Self-reported (recurrent wheezing)	Questionnaire	169/3245 (5.2%)	-	-	-
Khashabian 2003 ²⁷	Iran	Cross-sectional	Random	General population	Heartburn at least 3 times in the last 2 weeks	Self-reported (asthma or wheezing)	Physician interview	4/70 (5.7%)	-	-	-
Wang <i>et al</i> 2004 ²⁷	China	Cross-sectional	Random	General population	Symptom severity and frequency score of at least 3	Self-reported physician diagnosis	Questionnaire	28/430 (6.5%)	46/2102 (2.2%)‡	OR 3.1 (1.9 to 5.0)†	-
Ruigomez <i>et al</i> 2005 ²⁸	UK	Cohort study with nested case-control analysis	Random	General population	Physician diagnosis	Physician diagnosis	Database review	Incidence 6.0 cases per 1000 py (4.9-7.3) (103/5653)	Incidence 3.8 cases per 1000 py (3.1-4.6)‡‡	-	RR 1.2 (0.9 to 1.6)§§
Cho <i>et al</i> 2005 ²⁸	Korea	Cross-sectional	Random	General population	Heartburn and/or acid regurgitation at least weekly	Self-reported	Questionnaire	131/50 (26.0%)	391/1205 (3.2%)‡	OR 10.5 (5.2 to 21.3)†	OR 2.6 (1.4 to 4.8)
Nordenstedt <i>et al</i> 2006 ²⁹	Norway	Cross-sectional	Random	General population	Severe heartburn and/or acid regurgitation	Self-reported	Questionnaire	420/3153 (13.3%)	2940/40210 (7.3%)‡	OR 1.9 (1.7 to 2.2)†	OR 1.6 (1.4 to 1.9)§§
Rey <i>et al</i> 2006 ³⁰	Spain	Cross-sectional	Random	General population	Heartburn and/or acid regurgitation at least weekly	Self-reported	Questionnaire	301/245 (12.2%)	911/1709 (5.3%)‡	OR 2.5 (1.6 to 3.8)†	OR 1.0 (0.7 to 1.5)***

CI, Confidence interval; GORD, gastro-oesophageal reflux disease; OR, odds ratio; py, person-years; RR, relative risk.

*Controls were patients without a diagnosis of oesophagitis or oesophageal stricture.

†Calculated based on values given in publication.

‡Controls were individuals from the general population who had no symptoms of GORD.

§Adjusted for age, sex, psychosomatic symptom score and each of the remaining atypical symptoms.

|| Controls were individuals from the general population without hiatal hernia, oesophagitis, asthma, chronic bronchitis, emphysema or chronic cough at baseline.

#Adjusted for age, sex, marital status, education, body mass index, physical activity, smoking and diabetes.

††Controls were individuals from the general population without nocturnal symptoms of GORD.

‡‡Adjusted for possible confounders.

§§Controls were age and sex-matched primary care patients without a diagnosis of asthma or GORD at baseline.

||| Adjusted for age, sex, smoking, previous morbidity and healthcare utilisation.

§§§Adjusted for age, sex and remaining atypical symptoms.

||| | Adjusted for age, sex and remaining atypical symptoms.

Adjusted for age, sex, body mass index, smoking and use of asthma medication.

Adjusted for age, sex, coffee consumption, alcohol consumption, smoking, bingeing, psychosomatic symptom score and all other atypical symptoms.

Figure 6 The severity–response relationship between symptoms of gastro-oesophageal reflux disease and asthma. GORD, Gastro-oesophageal reflux disease.

removal of stimuli related to one condition (such as GORD) and the corresponding response (such as the effect on asthma symptoms or pulmonary function) have been used to examine the potential for a causal relationship between the two conditions. This has been investigated using oesophageal acid perfusion testing, but no consistent effect has been found.⁸⁶ GORD treatment in patients with asthma has also had mixed results. A recent systematic review showed minimal improvement of asthma symptoms with GORD therapy, but no improvement in objective pulmonary function indices.⁸⁷ Antireflux therapy does, however, allow a reduction in asthma medication use.⁴ This difference in effect may be related to the fact that pharmacological treatments for GORD change the composition of refluxate without preventing reflux itself, whereas antireflux surgery reduces the number of reflux events. It may be that only some patients are sensitive to acid reflux, or that GORD may affect asthma symptoms but not the pulmonary function measures used in those studies.^{4, 9} For example, increased respiratory effort could be a result of the pain of heartburn causing increased minute ventilation rather than triggering bronchospasm. Even if asthma is GORD related, in some cases there may be resistance to GORD treatment through chronic or irreversible changes.

Our systematic review has both strengths and limitations. We present the most comprehensive systematic review of the epidemiological and clinical literature in this area. In particular, consistent definitions of GORD and asthma provide as true a representation of the prevalence of these diseases as possible. The results of our statistical funnel plot testing argue against the presence of publication bias. We have attempted to minimise selection bias by excluding studies in which investigated asthma patients were referred on the basis of a suspicion of underlying GORD. Most of the studies included in

our review were, however, based in secondary and tertiary referral centres and thus have limited generalisability because they are subject to selection bias. In particular, the results from endoscopic and oesophageal pH monitoring studies may have limited generalisability because a large proportion of eligible patients will not give consent for these invasive and sometimes uncomfortable procedures, especially if the procedure is for the purpose of research only. Most studies had no internal controls and this aspect may have led to an overestimation of the association between GORD and asthma, as patients with difficult-to-control disease or suspicion of another causative factor may be overrepresented in these populations. Although we did not employ formal scoring of the studies in this review based on quality, we did perform a secondary analysis of studies with greater generalisability including only those that were population based and included internal controls. Six studies measuring the prevalence of asthma in individuals with GORD met these generalisability criteria, and the pooled odds ratios from those studies were indicative of a significant positive association between GORD symptoms and asthma.

Although there are a great many studies reporting the prevalence of GORD in individuals with asthma, and vice versa, we found very few population-based studies, and very few studies that considered the temporal sequence relationship between the two diseases. This type of epidemiological research would add to our understanding of the link between GORD and asthma. Prospective studies of individuals with GORD that include long-term follow-up and systematic testing for the incidence of asthma, and vice versa, would be the most valuable strategy. Further studies should also evaluate the severity–response relationship between the two diseases. Ideally, studies should include internal controls and adequate numbers of patients to avoid type 2 errors. They should also document, using validated and reproducible measures, the severity of asthma, GORD and oesophagitis. Age is an important factor in relation to the onset of asthma. Most asthma diagnoses are made in childhood, whereas most ‘difficult to control’ asthma is thought to originate in adult life.⁸⁸ It would be interesting to investigate whether age plays a role in GORD-related asthma.

In conclusion, this systematic review quantifies the prevalence of GORD in individuals with asthma, and asthma in GORD, and so contributes to our understanding of the association between these two diseases. It also highlights that, despite the enormous volume of literature that exists on the subject, there is a shortage of high-quality data. We have identified a clear paucity of data on the direction of the temporal sequence association. Addressing this should be a focus for future epidemiological research in this area.

Competing interests: Declared (the declaration can be viewed on the Gut website at <http://www.gutjnl.com/> supplemental).

Table 8 Excluded studies reporting the prevalence or incidence of asthma in individuals with symptoms of gastro-oesophageal reflux disease, or evidence of oesophagitis, and reasons for exclusion

Reference	Sample size <50	Population source not reported	Patients referred for suspected asthma	Reflux symptoms not defined, or GORD or oesophagitis not diagnosed by physician	ATS criteria for asthma diagnosis not satisfied in secondary care setting	Respiratory symptoms or diagnosis not specific for asthma	Prevalence of asthma in individuals with GORD or oesophagitis (%)
Bagnato <i>et al</i> 2000 ⁸¹	✓						11/30 (36.7)
Riccioni <i>et al</i> 2004 ⁸²	✓						7/22 (31.8)
Roka <i>et al</i> 2005 ⁸³					✓		10/299 (3.3)
Li <i>et al</i> 2006 ⁸⁴				✓			4/200 (2.0)

ATS, American Thoracic Society; GORD, gastro-oesophageal reflux disease.

Authors' affiliations

B D Havemann, H B El-Serag, Sections of Gastroenterology and Health Services Research at the Houston Department of Veterans Affairs Medical Center, and Department of Medicine, Baylor College of Medicine, Houston, Texas, USA

C A Henderson, Research Evaluation Unit, Oxford PharmaGenesis Ltd, Oxford, UK

Funding: This study was funded by a research grant from AstraZeneca R&D Mölndal, Sweden.

REFERENCES

- Vakil N, Veldhuyzen van Zanten S, Kahrilas P, *et al*. The Montreal definition and classification of gastro-esophageal reflux disease (GERD)—a global evidence-based consensus. *Am J Gastroenterol* 2006;**101**:1900–20.
- Shaheen N, Ransohoff DF. Gastroesophageal reflux, Barrett esophagus, and esophageal cancer: scientific review. *JAMA* 2002;**287**:1972–81.
- Hungin AP, Raghunath A, Wiklund I. Beyond heartburn: a review of the spectrum of reflux-induced disease. *Fam Pract* 2005;**22**:591–603.
- Field SK. Gastroesophageal reflux and asthma: are they related? *J Asthma* 1999;**36**:631–44.
- Sontag SJ. Gastroesophageal reflux disease and asthma. *J Clin Gastroenterol* 2000;(Suppl 3):S9–30.
- Castell DO, Schnatz PF. Gastroesophageal reflux disease and asthma. Reflux or reflex? *Chest* 1995;**108**:1186–7.
- Field SK. Asthma and gastroesophageal reflux: another piece in the puzzle? *Chest* 2002;**121**:1024–7.
- Sloan S, Rademaker AW, Kahrilas PJ. Determinants of gastroesophageal junction incompetence: hiatal hernia, lower esophageal sphincter, or both? *Ann Intern Med* 1992;**117**:977–82.
- Field SK. A critical review of the studies of the effects of simulated or real gastroesophageal reflux on pulmonary function in asthmatic adults. *Chest* 1999;**115**:848–56.
- Ronkainen J, Aro P, Storskrubb T, *et al*. Gastro-oesophageal reflux symptoms and health-related quality of life in the adult general population—the Kalixanda study. *Aliment Pharmacol Ther* 2006;**23**:1725–33.
- Dent J, El-Serag HB, Wallander MA, *et al*. Epidemiology of gastro-oesophageal reflux disease: a systematic review. *Gut* 2005;**54**:710–17.
- Vaezi MF, Schroeder PL, Richter JE. Reproducibility of proximal probe pH parameters in 24-hour ambulatory esophageal pH monitoring. *Am J Gastroenterol* 1997;**92**:825–9.
- Bhat YM, McGrath KM, Bielefeldt D. Wireless esophageal pH monitoring: new technique means new questions. *J Clin Gastroenterol* 2006;**40**:116–21.
- Ronkainen J, Aro P, Storskrubb T, *et al*. High prevalence of gastroesophageal reflux symptoms and esophagitis with or without symptoms in the general adult Swedish population: a Kalixanda study report. *Scand J Gastroenterol* 2005;**40**:275–85.
- Zagari RM, Pozzato P, Damian S, *et al*. Relationship between gastroesophageal reflux symptoms and esophagitis in the general population: results from the Loiano-Monghidoro study. *Gastroenterology* 2006;**130**:T1018.
- Thomson AB, Barkun AN, Armstrong D, *et al*. The prevalence of clinically significant endoscopic findings in primary care patients with uninvestigated dyspepsia: the Canadian Adult Dyspepsia Empiric Treatment–Prompt Endoscopy (CADET–PE) study. *Aliment Pharmacol Ther* 2003;**17**:1481–91.
- Cropp GJA, Bernstein IL, Boushey HA. Guidelines for bronchial inhalation challenges with pharmacologic and antigenic agents. *Am Thorac Soc Bull* 1980;**6**:11–19.
- Higgins JP, Thompson SG, Deeks JJ, *et al*. Measuring inconsistency in meta-analyses. *BMJ* 2003;**327**:557–60.
- Peters JL, Sutton AJ, Jones DR, *et al*. Comparison of two methods to detect publication bias in meta-analysis. *JAMA* 2006;**295**:676–80.
- Ruigomez A, Garcia Rodriguez LA, Wallander MA, *et al*. Gastroesophageal reflux disease and asthma: a longitudinal study in UK general practice. *Chest* 2005;**128**:85–93.
- Gatto G, Peri V, Cuttitta G, *et al*. Gastroesophageal reflux symptoms are more frequent in patients with severe asthma. *Gastroenterol Int* 2000;**13**:139–42.
- Field SK, Underwood M, Brant R, *et al*. Prevalence of gastroesophageal reflux symptoms in asthma. *Chest* 1996;**109**:316–22.
- Al-Asoom L, Al-Rubaish AM, El-Munshid HA, *et al*. Gastroesophageal reflux in bronchial asthma patients. A clinical note. *Saudi Med J* 2003;**24**:1364–9.
- Leggett JJ, Johnston BT, Mills M, *et al*. Prevalence of gastroesophageal reflux in difficult asthma: relationship to asthma outcome. *Chest* 2005;**127**:1227–31.
- Suzuki J, Sasaki K, Adachi T, *et al*. [A study of gastroesophageal reflux by 24-hour esophageal pH monitoring in patients with bronchial asthma] (in Japanese). *Nippon Shokakibyo Gakkai Zasshi* 1997;**94**:519–25.
- Vincent D, Cohen-Jonathan AM, Leport J, *et al*. Gastro-oesophageal reflux prevalence and relationship with bronchial reactivity in asthma. *Eur Respir J* 1997;**10**:2255–9.
- Khoshbaten M. Gastro-esophageal reflux disease in northwestern Tabriz, Iran. *Indian J Gastroenterol* 2003;**22**:138–9.
- Carmona-Sanchez R, Valdivinos-Diaz MA, Facha MT, *et al*. [Hiatal hernia in asthmatic patients: prevalence and its association with gastroesophageal reflux] (in Spanish). *Rev Invest Clin* 1999;**51**:215–20.
- Sontag SJ, O'Connell S, Miller TQ, *et al*. Asthmatics have more nocturnal gasping and reflux symptoms than nonasthmatics, and they are related to bedtime eating. *Am J Gastroenterol* 2004;**99**:789–96.
- Sontag SJ, O'Connell S, Khandelwal S, *et al*. Most asthmatics have gastroesophageal reflux with or without bronchodilator therapy. *Gastroenterology* 1990;**99**:613–20.
- Avidan B, Sonnenberg A, Schnell TG, *et al*. Temporal associations between coughing or wheezing and acid reflux in asthmatics. *Gut* 2001;**49**:767–72.
- Sontag SJ, Schnell TG, Miller TQ, *et al*. Prevalence of oesophagitis in asthmatics. *Gut* 1992;**33**:872–6.
- El-Serag HB, Sonnenberg A. Comorbid occurrence of laryngeal or pulmonary disease with esophagitis in United States military veterans. *Gastroenterology* 1997;**113**:755–60.
- Compte I, Garrigues V, Perpina M, *et al*. Prevalence of gastroesophageal reflux in asthma. *J Asthma* 2000;**37**:175–82.
- Kiljander TO, Laitinen JO. The prevalence of gastroesophageal reflux disease in adult asthmatics. *Chest* 2004;**126**:1490–4.
- Shimizu Y, Dobashi K, Kobayashi S, *et al*. High prevalence of gastroesophageal reflux disease with minimal mucosal change in asthmatic patients. *Tohoku J Exp Med* 2006;**209**:329–36.
- Perrin-Fayolle M, Bel A, Kofman J, *et al*. [Asthma and gastro-esophageal reflux. Results of a survey over 150 cases] (in French). *Poumon Coeur* 1980;**36**:225–30.
- Luo JY. A study on the morphologic and dynamics of abnormality of the esophagus and gastroesophageal reflux in patients with asthma. *Zhonghua jie he he hu xi za zhi = Zhonghua jiehe he huxi zazhi (Chinese J Tuberc Respir Dis)* 1989;**12**:135–7, 90.
- Harding SM, Guzzo MR, Richter JE. 24-h Esophageal pH testing in asthmatics: respiratory symptom correlation with esophageal acid events. *Chest* 1999;**115**:654–9.
- Nakase H, Itani T, Mimura J, *et al*. Relationship between asthma and gastro-oesophageal reflux: significance of endoscopic grade of reflux oesophagitis in adult asthmatics. *J Gastroenterol Hepatol* 1999;**14**:715–22.
- Harding SM, Guzzo MR, Richter JE. The prevalence of gastroesophageal reflux in asthma patients without reflux symptoms. *Am J Respir Crit Care Med* 2000;**162**:34–9.
- Magica Martinez MD, Paredes Cruz E, Tenorio Pastrana MA, *et al*. [Prevalence of hiatal hernia and chronic esophagitis in adult asthmatic patients] (in Spanish). *Rev Alerg Mex* 2001;**48**:145–50.
- Heaney LG, Conway E, Kelly C, *et al*. Predictors of therapy resistant asthma: outcome of a systematic evaluation protocol. *Thorax* 2003;**58**:561–6.
- Liou A, Grubb JR, Schechtman KB, *et al*. Causative and contributive factors to asthma severity and patterns of medication use in patients seeking specialized asthma care. *Chest* 2003;**124**:1781–8.
- Bochenska-Marciniak M, Gorski P. [Prevalence of gastroesophageal reflux disease in patients with bronchial asthma] (in Polish). *Pol Merkuriusz Lek* 2004;**16**:527–31.
- Carlo C, Anna F, Alessandra A, *et al*. Asthma and gastroesophageal reflux disease: effect of long-term pantoprazole therapy. *World J Gastroenterol* 2005;**11**:7657–60.
- Gopal B, Singhal P, Gaur SN. Gastroesophageal reflux disease in bronchial asthma and the response to omeprazole. *Asian Pac J Allergy Immunol* 2005;**23**:29–34.
- Chunlerith K, Boonsawat W, Zaeoue U. Prevalence of gastroesophageal reflux symptoms in asthma patients at Srinagarind Hospital. *J Med Assoc Thai* 2005;**88**:668–71.
- Kiljander TO, Salomaa ER, Hietanen EK, *et al*. Gastroesophageal reflux in asthmatics: a double-blind, placebo-controlled crossover study with omeprazole. *Chest* 1999;**116**:1257–64.
- Perpina M, Ponce J, Marco V, *et al*. The prevalence of asymptomatic gastroesophageal reflux in bronchial asthma and in non-asthmatic individuals. *Eur J Respir Dis* 1983;**64**:582–7.
- Ducolone A, Vandevenne A, Jouin H, *et al*. Gastroesophageal reflux in patients with asthma and chronic bronchitis. *Am Rev Respir Dis* 1987;**135**:327–32.
- Ekstrom T, Tibbling L. Gastro-oesophageal reflux and nocturnal asthma. *Eur Respir J* 1988;**1**:636–8.
- Nagel RA, Brown P, Perks WH, *et al*. Ambulatory pH monitoring of gastro-oesophageal reflux in "morning dipper" asthmatics. *BMJ* 1988;**297**:1371–3.
- Giudicelli R, Dupin B, Surpas P, *et al*. [Gastroesophageal reflux and respiratory manifestations: diagnostic approach, therapeutic indications and results] (in French). *Ann Chir* 1990;**44**:552–4.
- DeMeester TR, Bonavina L, Iacone C, *et al*. Chronic respiratory symptoms and occult gastroesophageal reflux. A prospective clinical study and results of surgical therapy. *Ann Surg* 1990;**211**:337–45.
- Larrain A, Carrasco E, Galleguillos F, *et al*. Medical and surgical treatment of nonallergic asthma associated with gastroesophageal reflux. *Chest* 1991;**99**:1330–5.
- Herve P, Escourrou P, Salmeron S, *et al*. [Risk factors for gastroesophageal reflux in chronic asthma] (in French). *Rev Mal Respir* 1993;**10**:527–30.
- Gastal OL, Castell JA, Castell DO. Frequency and site of gastroesophageal reflux in patients with chest symptoms. Studies using proximal and distal pH monitoring. *Chest* 1994;**106**:1793–6.
- Schnatz PF, Castell JA, Castell DO. Pulmonary symptoms associated with gastroesophageal reflux: use of ambulatory pH monitoring to diagnose and to direct therapy. *Am J Gastroenterol* 1996;**91**:1715–18.
- Suh JK, Kwang Ho In, So Ra Lee, *et al*. Gastro-esophageal reflux in asthmatic patients. *Tuberc Respir Dis* 1997;**44**:836–43.
- Campo S, Morini S, Re MA, *et al*. Esophageal dysmotility and gastroesophageal reflux in intrinsic asthma. *Dig Dis Sci* 1997;**42**:1184–8.

- 62 **Dal Negro R**, Pomari C, Micheletto C, *et al*. Prevalence of gastro-oesophageal reflux in asthmatics: an Italian study. *Ital J Gastroenterol Hepatol* 1999;**31**:371–5.
- 63 **Garcia-Compean D**, Gonzalez MV, Galindo G, *et al*. Prevalence of gastroesophageal reflux disease in patients with extraesophageal symptoms referred from otolaryngology, allergy, and cardiology practices: a prospective study. *Dig Dis* 2000;**18**:178–82.
- 64 **Oldigs M**. [pH-Testing—experience in clinical practice in a pulmonary division] (in French). *Pneumologie* 2004;**58**:479–82.
- 65 **Paleev NR**, Isakov VA, Chereiskaia NK, *et al*. [Gastroesophageal reflux disease and respiratory apparatus pathology: the evidence of interrelation and unsolved problems] (in Russian). *Vestn Ross Akad Med Nauk* 2005;3–7.
- 66 **Hsu JY**, Lien HC, Chang CS, *et al*. Abnormal acid reflux in asthmatic patients in a region with low GERD prevalence. *J Gastroenterol* 2005;**40**:11–15.
- 67 **Sami K**, Lilia Z, Samira M, *et al*. Findings of 24-hour esophageal pH monitoring in patients with respiratory and otorhinolaryngologic manifestations of gastroesophageal reflux disease. *Tunisie Medicale* 2006;**84**:165–9.
- 68 **Wong CH**, Chua CJ, Liam CK, *et al*. Gastro-oesophageal reflux disease in “difficult-to-control” asthma: prevalence and response to treatment with acid suppressive therapy. *Aliment Pharmacol Ther* 2006;**23**:1321–7.
- 69 **Kjellen G**, Brundin A, Tibbling L, *et al*. Oesophageal function in asthmatics. *Eur J Respir Dis* 1981;**62**:87–94.
- 70 **Mays EE**. Intrinsic asthma in adults. Association with gastroesophageal reflux. *JAMA* 1976;**236**:2626–8.
- 71 **Diez Gomez ML**, Rubio Sotes M, Ortega Nunez A. Relation of hiatal hernia and respiratory tract pathology. *Revista Clin Esp* 1979;**154**:221–3.
- 72 **Ben-Noun L**. Characteristics of comorbidity in adult asthma. *Public Health Rev* 2001;**29**:49–61.
- 73 **Locke GR III**, Talley NJ, Fett SL, *et al*. Prevalence and clinical spectrum of gastroesophageal reflux: a population-based study in Olmsted County, Minnesota. *Gastroenterology* 1997;**112**:1448–56.
- 74 **Ruhl CE**, Sonnenberg A, Everhart JE. Hospitalization with respiratory disease following hiatal hernia and reflux esophagitis in a prospective, population-based study. *Ann Epidemiol* 2001;**11**:477–83.
- 75 **Gislason T**, Janson C, Vermeire P, *et al*. Respiratory symptoms and nocturnal gastroesophageal reflux: a population-based study of young adults in three European countries. *Chest* 2002;**121**:158–63.
- 76 **Jaspersen D**, Kulig M, Labenz J, *et al*. Prevalence of extra-oesophageal manifestations in gastro-oesophageal reflux disease: an analysis based on the ProGERD Study. *Aliment Pharmacol Ther* 2003;**17**:1515–20.
- 77 **Wang JH**, Luo JY, Dong L, *et al*. Epidemiology of gastroesophageal reflux disease: a general population-based study in Xi'an of Northwest China. *World J Gastroenterol* 2004;**10**:1647–51.
- 78 **Cho YS**, Choi MG, Jeong JJ, *et al*. Prevalence and clinical spectrum of gastroesophageal reflux: a population-based study in Asan-si, Korea. *Am J Gastroenterol* 2005;**100**:747–53.
- 79 **Nordenstedt H**, Nilsson M, Johansson S, *et al*. The relation between gastroesophageal reflux and respiratory symptoms in a population-based study: the Nord-Trondelag health survey. *Chest* 2006;**129**:1051–6.
- 80 **Rey E**, Elola-Olaso CM, Rodriguez-Artalejo F, *et al*. Prevalence of atypical symptoms and their association with typical symptoms of gastroesophageal reflux in Spain. *Eur J Gastroenterol Hepatol* 2006;**18**:969–75.
- 81 **Bagnato GF**, Gulli S, Giacobbe O, *et al*. Bronchial hyperresponsiveness in subjects with gastroesophageal reflux. *Respiration* 2000;**67**:507–9.
- 82 **Ricconi G**, Della Vecchia R, Menna V, *et al*. Prevalence of bronchial asthma in patients with endoscopically-documented esophagitis. *Ann Clin Lab Sci* 2004;**34**:94–8.
- 83 **Roka R**, Rosztoczy A, Izbeki F, *et al*. Prevalence of respiratory symptoms and diseases associated with gastroesophageal reflux disease. *Digestion* 2005;**71**:92–6.
- 84 **Li ZS**, Xu XR, Zou DW, *et al*. [A study of extraesophageal presentations in gastroesophageal reflux disease] (in Chinese). *Zhonghua Nei Ke Za Zhi* 2006;**45**:13–16.
- 85 **Mishima I**, Adachi K, Arima N, *et al*. Prevalence of endoscopically negative and positive gastroesophageal reflux disease in the Japanese. *Scand J Gastroenterol* 2005;**40**:1005–9.
- 86 **Sontag SJ**. Why do the published data fail to clarify the relationship between gastroesophageal reflux and asthma? *Am J Med* 2000;**108**(Suppl 4a):159S–69S.
- 87 **Gibson PG**, Henry RL, Coughlan JL. Gastro-oesophageal reflux treatment for asthma in adults and children. Cochrane Review. Issue 3. Oxford: update Software, 2002. Available: <http://www.cochrane.org/reviews> (accessed 9 September 2004).
- 88 **Barnes N**. Most difficult asthma originates primarily in adult life. *Paediatr Respir Rev* 2006;**7**:141–4.

EDITOR'S QUIZ: GI SNAPSHOT .....

A rare complication of peptic ulcer disease

Robin Spiller, editor

Clinical presentation

A 67-year-old Caucasian retired miner was referred by his general practitioner for an open access gastroscopy to evaluate dyspepsia of 6 months' duration. There were no associated “alarm symptoms”, comorbidities, past history of similar symptoms or gastroscopy. His only drug, ranitidine 150 mg twice daily, recently prescribed by his general practitioner, partially relieved his symptoms. He was an ex-smoker and drank alcohol in moderation. Gastroscopy in July 2003 showed findings in the antral area (fig 1).

Questions

What are the findings in the figure?

What is the diagnosis?

See page 1677 for answers

M Srinivas, P Basumani

Department of Gastroenterology, Rotherham Hospitals NHS Foundation Trust, UK

Correspondence to: Dr P Basumani, Level D, Rotherham General Hospital, Moorgate Road, Rotherham S60 2UD, UK; drsrinivasm@hotmail.com

doi: 10.1136/gut.2006.105304

Figure 1 Endoscopic appearance of gastric antrum at initial gastroscopy.