OC-050

ORIGIN OF INCREASED FECAL SERINE PROTEASE IN PATIENTS WITH IRRITABLE BOWEL SYNDROME AND DIARRHOEA (IBS-D)

doi:10.1136/gut.2011.239301.50

K Garsed,^{1,*} G Singh,¹ D Tooth,² C Lam,¹ R Banwait,¹ M Lingaya,¹ R Spiller¹ ¹Nottingham Digestive Diseases Centre NHR BRU, UK; ²School of Biomedical Sciences, University of Nottingham, Nottingham, UK

Introduction Patients with IBS-D have been reported to have increased faecal serine protease activity which could induce visceral hypersensitivity and increase gut permeability by activating proteinase activated receptor-2 (PAR-2). These proteases could be either endogenous (pancreatic or mast cell) or bacterial in origin.

Aim To define the type and origin of proteases using a proteomic method.

Methods *Study 1:* Serine protease activity in stool supernatant from 36 patients meeting the Rome III criteria for IBS-D and 9 healthy controls was evaluated using a previous published method.¹ Patients completed a one week stool diary and a hospital anxiety and depression score. *Study 2:* Serine proteases from a subset of 8 pooled patient samples were purified using a combination of affinity chromatography and SDS-PAGE. Protease identification using proteolysis and mass-spectrometry was performed. *Study 3:* Assessment of stool amylase and elastase was performed using two commercially available ELISAs.

Results *Study 1:* Serine protease activity was significantly increased in patients, 621 ± 94 versus 211 ± 46 units of trypsin/mg protein in controls, $p = 0.04$. Serine protease activity was significantly correlated with anxiety $r = 0.5$ $p = 0.002$. *Study 2:* The predominant proteases were pancreatic in origin including trypsin, elastase and carboxypeptidase-B. Additionally there was abundant pancreatic amylase. *Study 3:* Stool amylase but not elastase was significantly increased in IBS-D 183 ± 35 103 units/ml versus controls 78 ± 44 . Amylase also correlated with serine protease activity, $r = 0.33$, $p < 0.05$.

Conclusion We confirm previous reports of increased serine proteases in IBS-D but these appear to be of pancreatic origin and associated with increased anxiety. Anxiety may cause rapid transit which reduces degradation by colonic bacteria and increases fecal protease.

Competing interests K. Garsed: None Declared, G. Singh: None Declared, D. Tooth: None Declared, C. Lam: None Declared, R. Banwait: None Declared, M. Lingaya: None Declared, R. Spiller Grant / Research Support from: Norgine, Consultant for: Boehringer, Alberio.

Keywords faecal elastase, faecal amylase, irritable bowel syndrome, serine proteases.

REFERENCE

1. Gecse K, Róka R, Ferrier L, Leveque M, Eutamene H, Cartier C, et al. Increased faecal serine protease activity in diarrhoeic IBS patients: a colonic luminal factor impairing colonic permeability and sensitivity. *Gut* 2008;57:591-9.