A total of 1414 adult patients underwent liver transplantation. 81 (6%) patients were diagnosed with biliary anastomotic strictures during this period.

Results Endoscopic therapy was successful in 61% of patients referred with anastomotic strictures with a median intervention/stent free follow-up of 54 months. Patients required a median of 3 endoscopic procedures and two 24 F balloon dilatations to adequately treat the stricture. Three patients (3%) had recurrent stricture and two of these were successfully retreated with endotherapy. A total of 261 endoscopic procedures were undertaken and complications included mild to moderate pancreatitis (n = 9), Cholangitis (n = 4) and stent migration (n = 2). The rate of pancreatitis per endoscopic procedure was 3%. Severe pancreatitis or death were not reported. Twenty patients underwent biliary reconstruction.

Conclusion We report one of the very few prospective studies looking at the long-term efficacy and safety of endoscopic therapy in the management of anastomotic biliary strictures. Endoscopic balloon dilatation and stenting is a safe and efficacious means of treating biliary anastomotic strictures complicating liver transplantation.

Competing interests None.

REFERENCES

1. Morelli J, Mulcahy HE, Willner IR, Cunningham JT, Draganov P. Long-term outcomes for patients with post-liver transplant anastomotic biliary strictures treated by endoscopic stent placement. *Gastrointest Endosc* 2003;58:374-9.
2. Thuluvath PJ, Pfau PR, Kimmey MB, Ginsberg GG. Biliary complications after liver transplantation: the role of endoscopy. *Endoscopy* 2005;37:857-63.

BSG ENDOSCOPY SECTION SYMPOSIUM AND FREE PAPERS

OC-079

LONG-TERM EFFICACY AND SAFETY OF ENDOTHERAPY FOR MANAGEMENT OF BILIARY ANASTAMOTIC STRICTURES FOLLOWING LIVER TRANSPLANTATION

doi:10.1136/gut.2011.239301.79

S Putta,^{1,*} V Sagar,¹ B Gunson,¹ G Haydon,¹ D Freshwater,² A Holt,² S Pathmakanthan,² J Ferguson² ¹Liver Medicine, Queen Elizabeth Hospital, Birmingham, B15 2TH, UK; ²Queen Elizabeth Hospital, Birmingham, UK

Introduction Biliary anastomotic strictures are among the commonest biliary complications following liver transplantation. Endoscopic therapy including balloon dilatation of stricture and maximal stent placement, is widely used as the preferred means of managing these patients.^{1,2} However, endoscopic therapy is not standardized and there is very limited prospective data to support the long-term efficacy of this approach. The aim of our study is to assess the safety and long-term efficacy of endotherapy in the management of biliary anastomotic strictures complicating liver transplantation.

Methods This was a prospective observational study carried out at the liver transplantation unit, Queen Elizabeth Hospital, Birmingham, United Kingdom, between June 2000 and September 2010. In June 2000, the liver transplant unit in Birmingham introduced a new protocol for managing biliary anastomotic strictures complicating orthotopic liver transplantation. The new protocol entailed a sequence of clearly defined endoscopic interventions including balloon dilatations and stent insertions at regular intervals.