underwent ERCP with sphincterotomy and 9/13 were treated conservatively. In patients undergoing sphincterotomy, there were no procedural complications and all had relief of pain (two patients later solicited further ERCP and sphincterotomy).

Conclusion In this series, secretin-MRCP was valuable in a group of patients with suspected SOD. Most scan findings for this indication are normal, but in some patients an abnormal scan is valuable in giving a positive diagnosis. A subgroup of these may benefit from ERCP with sphincterotomy, while others respond adequately to conservative therapy.

Competing interests None declared.

OC-150 A DECADE OF CHANGE IN THE MANAGEMENT OF SEVERE GASTROINTESTINAL HAEMORRHAGE

doi:10.1136/gutjnl-2012-302514a.150

¹A S Dhadwal,* ¹E Platt, ¹E Morad, ¹A Osborne, ²A Watkinson, ¹S Wajed. ¹Department of Surgery, Royal Devon and Exeter Hospital, Exeter, UK; ²Department of Radiology, Royal Devon and Exeter Hospital, Exeter, UK

Introduction Acute major gastrointestinal bleeding represents a serious and complex clinical challenge requiring a multi-modality approach involving endoscopic, radiological and surgical intervention. The optimal management of this condition has evolved significantly in recent years, and this retrospective study evaluated the changing trends in practice over a 10-year period.

Methods A comprehensive and retrospective analysis of all the patients treated for acute serious gastrointestinal haemorrhage (SGIH), not controllable by endoscopic therapy alone between 2001 and 2011 was performed. This time period overlaps the introduction of a dedicated 24-h interventional radiology service (2009). Baseline demographics, including Rockall and Glasgow Scores, and outcomes from interventional radiological and/or surgical intervention were analysed.

Results Ninety-nine patients in the 10-year period required radiological or surgical intervention for SGIH with a median age of 70 years old (range 22–93). Sixty-seven patients (68%) were treated with surgery alone, with a successful outcome in 63 (94%), and a mortality rate of 6%. Radiological intervention was carried out in 32 patients with a successful outcome for this modality alone in 23 (72%), with one death. Eight patients underwent surgery due to incomplete radiological control, with a successful outcome in 7 (88%), but with one death, giving an overall mortality of 6%. Throughout the 10-year period, there was a sequential decline in the use of primary surgical intervention in favour of a multi-modal radiology-surgery approach.

Conclusion Severe gastro-intestinal haemorrhage in the acute setting, when not amenable to successful endoscopic therapy remains a serious condition with significant mortality (6%). The introduction of interventional radiological techniques however now means that the majority of patients can be successfully managed without operative surgery, but this still plays an important role for complex or refractory patients.

Competing interests None declared.

OC-151 PREVALENCE OF, AND PREDICTORS OF, A POSITIVE SEHCAT SCAN FOR BILE ACID DIARRHOEA IN OUTPATIENTS WITH CHRONIC DIARRHOEA

doi:10.1136/gutjnl-2012-302514a.151

¹D J Gracie,* ²F Chowdhury, ¹J Kane, ¹S Mumtaz, ^{1,3}A C Ford. ¹Leeds Gastroenterology Institute, Leeds, UK; ²Department of Nuclear Medicine, St. James's

University Hospital, Leeds, UK; ³Leeds Institute of Molecular Medicine, Leeds University, Leeds, UK

Introduction SeHCAT scanning is not commonly performed in patients with chronic diarrhoea, despite British Society of Gastroenterology guidelines advocating its use. Greater awareness of bile acid diarrhoea (BAD) may increase uptake of testing. We aimed to assess the prevalence of BAD in consecutive patients with chronic diarrhoea undergoing SeHCAT scanning, and to identify factors predicting a positive result.

Methods A retrospective review was undertaken over 6 years at Leeds Teaching Hospitals. A standardised scanning protocol was adopted. Medical records were reviewed to obtain information regarding previous cholecystectomy, terminal ileal (TI) Crohn's disease (CD), TI resection or right hemicolectomy for CD, right hemicolectomy for other reasons, radiation exposure, enteric infection and microscopic colitis. BAD was defined as present when SeHCAT retention was <15% (mild <15%, moderate <10%, and severe <5%), and classified according to underlying aetiology (type I secondary to TI resection or right hemicolectomy, type II idiopathic, and type III secondary to all other causes). Presence of bloating and abdominal pain were used to define IBS.

Results 163 (51.4%) of 317 patients had some degree of BAD. A greater proportion of patients with a positive test had undergone cholecystectomy (29.4% vs 13.9%, $p=0.001$), TI resection for CD (17.8% vs 1.9%, $p<0.001$) or TI resection or right hemicolectomy for other reasons (8.6% vs 1.9%, $p=0.009$). Radiotherapy, enteric infection, or microscopic colitis were not associated with BAD. 54 (37%) of 145 individuals with no obvious risk factors had some degree of BAD, with 17 (32.1%) having severe BAD. Fewer individuals with BAD reported bloating (16.0% vs 25.0%, $p=0.05$), and fewer met criteria for IBS (40.5% vs 52.6%, $p=0.03$). More patients with severe disease had previous right hemicolectomy or TI resection for CD (32.1%, $p<0.001$). A greater proportion of those with mild BAD fulfilled criteria for IBS, compared with those with moderate or severe BAD (65.9% vs 46.3% and 24.7%, $p<0.001$).

Conclusion BAD is common in patients with chronic diarrhoea. Prior surgery is a strong predictor of a positive test, and may negate need for testing. However, among those with no proposed risk factors, yield of testing was almost 40%. The prevalence of BAD in patients with chronic diarrhoea is high, and SeHCAT scanning is a worthwhile diagnostic strategy.

Competing interests None declared.

OC-152 "EXPECTING THE UNEXPECTED": A REVIEW OF EXTRACOLONIC FINDINGS FOUND AT CT COLONOGRAPHY

doi:10.1136/gutjnl-2012-302514a.152

¹H M Owen, ¹R Foulkes,* ¹P Billings, ¹P Chandran, ²C Corr. ¹Department of Colorectal Surgery, Wrexham Maelor Hospital, Wrexham, UK; ²Department of Radiology, Wrexham Maelor Hospital, Wrexham, UK

Introduction The aim of this paper is to report the high prevalence of important extracolonic findings, including cancer, at CT colonography.

Methods Using the PACS system all CT colonograms performed for symptomatic indications between December 2006 and June 2011 were retrieved as part of our ongoing audit. Extracolonic findings were identified and analysed. They were categorised into extracolonic malignancies, benign and important benign findings which were findings that required further investigation or management.

Results 830 patients underwent CT Colonography during this time period (518 females, 313 males, average age 74). 85 colonic cancers were identified and 103 patients had colonic polyps with or without