Introduction It is proposed that the presence of oesophageal mucosa dilated intercellular spaces (DIS) underlies heartburn in patients with NERD. However, 20% of asymptomatic subjects display DIS in distal oesophageal biopsies. Furthermore, oesophageal experimental acidification in healthy volunteers provokes DIS, but often without heartburn despite ongoing acid perfusion. These observations suggest that mucosal impairment above and beyond DIS may be relevant in heartburn perception. Functional mucosal integrity in response to acid challenge can be tested in oesophageal mucosal biopsies “in vitro” by measuring changes in transepithelial electrical resistance (TER).

We aimed to further assess the relationship between presence of DIS, changes in functional oesophageal mucosal integrity (TER) and heartburn perception.

Methods We took distal oesophageal biopsies from patients with and without heartburn. Histological examination for epithelial intercellular space diameter (ISD) was done using light microscopy by taking 50 random measurements at several levels from the basal epithelial layers for each biopsy. DIS was declared at above the 95% CI for normal values as previously published ($> 0.72 \mu\text{m}$).

We identified 11 subjects with DIS (4 with predominant daily troublesome heartburn, 7 with predominant dyspepsia and no heartburn). Biopsies were placed in mini-Ussing chambers, and baseline TER was measured. The luminal aspect of the biopsy was then exposed for 30min to an acidic solution (pH2 + 1 mg/ml pepsin + 1 mM taurodeoxycholate). During exposures, % changes in TER relative to baseline were analysed.

Results

The mean ISD in all subjects was $0.94 \mu\text{m}$ (range $0.74\text{--}1.18 \mu\text{m}$). There was no significant difference in ISD (0.95 vs. $0.91 \mu\text{m}$) or baseline TER (144 vs. $165 \Omega\cdot\text{cm}^2$) between predominant heartburn and predominant dyspepsia groups.

Despite the pre-existing DIS in all subjects, 30 minutes acid exposure was able to induce a further reduction in TER (-23.0% change from baseline, $p < 0.01$).

The reduction in TER was greater in subjects with predominant reflux vs. those with predominant dyspepsia ($-40.4 \pm 10.3\%$ vs. $-11.2 \pm 3.7\%$, $p = 0.01$).

Conclusion In subjects with pre-existing DIS with and without heartburn, trans-epithelial electrical resistance can be further impaired with in vitro acid exposure, suggesting that the mechanism for acid-induced mucosal integrity impairment is not limited to DIS. Furthermore, acid-induced functional integrity impairment was greater in patients with heartburn. This difference in mucosal behaviour in the presence of acid suggests that other mechanisms beyond DIS might be needed to further stimulate afferent nerves in heartburn generation.

Disclosure of Interest None Declared

PTU-163 PROXIMAL OESOPHAGEAL FUNCTIONAL MUCOSAL INTEGRITY DURING DISTAL OESOPHAGEAL ACIDIFICATION IN PATIENTS WITH NERD AND FH

doi:10.1136/gutjnl-2013-304907.253

¹P Woodland, ¹M Al-Zinaty, ¹E Yazaki, ¹J Jafari, ¹D Sifrim. *Barts & The London School of Medicine, Queen Mary University of London, London, UK*

Introduction Although macroscopically normal, the oesophageal mucosa in patients with non-erosive reflux disease (NERD) displays epithelial dilated intercellular spaces (DIS). Such loss of barrier integrity is thought to be important in disease pathogenesis. Patients with NERD have DIS in both the distal and proximal oesophagus. Patients with functional heartburn (FH) do not have DIS. In healthy volunteers acid perfusion of the distal oesophagus induces DIS not only in the exposed distal, but also the unexposed proximal oesophagus. It is unknown what impact distal acid perfusion has in the proximal oesophageal mucosa in symptomatic

patients. Oesophageal impedance measurements can be an “in vivo” surrogate marker of functional mucosal integrity.

We aimed to assess with impedance the proximal oesophageal mucosal integrity during distal acid perfusion in symptomatic patients with NERD (+ve DIS) and FH (-ve DIS).

Methods We studied 22 patients (mean age 48, range 20–75) with typical reflux symptoms and negative endoscopy. 24-hour MII-pH monitoring identified 11 patients with NERD and 11 with FH. A pH-impedance catheter was inserted transnasally such that the proximal impedance segment was 15cm above the LOS. A 10 minute perfusion (10ml/min) of a neutral (pH7), then acidic (pH1) solution was performed in the distal oesophagus (10 cm from LOS) whilst simultaneously measuring proximal impedance. The absence of proximal fluid contamination was suggested by a lack of impedance drop during the neutral perfusion. Baseline impedance, and percentage change from baseline during the perfusion was calculated.

Results Distal perfusion of neutral solution caused no fall in proximal impedance (mean impedance $107 \pm 2.8\%$ of baseline). Distal perfusion of acidic solution caused a significant reduction in proximal impedance (mean impedance $80.3 \pm 5.3\%$ of baseline, $p < 0.01$). The drop of proximal impedance started at a mean of 175 seconds after commencement of acid perfusion. Proximal impedance did increase to a mean of 90% of baseline until 30 minutes after cessation of the acid perfusion. There was no significant difference in baseline proximal impedance ($3165 \pm 216\Omega$ vs. $3116 \pm 232\Omega$) or acid-induced change in proximal impedance ($79.4 \pm 8.1\%$ vs. $81.2 \pm 7.1\%$ of baseline) between FH and NERD patients respectively.

Conclusion In keeping with morphological studies of healthy volunteers, distal oesophageal acidification modifies mucosal functional integrity of the proximal, unexposed oesophagus. This appears to occur regardless of the presence of pre-existing proximal DIS, suggesting that mechanisms of functional mucosal integrity as measured by impedance are beyond DIS formation.

Disclosure of Interest None Declared.

PTU-164 COST EFFECTIVENESS OF AN ER DOMINANT APPROACH IN THE MANAGEMENT OF HIGH GRADE INTRAEPITHELIAL NEOPLASIA AND MUCOSAL CANCER IN BARRETT'S OESOPHAGUS

doi:10.1136/gutjnl-2013-304907.254

¹P J Basford, ¹G Longcroft-Wheaton, ¹R Mead, ¹P Bhandari. *Gastroenterology, Portsmouth Hospitals NHS Trust, Portsmouth, UK*

Introduction Endoscopic resection (ER) is an established effective treatment for high grade intraepithelial neoplasia (HGIN) and intramucosal cancer (IMC) arising in Barrett's oesophagus. ER can lead to recurrence so it is suggested that all patients should undergo radiofrequency ablation (RFA) after ER as a complimentary treatment strategy. However no comparative study to support this concept has been performed. We aimed to compare the cost-effectiveness of an EMR-dominant approach vs an EMR-RFA approach for the treatment of HGIN and IMC in Barrett's oesophagus.

Methods All ER procedures between 2005 and 2012 were recorded in a prospective database which was analysed. Demographic data, histology, procedure success, long-term outcome and complications were assessed. Costs were calculated using NHS HRG codes plus equipment costs for ER and RFA.

Results 92 patients were treated for dysplastic Barrett's oesophagus or early Barrett's cancer by ER. The mean age at first procedure was 69 years and 87% of the patients were male. 21 of 92 patients had advanced histological features on the initial ER specimen and were referred for surgical or oncological treatment. Of the remaining 71 cases, 63 have follow-up data with a mean duration of 4.3 years. 59 of 63 cases (94%) had successful eradication of HGIN/IMC by ER. The remaining 4 patients were referred for surgery for advanced disease (3) or extensive bulky disease not amenable to ER

(1). ER was successful in a mean of 1.46 procedures per patient (range 1–3). Complication rate was 5.2% (4 bleeds, 1 microperforation, 2 strictures). Additional RFA was used in 11 cases. 12 (20%) of patients developed recurrence of HGIN/IMC during follow-up requiring further endoscopic therapy. 2 (3.4%) patients developed more advanced Barrett's neoplasia during follow-up. The calculated cost per patient of an ER-dominant approach is £4125 compared to £8868 per patient for an RFA dominant approach.

Conclusion ER acted as an accurate and safe staging procedure in up to 23% of cases found to have advanced histology. ER is an effective and safe treatment for HGIN/IMC within Barrett's oesophagus without the need for routine RFA and can be performed successfully in a UK centre. However the recurrence of HGIN/IMC is not uncommon and therefore close follow-up is required to identify and treat it at an early stage. An ER-dominant approach may offer significant cost-savings compared to an RFA-dominant approach without compromising overall outcomes.

Disclosure of Interest None Declared

PTU-165 THE BETTER DEFINITION OF NODAL STAGING IN THE 7TH EDITION OF TNM MANUAL DOES NOT PREDICT SURVIVAL OR TRANSLATES INTO BETTER PROGNOSTICATING ABILITY IN OESOPHAGO-GASTRIC JUNCTIONAL ADENOCARCINOMA

doi:10.1136/gutjnl-2013-304907.255

¹R Y Kannan, ¹M Davies, ¹C Jenkins, ¹A Rasheed. ¹Institute of Minimal Access Surgery, Royal Gwent Hospital, Newport, UK

Introduction The 7th TNM staging defines a minimum number of nodes, recommends an optimal number for each T stage, emphasises the prognostic importance of number of regional nodes involved and upstages based on the number of metastatic lymph nodes. We intend on studying the impact of application of 7th TNM rules on nodal staging (N) of resected and pathologically reported oesophago-gastric junctional (OGJ) adenocarcinomas during the last 10 years stratifying them according to the 7th edition TNM staging and to compare against the original staging and assess possible impact of nodal neo-staging on survival.

Methods A retrospective database was used to capture the clinicopathological data of all consecutive curative resections of OGJ adenocarcinomas over the last 10 years in two UK Upper GI Units. Any report with less than 12 lymph nodes was considered inadequate and denoted as (Nx). All cases were re-reported and re-staged according to the 7th TNM staging rules. We compared the impact of the 7th TNM staging rules on neo-staging. Overall survival was analysed using the 6th and 7th TNM staging respectively. Overall survival was sub-stratified into 2 years, 5 years and 10 years post curative resection.

Results Fifty seven (57) pathology reports confirming OGJ adenocarcinomas were reviewed. Adequate lymphadenectomy (minimum of 12 nodes) was noted in 33 patients. Overall stage migration was noted in 36 (63%) reports with the 7th TNM staging. Of those who had adequate lymphadenectomy (33), 20 reports (60.6%) had stage migration. Survival was calculated from the time of initial surgery. Two year survival was assessed in the whole group (n = 57). Five year survival for patients operated between 2000 to 2007 (n = 34) and 10 year survival for those operated on between 2000 to 2002 (n = 10). For stage 3b and stage 3c (7th TNM) there was a 12.5%, 8.9% and 8.9% higher survival rate respectively (for 2.5 and 10 years), compared to the original 6th TNM staging for stage 3. Correspondingly for stage 1b, the survival rate was 5.3%, 3.6% and 3.6% respectively.

Conclusion The 7th edition of TNM staging provides a detailed documentation of the lymphatic staging. This better defined lymphatic staging does not seem to predict survival or have a superior prognosticating ability.

Disclosure of Interest None Declared

PTU-166 DIAGNOSIS OF GASTRO-OESOPHAGEAL REFLUX DISEASE (GORD) BY HISTOLOGY OF MUCOSAL BIOPSIES FROM DISTAL OESOPHAGUS: AGREEMENT WITH PROLONGED PH MONITORING

doi:10.1136/gutjnl-2013-304907.256

¹R Sweis, ²F Chang, ³M Fox, ⁴A Anggiansah, ¹A Lee, ¹A Valdes, ⁵T Wong. ¹Oesophageal lab; ²Histopathology Department, St Thomas' Hospital, London; ³NIHR Biomedical Research Unit and Digestive Diseases Centre, Nottingham University Hospitals, Nottingham; ⁴Oesophageal lab; ⁵Gastroenterology Department, St Thomas' Hospital, London, UK

Introduction Prolonged wireless pH-monitoring (Bravo) increases diagnostic yield compared to 24 hr pH-studies. Studies have shown a weak association between oesophageal acid exposure from 24 hr pH-studies and mucosal disease on distal oesophageal histology. This study assessed the association between Bravo, endoscopic findings and distal oesophageal histology.

Methods From July 2009 to August 2010, 63 consecutive patients with typical reflux symptoms had endoscopy with biopsies taken from 3 & 9 o'clock position at and 2cm proximal to the Z-line prior to pH capsule fixation 6cm proximal to the Z-line for prolonged (up to 96 hr) Bravo. All biopsies were assessed by the 6-parameter Zentilin histology score (Zentilin *et al* Gastroenterol 2005). GORD diagnosis was based on "Average" acid exposure (Total Reflux; TR > 5.3% time pH < 4) over the time period measured and/or symptom-association (symptom index; SI > 50%)

Results Adequate biopsy samples were available from 57/63 patients (mean age 44 (range 17–78); 27M). 37/57 (65%) patients had GORD based on either TR (n = 30) or SI (n = 32). 20/57 (35%) were both TR & SI negative (Functional Heartburn; FH). 18 FH patients had no mucosal changes, 2 had grade A oesophagitis. There was no difference in individual histology parameters between GORD vs. FH (p > 0.05) apart from increased 'intra-epithelial neutrophils' (IEN) at the Z-line (9/37 vs. 1/20 positive; p = 0.031) and 2cm proximally (6/37 vs. 0/20 positive; p = 0.012). The combined Zentilin histology score was also higher in GORD at the Z-line (p = 0.079) and 2 cm proximally (p = 0.05).

Using GORD diagnosis from 96 hr Bravo as reference, ROC analysis revealed that, although sensitivity remained poor, specificity of GORD diagnosis based on histology improved with IEN and total histology score. With increased IEN, sensitivity was 30% at the Z-line and 20% 2 cm above while specificity was 92.6% at the Z-line and 100% 2cm proximally. For the optimal Zentilin histology score of ≥7, sensitivity was 40.5% at the Z-line and 18.9% 2cm above while specificity was 95% at the Z-line and 100% 2 cm proximally.

Histology corroborated GORD diagnosis (based on positive TR) in 11/30 and 20/30 patients at the Z-line and in 6/30 and 11/30 patients 2 cm proximally.

Conclusion Histology lacks sensitivity as a stand-alone diagnostic test; however high IEN or total histology scores have high specificity for GORD diagnosis based on pH-study results. Thus, routine biopsy of the distal oesophagus may be sufficient to diagnose GORD and obviate the need for ambulatory pH-studies in this subgroup of patients.

Disclosure of Interest None Declared.

PTU-167 DIFFERENCES IN BARRETT'S OESOPHAGUS IN AN ETHNICALLY DIVERSE SOUTH LONDON POPULATION

doi:10.1136/gutjnl-2013-304907.257

¹R Hewett, ¹J-Y Kang, ¹A Poullis. ¹Gastroenterology, St George's, University of London, London, UK

Introduction Barrett's oesophagus (BO) is a metaplastic change of the lining of the oesophagus; the normal squamous epithelium is replaced by specialised columnar epithelium. BO is a risk factor for