Effect of a synthetic gastrin-like pentapeptide upon the intestinal transport of sodium, potassium, and water¹

J. C. GINGELL, M. W. DAVIES, AND R. SHIELDS

From the Department of Surgery, The Royal Infirmary, Cardiff

In addition to its marked stimulation of the stomach to secrete acid, the antral hormone, gastrin, can affect other functions of the alimentary tract. Thus, the highly active preparation of gastrin extracted from hog antral mucosa has been shown to stimulate the motility of the stomach and jejunum, to increase the tone of the gall bladder, and to augment the flow of pancreatic juice and hepatic bile (Gregory and Tracy, 1964).

The entire range of physiological activity displayed by natural gastrin is possessed by the C-terminal tetrapeptide-amide, tryptophan, methionine, aspartic acid, and phenylalanine NH₂ (Tracy and Gregory, 1964). The synthetic analogue, which has been produced by the Imperial Chemical Industries (I.C.I. 50,123) and has t-butyloxycarbonyl- β -alanine added to the functional tetrapeptide sequence, seems to have the properties of natural gastrin although it is probably less potent on a molar basis (Konturek and Grossman, 1966; Wormsley, Mahoney, and Ng, 1966; Logan and Connell, 1966). Because one of the important functions of the gastrointestinal tract is the absorption of water and electrolytes, we decided to study the effect of the gastrin-like pentapeptide upon the intestinal handling of sodium, potassium, and water.

METHODS

The experiments were performed in trained, placid mongrels, weighing 10-12 kg., in whom segments of jejunum (two dogs), ileum (two dogs), and colon (one dog) were isolated as Thiry-Vella fistulas. The serosal surface area of the intestinal segments measured approximately 100 cm.². The details of the animal preparation and of the technique of measuring the rates of net transport and unidirectional fluxes of sodium, potassium, and water have been given elsewhere (Shields, Mulholland, and Elmslie, 1966).

¹A preliminary report of this work was read to the Surgical Research Society (Gingell, Davies, and Shields, 1966), and it was carried out in laboratories which were built from funds given by the Wellcome Trustees.

Briefly, the isolated intestinal segments were first rinsed with modified Tyrode's solution (Code and McIntire, 1956) at pH 7 and 37°C, until the returning fluid was clear. After 30 minutes, exactly 25 ml. test solution was instilled into the lumen of the fistula. At 10 minutes as much as possible of the luminal solution was withdrawn and the segment immediately irrigated with 100 ml. non-radioactive Tyrode's solution. From the radioactivity acquired by this solution, the volume of test solution, which could not be aspirated at 10 minutes, was calculated.

The instillation of solutions into, and their withdrawal from, the isolated intestinal segments were facilitated by modified Foley catheters (Code, Bass, McClary, Newnum, and Orvis, 1960) which were inserted into each end of the fistula, leakage being prevented by inflation of the balloons of the catheters.

Two further 10-minute absorption tests were carried out in a similar manner at 1½ and 2½ hours after the end of the first test. The three absorption tests constituted a single experiment.

During all experiments an isotonic solution of sodium chloride was infused intravenously at a rate of 10 ml. per hour by a Palmer constant-rate infusion pump. In some experiments, the pentapeptide (I.C.I. 50,123) was added to the infusion solution to supply a dose of 4 µg. or 8 µg. per kilogram body weight per hour. In control experiments the saline solution was infused alone.

TEST SOLUTION The test solution was modified Tyrode's solution (Code and McIntire, 1956) containing the radioactive isotopes of sodium (²⁴Na—2 µc./litre solution) and of potassium (⁴²K—4 µc./litre solution), and the stable isotope of water, deuterium oxide (D₂O—1% v/v). The reaction of the solution, the temperature of which was maintained at 37°C, was brought to pH 7 with 0.1N hydrochloric acid.

ESTIMATIONS The concentrations of sodium and potassium in the test solution were estimated by flame photometry. The separate activities of ²⁴Na and ⁴²K in mixtures were determined by differential counting in a well-type scintillation counter and in a Geiger-Muller M6 tube, with a wall 1 mm. thick (Veall and Vetter, 1958). The concentration of D₂O was determined by infrared spectroscopy using a modification (Shields *et al.*, 1966) of the method of Berglund-Larsson (1956).

CALCULATIONS The rates of movement of sodium, potassium, and water into and out of the intestinal lumen were determined from the formulae of Visscher, Fetcher, Carr, Gregor, Bushey, and Barker (1944a) and Visscher, Varco, Carr, Dean, and Erickson (1944b).

The errors of the method and the assumptions inherent in the calculations have been discussed in detail (Shields *et al.*, 1966).

TERMINOLOGY Sodium, potassium, and water move simultaneously in both directions across the intestinal mucosa. 'Absorption' of a substance is applied to the situation where more of the substance leaves the intestinal lumen and enters the body than moves in the opposite direction into the intestinal lumen. 'Secretion' represents the converse situation, where the substance enters the intestinal lumen more rapidly than it leaves. The terminology describing the directions of movement of water and electrolytes across the intestinal mucosa has been reviewed (Shields, 1964).

RESULTS

NET TRANSPORT (TABLE I) In control experiments sodium and water were absorbed by the isolated

segments of jejunum, ileum, and colon. The rates of net transport of these substances were greater in the ileum and colon than in the jejunum. Potassium was secreted by the small intestine but was absorbed by the colon.

When the synthetic gastrin-like pentapeptide was infused at rates of 4 and 8 $\mu\text{g.}$ per kilogram per hour, the rates of absorption of sodium and water were significantly reduced in the ileum. The pentapeptide had no effect upon the net transport of these substances in the jejunum or in the colon.

In the jejunum the secretion of potassium was significantly increased when the pentapeptide was infused intravenously at rates of 4 and 8 $\mu\text{g.}$ per kilogram per hour. In the ileum, the rate of potassium secretion was increased during the infusion of 4 $\mu\text{g.}$ per kilogram per hour, but not with the larger dose of pentapeptide. The net transport rate of potassium in the colon was not affected by the intravenous infusion of the pentapeptide.

UNIDIRECTIONAL MOVEMENTS (TABLE I) When the pentapeptide was given at a rate of 4 $\mu\text{g.}$ per kilogram

TABLE I
THE MEAN RATES ($\pm$ S.E. MEAN) OF TRANSPORT OF SODIUM, POTASSIUM, AND WATER IN JEJUNUM, ILEUM, AND COLON IN CONTROL EXPERIMENTS AND DURING THE INTRAVENOUS INFUSION OF SYNTHETIC PENTAPEPTIDE AT RATE OF 4 $\mu\text{g.}$ and 8 $\mu\text{g./kg./hr.}$

Site	Experiment	Sodium Transport ($\mu\text{-equiv./10 min.}$)			Potassium Transport ($\mu\text{-equiv./10 min.}$)			Water Transport (ml./10 min.)		
		Net ¹	Out of Lumen	Into Lumen	Net ¹	Out of Lumen	Into Lumen	Net ¹	Out of Lumen	Into Lumen
Jejunum	1 Control (9) ^a	+2 $\pm$ 26	417 $\pm$ 31	415 $\pm$ 32	-3 $\pm$ 2	14 $\pm$ 1	17 $\pm$ 2	+0.2 $\pm$ 0.1	7.0 $\pm$ 0.4	6.8 $\pm$ 0.5
	2 Pentapeptide—4 $\mu\text{g./kg./hr.}$ (12)	-64 $\pm$ 44	564 $\pm$ 43	628 $\pm$ 58	-11 $\pm$ 3	17 $\pm$ 1	28 $\pm$ 3	-0.1 $\pm$ 0.2	7.5 $\pm$ 0.8	7.6 $\pm$ 0.9
	Difference between 1 and 2									
	Mean $\pm$ S.E.	66 $\pm$ 55	147 $\pm$ 56	213 $\pm$ 73	8 $\pm$ 4	3 $\pm$ 1	11 $\pm$ 4	0.3 $\pm$ 0.3	0.5 $\pm$ 1.0	0.8 $\pm$ 1.2
	t	1.2	2.6	2.9	2.6	3.0	2.7	1.0	0.5	0.7
	P	>0.2	<0.02	<0.01	<0.02	<0.01	<0.02	>0.3	>0.6	>0.5
Jejunum	3 Pentapeptide—8 $\mu\text{g./kg./hr.}$ (5)	+21 $\pm$ 52	557 $\pm$ 88	536 $\pm$ 112	-12 $\pm$ 2	17 $\pm$ 1	29 $\pm$ 2	+0.3 $\pm$ 0.2	6.6 $\pm$ 0.3	6.3 $\pm$ 0.4
	Difference between 1 and 3									
	Mean $\pm$ S.E.	19 $\pm$ 52	140 $\pm$ 76	121 $\pm$ 91	9 $\pm$ 3	3 $\pm$ 1	12 $\pm$ 3	0.1 $\pm$ 0.2	0.4 $\pm$ 0.6	0.5 $\pm$ 0.7
	t	0.4	1.8	1.3	3.0	3.0	4.0	0.5	0.7	0.8
	P	>0.7	>0.05	>0.2	<0.02	<0.02	<0.01	>0.6	>0.5	>0.3
Ileum	1 Control (20)	+190 $\pm$ 25	671 $\pm$ 36	481 $\pm$ 34	-3 $\pm$ 1	20 $\pm$ 1	23 $\pm$ 1	+1.2 $\pm$ 0.1	10.6 $\pm$ 0.6	9.4 $\pm$ 0.6
	2 Pentapeptide—4 $\mu\text{g./kg./hr.}$ (6)	+31 $\pm$ 48	619 $\pm$ 16	588 $\pm$ 53	-15 $\pm$ 4	19 $\pm$ 2	34 $\pm$ 6	+0.5 $\pm$ 0.3	9.0 $\pm$ 0.5	8.5 $\pm$ 0.6
	Difference between 1 and 2									
	Mean $\pm$ S.E.	159 $\pm$ 53	52 $\pm$ 67	107 $\pm$ 69	12 $\pm$ 3	1 $\pm$ 2	10 $\pm$ 4	0.7 $\pm$ 0.3	1.6 $\pm$ 1.2	0.9 $\pm$ 1.2
	t	3.0	0.8	1.5	3.4	0.5	2.5	2.3	1.3	0.7
	P	>0.01	>0.4	>0.1	<0.001	>0.6	<0.02	<0.05	>0.2	>0.4
Ileum	3 Pentapeptide—8 $\mu\text{g./kg./hr.}$ (3)	+11 $\pm$ 17	377 $\pm$ 135	366 $\pm$ 119	-2 $\pm$ 2	20 $\pm$ 6	22 $\pm$ 4	+0.3 $\pm$ 0.1	6.2 $\pm$ 0.2	5.6 $\pm$ 0.1
	Difference between 1 and 3									
	Mean $\pm$ S.E.	179 $\pm$ 67	294 $\pm$ 105	115 $\pm$ 98	1 $\pm$ 10	0 $\pm$ 1	1 $\pm$ 10	0.9 $\pm$ 0.3	4.4 $\pm$ 1.7	3.8 $\pm$ 1.6
	t	2.7	2.8	1.2	0.1	N.S.	0.1	2.5	2.6	2.3
	P	<0.02	<0.02	>0.2	>0.9	N.S.	>0.9	<0.02	<0.02	<0.05
Colon	1 Control (3)	+158 $\pm$ 115	389 $\pm$ 97	231 $\pm$ 211	+4 $\pm$ 1	12 $\pm$ 1	8 $\pm$ 2	+1.2 $\pm$ 0.3	5.4 $\pm$ 0.4	4.2 $\pm$ 0.7
	2 Pentapeptide—4 $\mu\text{g./kg./hr.}$ (3)	+120 $\pm$ 80	381 $\pm$ 180	261 $\pm$ 105	-2 $\pm$ 4	15 $\pm$ 3	17 $\pm$ 2	+0.7 $\pm$ 0.7	4.8 $\pm$ 0.6	4.2 $\pm$ 0.7
	Difference between 1 and 2									
	Mean $\pm$ S.E.	38 $\pm$ 134	8 $\pm$ 243	30 $\pm$ 207	6 $\pm$ 5	3 $\pm$ 4	9 $\pm$ 3	0.5 $\pm$ 0.8	0.6 $\pm$ 0.8	0 $\pm$ 0.7
	t	0.3	0.03	0.1	1.2	0.8	3.0	0.5	0.6	N.S.
	P	>0.7	>0.9	>0.8	>0.3	>0.4	<0.05	0.6	>0.5	N.S.

¹The plus and minus signs preceding the mean rates of net transport indicate absorption and secretion respectively.

^aThe figures in parenthesis indicate the number of experiments.

per hour, potassium entered the lumen of the jejunum, ileum, and colon more rapidly than in the control experiments. When the rate of infusion of pentapeptide was doubled, the increase in the rate of potassium movement into the lumen was maintained in the jejunum. Also, in the jejunum, at both rates of pentapeptide infusion, potassium moved more rapidly out of the lumen than in control experiments. In the ileum at the higher rate of infusion the rates of unidirectional movement of potassium did not differ significantly from those in control experiments.

The pentapeptide had no constant effect upon the unidirectional fluxes of sodium and water.

CONCENTRATION OF ELECTROLYTES IN THE LUMINAL SOLUTION (TABLE II) In control experiments the concentration of sodium in the jejunal and ileal solutions did not change significantly during 10 minutes. Also, in the jejunum, no change was observed in the concentration of potassium. There was a slight but significant increase in the ileal concentration of potassium during an absorption test.

During the infusion of the synthetic pentapeptide (at a rate of 4 $\mu\text{g./kg./hr.}$), the concentration of sodium in the ileum rose slightly. The concentration of potassium in both jejunum and ileum increased significantly when the pentapeptide was infused at

this dose rate. During the intravenous infusion of the pentapeptide the concentrations of potassium in the jejunal and ileal lumen after 10 minutes were significantly greater ($P < 0.02$ and $P < 0.05$, respectively) than the corresponding final concentrations of potassium in the control experiments.

An increase of similar magnitude was observed in the concentration of potassium in the jejunum when the pentapeptide was infused at 8 $\mu\text{g. per kg. per hour.}$

DISCUSSION

These results show that, when the synthetic gastrin-like pentapeptide was infused intravenously into dogs, the rates of movement of sodium, potassium, and water into and out of isolated segments of small intestine were markedly altered. The intestinal action of the pentapeptide seemed to be selective. Thus in the ileum, but not in the jejunum, the rates of absorption of sodium and water were greatly reduced. In both the ileum and jejunum, when pentapeptide was infused at 4 $\mu\text{g. per kilogram per hour}$, potassium was secreted more rapidly than in control studies. At the higher rate of infusion of pentapeptide (8 $\mu\text{g./kg./hr.}$), the increase in potassium secretion was maintained in the jejunum but, in the ileum, the rate of secretion returned to control levels. There is no obvious explanation for the failure of the larger dose of pentapeptide to stimulate the ileal secretion of potassium. The colonic handling of water and electrolytes was not affected by the infusion of the gastrin-like pentapeptide.

Recently Gardner, Peskin, Cerda, and Brooks (1967) studied the effect of several gastrointestinal hormones upon the transport of water and electrolytes by everted sacs of hamster intestine *in vitro*. The introduction of a crude porcine extract of gastrin into the solution bathing the mucosal surface of sacs formed from the distal third of small intestine was followed by a reduction in the net absorption of sodium, chloride, and water. Potassium transport was not affected nor did the remainder of the small intestine show any alteration in the handling of water and electrolytes. The unidirectional flux rates of sodium were studied: the mucosal-to-serosal movement of sodium was reduced when gastrin was instilled into the mucosal solution. The selective action of the gastrin extract upon the ileal absorption of sodium and water is similar to that found in the present study. However, close comparison of the results must be avoided in view of the differences in the nature of the gastrin, in the type of experimental preparation, and in the species of animal.

The changes in the intestinal transport of water

TABLE II

MEAN CONCENTRATION OF ELECTROLYTE (m-equiv/l $\pm$ S.E. of mean) IN LUMEN OF JEJUNUM AND ILEUM AT BEGINNING (t_0) AND END (t_{10}) OF 10-MINUTE ABSORPTION TEST IN CONTROL EXPERIMENTS AND DURING THE INTRAVENOUS INFUSION OF THE SYNTHETIC PENTAPEPTIDE AT 4 $\mu\text{g./kg./hr.}$

Segment	Solution	Control Experiments	Pentapeptide Infusion
Sodium			
Jejunum	Concentration at t_0	150 $\pm$ 0.4 (9) ¹	152 $\pm$ 0.8 (12)
	Concentration at t_{10}	151 $\pm$ 1.0 (9)	154 $\pm$ 1.0 (12)
	Mean difference $\pm$ S.E. mean	1.0 $\pm$ 1.0	2 $\pm$ 1.3
	t	0.87	1.5
	P	>0.3	>0.1
Ileum	Concentration at t_0	152 $\pm$ 0.7 (20)	149 $\pm$ 1.1 (6)
	Concentration at t_{10}	152 $\pm$ 0.6 (20)	151 $\pm$ 0.7 (6)
	Mean difference $\pm$ S.E. mean	0 $\pm$ 0.9	2 $\pm$ 0.8
	t	N.S.	2.3
	P	N.S.	<0.05
Potassium			
Jejunum	Concentration at t_0	3.84 $\pm$ 0.08 (9)	3.98 $\pm$ 0.14 (12)
	Concentration at t_{10}	3.99 $\pm$ 0.06 (9)	4.41 $\pm$ 0.12 (12)
	Mean difference $\pm$ S.E. mean	0.15 $\pm$ 0.10	0.43 $\pm$ 0.19
	t	1.5	2.3
	P	>0.1	<0.05
Ileum	Concentration at t_0	3.92 $\pm$ 0.10 (20)	4.00 $\pm$ 0.13 (6)
	Concentration at t_{10}	4.25 $\pm$ 0.10 (20)	4.90 $\pm$ 0.30 (6)
	Mean difference $\pm$ S.E. mean	0.33 $\pm$ 0.14	0.90 $\pm$ 0.35
	t	2.3	2.9
	P	<0.05	<0.05

¹Figures in parenthesis indicate number of experiments.

and electrolytes may have been brought about by the direct action of the pentapeptide upon the mucosal cells of the small intestine. On the other hand, the observed effects may have been both secondary and non-specific. First, the reduction in the rates of absorption of sodium and water and the increase in potassium secretion, following the infusion of the pentapeptide, may be secondary to changes in, for example, intestinal motility. However, not only is the influence of motility on intestinal absorption far from clear (Shields, 1964) but also the alterations in the motility of the small intestine produced by the pentapeptide are not marked (Logan and Connell, 1966; Neely, 1967; Bennett, Misiewicz, and Waller, 1967).

Second, the observed alterations in absorption and secretion may have been secondary to the stimulation of gastric secretion of acid. Indeed the doses of pentapeptide used in this study were those which Thomas and Forrest (1967), working in this laboratory, have found to produce the highest output of acid from denervated Heidenhain gastric pouches. However, there is no evidence to suggest that such hypersecretion of acid by the intact stomach and, particularly, the entry of considerable quantities of the acid into the small intestine would *per se* affect the absorptive and secretory behaviour of isolated segments of intestine (Miles, Davies, and Shields, 1965).

Finally the effect of the pentapeptide on the intestinal transport of water and electrolytes may be the non-specific action of a foreign peptide. For this reason the influence of pure antral gastrins isolated by Gregory and Tracy (1964) upon intestinal absorption should be studied.

These changes in the handling of water and electrolytes by the small intestine were observed over three hours, in isolated segments of bowel with a serosal surface area of 100 cm.². From the data given by Davenport (1961) it can be calculated that the mucosal area of these segments would probably represent one-seventh of the mucosal surface area of the small bowel. If the entire intestine responded to the pentapeptide in a manner similar to that of the isolated fistulas the losses of fluid and electrolytes would be considerable. However, certain additional points must be taken into consideration. First, the absorptive ability of the colon was quite unaffected by the pentapeptide. Because the colon has a reserve capacity to absorb five to seven times as much water and salt as it is normally called upon to do (Shields and Miles, 1965), increased absorption of sodium and water by the colon may very well compensate for the observed reduction in absorption by the ileum. Second, the artificial nature of these experiments must be borne in mind. Before any

definitive conclusions can be reached upon the effect of the natural antral hormone upon the intestinal handling of water and electrolytes, the intestine will have to be studied with its normal anatomical relationships preserved so that intestinal mucosa is in contact with food and alimentary secretions.

The relationship between the results of the present study and certain aspects of the Zollinger-Ellison syndrome remains to be discussed. The cause of the diarrhoea and hypokalaemia which are additional, occasionally dominating, features of this syndrome (Zollinger and Grant, 1964) remains obscure. The pancreatic adenoma is considered to produce a gastrin-like hormone which stimulates the hypersecretion of acid by the stomach (Gregory and Tracy, 1964). Some have attributed the accompanying diarrhoea to irritation of the small intestine by large volumes of acid (Donaldson, Vom Eigen, and Dwight, 1957; Summerskill, 1959; Rawson, England, Gillam, French, and Stammers, 1960; Parker, Soergel, and Ellison, 1963; Miles *et al.*, 1965). However, it may be that the hormone liberated by the pancreatic adenoma has a dual effect upon the intestinal handling of water and electrolytes. In addition to the secondary irritation of the small intestine with gastric acid, the gastrin-like hormone may, as suggested by the present study, act directly upon the small intestine to augment the secretion of potassium by the jejunum and ileum, and to reduce the absorption of sodium and water in the ileum. The critical experiment remains to study the effect upon intestinal absorption of a potent extract of pancreatic adenoma from a patient with the Zollinger-Ellison syndrome.

A less common form of the syndrome consists of a non-beta islet cell tumour of the pancreas associated with profuse watery diarrhoea and severe hypokalaemia with hypochlorhydria, or even achlorhydria, and without peptic ulceration (Verner and Morrison, 1958; Matsumoto, Peter, Schultze, Hakim, and Franck, 1966). In one patient, presenting such features, the increased secretion of potassium seemed to be caused by an increase in the movement of potassium ions into the intestinal lumen (Espiner and Beaven, 1962)—a response similar in character to the one observed in the present study. In this context, it is of interest to note that increased movement of potassium into the intestinal lumen has also been observed in other situations in which hormones seem to be implicated, for example, following the administration of aldosterone (Shields *et al.*, 1966), in primary hyperaldosteronism (Shields, 1966), and during salt depletion (Clarke, Miller, and Shields, 1967).

In two patients with diarrhoea and hypokalaemia

but without gastric hypersecretion and peptic ulceration, assay of the pancreatic adenoma did not reveal any gastrin-like activity (Matsumoto *et al.*, 1966). The mechanism of production of the diarrhoea remains obscure in these cases and the existence of another hormone, originally suggested by Telling and Smiddy (1961), seems most likely. Attempts by experiments *in vitro* to demonstrate that extracts from these tumours can affect intestinal absorption and secretion of fluid and electrolytes have not met with success (Matsumoto *et al.*, 1966). However, *in vitro* preparations are not entirely suitable for the study of intestinal transport of water and electrolytes (Shields, 1964).

SUMMARY

The movement of sodium, potassium, and water into and out of isolated segments of jejunum, ileum, and colon of dogs was studied *in vivo* during the intravenous infusion of a synthetic gastrin-like pentapeptide (I.C.I. 50,123) in doses of 4 μ g. and 8 μ g. per kilogram body weight per hour. There was a marked reduction in the rates of absorption of sodium and water in the ileum. In the jejunum and ileum the rates of secretion of potassium were increased when the pentapeptide was infused at a rate of 4 μ g. per kilogram per hour and, in the jejunum alone, with the higher dose of pentapeptide. The rates of transport of these substances in the colon were not affected by the pentapeptide.

There was a significant increase in the concentration of potassium in the lumen of the small intestine during the infusion of pentapeptide at 4 μ g. per kilogram per hour.

The action of the pentapeptide is considered to be a direct and specific one on the small intestinal mucosa.

The relationship between the results of this study and the variants of the Zollinger-Ellison syndrome is discussed.

We thank Professor A. P. M. Forrest for his advice on the dosage of the pentapeptide. We are grateful to Dr J. D. Fitzgerald of Imperial Chemical Industries Limited, for the supplies of I.C.I. (50,123). This work was supported by a grant from the Medical Research Council to Mr. R. Shields.

We are grateful to Professor C. A. Taylor of the Department of Physics, University College, Cardiff, for granting access to a Stantec Zebra computer on which the rates of net transport and of unidirectional fluxes were calculated.

REFERENCES

- Bennett, A., Misiewicz, J. J., and Waller, S. L. (1967). The effect of gastrin and a gastrin-like pentapeptide on human isolated gastro-intestinal muscle. (Abstract). *Brit. J. Surg.*, **54**, 232-233.
- Berglund-Larsson, U. (1956). Determination of small amounts of deuterium oxide by infra-red spectroscopy. *Acta chem. scand.*, **10**, 701-702.
- Clarke, A. M., Miller, M., and Shields, R. (1967). Intestinal transport of sodium, potassium, and water in the dog during sodium depletion. *Gastroenterology*, **52**, 846-858.
- Code, C. F., Bass, P., McClary, G. B., Jr., Newnum, R. L., and Orvis, A. L. (1960). Absorption of water, sodium and potassium in small intestine of dogs. *Amer. J. Physiol.*, **199**, 281-288.
- , and McIntire, F. C. (1956). Quantitative determination of histamine. *Meth. biochem. Anal.*, **3**, 49-95.
- Davenport, H. W. (1961). *Physiology of the Digestive Tract*, p. 161. Year Book Medical Publishers, Chicago.
- Donaldson, R. M., Jr., Vom Eigen, P. R., and Dwight, R. W. (1957). Gastric hypersecretion, peptic ulceration and islet-cell tumor of the pancreas (the Zollinger-Ellison syndrome): report of a case and review of the literature. *New Engl. J. Med.*, **257**, 965-970.
- Espiner, E. A., and Beaven, D. W. (1962). Non-specific islet-cell tumour of the pancreas with diarrhoea. *Quart. J. Med.*, **31**, 447-471.
- Gardner, J. D., Peskin, G. W., Cerda, J. J., and Brooks, F. P. (1967). Alterations of *in vitro* fluid and electrolyte absorption by gastrointestinal hormones. *Amer. J. Surg.*, **113**, 57-64.
- Gingell, J. C., Davies, M. W., and Shields, R. (1966). The effect of gastrin on the absorption and secretion of water and electrolytes in the small intestine. *Brit. J. Surg.*, **53**, 990.
- Gregory, R. A., and Tracy, H. J. (1964). The constitution and properties of two gastrins extracted from hog antral mucosa. *Gut*, **5**, 103-117.
- Konturek, S., and Grossman, M. I. (1966). Acid response to gastrin and related peptides. *Gastroenterology*, **50**, 650-652.
- Logan, C. J. H., and Connell, A. M. (1966). The effect of a synthetic gastrin-like pentapeptide (I.C.I. 50,123) on intestinal motility in man. *Lancet*, **1**, 996-999.
- Matsumoto, K. K., Peter, J. B., Schultze, R. G., Hakim, A. A., and Franck, P. T. (1966). Watery diarrhoea and hypokalemia associated with pancreatic islet cell adenoma. *Gastroenterology*, **50**, 231-242.
- Miles, J. B., Davies, M. W., and Shields, R. (1965). A study of the possible causes of the watery diarrhoea of the Zollinger-Ellison syndrome. *Brit. J. Surg.*, **52**, 625.
- Neely, J. (1967). A comparison of the effects of a gastrin extract and synthetic pentapeptide (I.C.I. 50,123) on gastro-intestinal motility in the cat. (Abstract). *Ibid.*, **54**, 232.
- Parker, P. E., Soergel, K., and Ellison, E. H. (1963). Effects of excessive hydrochloric acid on the canine gastrointestinal tract. *Surg. Forum*, **14**, 333-334.
- Rawson, A. B., England, M. T., Gillam, G. G., French, J. M., and Stammers, F. A. R. (1960). Zollinger-Ellison syndrome with diarrhoea and malabsorption. Observations on a patient before and after pancreatic islet-cell tumour removal without resort to gastric surgery. *Lancet*, **2**, 131-134.
- Shields, R. (1964). Surgical aspects of the absorption of water and electrolytes by the intestine. *Monogr. Surg. Sci.*, **1**, 119-172.
- (1966). The role of the adrenal cortex in the intestinal absorption of water and electrolytes. In *Postgraduate Gastroenterology*, edited by T. J. Thomson and I. E. Gillespie, pp. 30-43. Ballière, Tindall and Cassell, London.
- , and Miles, J. B. (1965). Absorption and secretion in the large intestine. *Postgrad. med. J.*, **41**, 435-439.
- , Mulholland, A. T., and Elmslie, R. G. (1966). Action of aldosterone upon the intestinal transport of potassium, sodium, and water. *Gut*, **7**, 686-696.
- Summerskill, W. H. J. (1959). Malabsorption and jejunal ulceration due to gastric hypersecretion with pancreatic islet-cell hyperplasia. *Lancet*, **1**, 120-123.
- Telling, M., and Smiddy, F. G. (1961). Islet tumours of the pancreas with intractable diarrhoea. *Gut*, **2**, 12-17.
- Thomas, W. R. G., and Forrest, A. P. M. (1967). Acid secretion from canine separated fundic pouches. *Gastroenterology*, **52**, 192-197.
- Tracy, H. J., and Gregory, R. A. (1964). Physiological properties of a series of synthetic peptides structurally related to Gastrin I. *Nature (Lond.)*, **204**, 935-938.
- Veall, N., and Vetter, H. (1958). *Radioisotope Techniques in Clinical Research and Diagnosis*. Butterworth, London.

- Verner, J. V., and Morrison, A. B. (1958). Islet cell tumor and a syndrome of refractory watery diarrhea and hypokalemia. *Amer. J. Med.*, 25, 374-380.
- Vischer, M. B., Fetcher, E. S., Jr., Carr, C. W., Gregor, H. P., Bushey, M. S., and Barker, D. E. (1944a). Isotopic tracer studies on the movement of water and ions between intestinal lumen and blood. *Amer. J. Physiol.*, 142, 550-575.
- , Varco, R. H., Carr, C. W., Dean, R. B., and Erickson, D. (1944b). Sodium ion movement between the intestinal lumen and the blood. *Ibid.*, 141, 488-505.
- Wormsley, K. G., Mahoney, M. P., and Ng, M. (1966). Effects of a gastrin-like pentapeptide (I.C.I. 50,123) on stomach and pancreas. *Lancet*, 1, 993-996.
- Zollinger, R. M., and Grant, G. N. (1964). Ulcerogenic tumor of the pancreas. *J. Amer. med. Ass.*, 109, 181-184.

The December 1967 Issue

THE DECEMBER 1967 ISSUE CONTAINS THE FOLLOWING PAPERS

- Recording of intestinal motility: routine or research? A. M. CONNELL
- Population control in the small bowel SHERWOOD L. GORBACH
- Observations on the epidemiology of appendicitis DAVID J. B. ASHLEY
- Evidence of vitamin E deficiency in patients with malabsorption or alcoholism and the effects of therapy M. S. LOSOWSKY and P. J. LEONARD
- Morphological observations on gastric ulcers treated with carbenoxolone sodium T. E. W. GOODIER, L. HORWICH, and R. W. GALLOWAY
- A study of the effect of saliva on the concentration of mucin in gastric juice and its possible relationship to the aetiology of peptic ulcer S. L. MALHOTRA
- Effect of the exocrine pancreatic secretions on iron absorption H. KAVIN, R. W. CHARLTON, P. JACOBS, R. GREEN, J. D. TORRANCE, and T. H. BOTHWELL
- Cystic arterial patterns in diseased human gall bladders K. C. D. GORDON
- An experimental method for recording the behaviour of human isolated colonic segments ANNE BUCKNELL and C. CLARK
- Influence of gastric pH on gastric and jejunal flora J. D. ALLAN GRAY and M. SHINER
- Gastric, haematological, and immunological abnormalities in Hashimoto's thyroiditis K. F. R. SCHILLER, L. MICHAEL SNYDER, and M. B. VALLOTTON
- Acid and pepsin responses to gastrin in Heidenhain pouch dogs following bilateral adrenal ectomy ALLAN R. COOKE
- Effect of intravenous calcium administration on gastric secretion of acid and pepsin in man R. A. SMALLWOOD
- Validity of polyethyleneglycol in the study of the total pancreatic secretion in man after stimulation by secretin, pancreozymin, or food H. WORNING
- Neuromuscular disease in patients with steatorrhea H. J. BINDER, G. B. SOLITARE, and H. M. SPIRO
- Reversal of non-Addisonian achlorhydria by a Roux-en-Y loop W. M. CAPPER, T. J. BUTLER, and J. O. KILBY
- Risk of anicteric hepatitis following blood transfusion B. N. SOMAYAJI, W. D. STONE, and P. B. GLOVER
- Surface area of the small intestine in man J. P. WILSON
- Absorption of lactose and its digestion products in the normal and malnourished Ugandan G. C. COOK, ANNE LAKIN, and R. G. WHITEHEAD

The British Society of Gastroenterology

Copies are still available and may be obtained from the PUBLISHING MANAGER,
BRITISH MEDICAL ASSOCIATION, TAVISTOCK SQUARE, W.C.1, price 18s. 6d.