"90% satisfactory response obtained in ambulant patients"

"The immediate clinical results were assessed after the first month's treatment (with Roter Tablets) . . . 81% of cases became symptom-free—70% of them during the first week and 30% during the second week; a further 9% were relieved of the majority of their symptoms. Thus there was a satisfactory response in 90% of cases In 75% of cases the patients found they were able to take foods which they had avoided for years "

Extract from the BRITISH MEDICAL JOURNAL (1955) 2, 827

IMPROVED THERAPY FOR **PEPTIC ULCER**

"In a series of 155 cases of peptic ulcer treated with . . . (Roter) tablets for three to six weeks, there was immediate relief of pain in 92% of cases and roent-genological healing of the ulcer in 81% . . . The special features of this medication are immediate and prolonged relief of pain, correction of gastric hyperacidity, absence of side reactions, effectiveness for ambulatory patients, and early healing of the ulcer in most cases."

Extract from the AMERICAN JOURNAL OF GASTROENTEROLOGY (1957) 28, 439

"Immediate relief of pain in 92% of cases"



FORMULA PER TABLET. Magnesium Carbonate (400 mg.), Bismuth Subnitrate Roter (350 mg.), Sodium Bicarbonate (200 mg.), and Cortex Rhamni Frangulae (25 mg.).

PACKINGS. Tins of 40, 120, and dispensing sizes 360 and 720 (P.T. exempt.). Basic N.H.S. price: 1/- for 10 tablets.

ROTER Tablets are not advertised to the public and may be prescribed on E.C.10 forms. Literature on request.

F.A.I.R. LABORATORIES LTD., TWICKENHAM, MIDDLESEX

ALUHYDE

for prompt, safe symptomatic medication in hyperchlorhydria and peptic ulcer

'ALUHYDE' tablets present a new combination with these advantages...

'ALUHYDE' effects safe, reliable, nonsystemic neutralization by means of two of the most preferred modern antacid adsorbents...



'ALUHYDE' permits of quick mental sedative-effect by its inclusion of a barbiturate recognized for minimal cumulative potential...



'ALUHYDE' provides antispasmodic action through its content of an agent long known for its favourable influence against hypermotility...



Formula-

Each tablet: Dried Aluminium

Hydroxide Gel. B.P.C. - 3.75 gr.

Magnesium trisilicate B.P. - 3.75 gr.

Quinalbarbitone sodium B.P. - 0.50 gr.

Tinct. Belladonnæ B.P. - 3 m.

Packings—Regular packs of 50; bottles of 500 (dispensing)


Samples available on personally signed request of qualified physicians only


MULTIPAX CHEMICALS LTD


142, 144, 146 LARKHALL LANE, LONDON, S.W.4

Phones: MACaulay 8493/6821









SALAZOPYRIN

BRAND OF SALICYLAZOSULPHAPYRIDINE

to be

a drug

of significance

in to-day's management

of this distressing

disease.







Outstanding characteristics of Salazopyrin treatment are

- Prompt remission of symptoms
- Bleeding and frequency of stools sharply reduced
- Abdominal cramp controlled
- In many cases rectal mucosa heals within a month
- Fever subsides, appetite improves
- Can be given safely over long periods of time


SALAZOPYRIN is also a safe and effective product in other forms of colitis and regional enteritis.



PHARMACIA UPPSALA: SWEDEN

Distributed by

Savory & Moore Ltd. 60/61 Welbeck Street, London W. 1



long-acting anticholinergic

Tyrimide*

Experimental studies and clinical use have established that:

- 1. 'Tyrimide' is <u>inherently long-acting</u>. One dose in the morning and another at night provide an uninterrupted 24-hour effect
- 2. 'Tyrimide' is <u>effective</u>. Tests of anti-secretory and anti-spasmodic activity have proved 'Tyrimide' to be more effective than other newer synthetics and atropine
- 3. 'Tyrimide' is remarkably free from troublesome side-effects

'Tyrimide'—for the anticholinergic management of peptic ulcer and gastro-intestinal disorders

EACH 'TYRIMIDE' TABLET CONTAINS 5 MG. ISOPROPAMIDE IODIDE [3-carbamoyl-3, 3-diphenylpropyl] methyl diisopropyl ammonium iodide

Smith Kline & French Laboratories Ltd Welwyn Garden City, Herts


*'Tyrimide' (Trade mark) brand of isopropamide iodide. Brit. Pat. No. 77292I TM:PA30

NEW

a more convenient and economical presentation of hydrocortisone for ulcerative colitis

EFCORTELAN SOLUTION


TRADE MARK



Given as a nightly rectal drip, soluble hydrocortisone (Efcortelan) has been demonstrated to afford rapid alleviation of symptoms of ulcerative colitis in most patients. Symptomatic relief is accompanied by an improvement in the sigmoidoscopic findings and in the colonic mucosa.

The method of treatment is quite suitable for the patient to carry out himself provided the attack is not severe.

With soluble Efcortelan, hormonal side-effects are avoided. If further attacks occur they can usually be



reduced by another course of treatment.

Only one of the new Efcortelan

Solution Tablets in the appropriate
amount of normal saline (100-120 ml)
is needed to prepare the retention
enema. It is therefore the ideal
presentation for home treatment.


EFCORTELAN SOLUTION TABLETS

Foil strip of 7 tablets (one week's treatment). Each tablet contains 100 mg hydrocortisone (as water soluble sodium succinate).

Subsidiary Companies or Agents in most countries

GLAXO LABORATORIES LTD, GREENFORD, MIDDLESEX BYRON 3434





... 'Streptotriad' is indicated for the rapid control of the acute attack and is likely to prevent the patient from becoming a carrier, either during the convalescent or chronic stage of the disease. 'Streptotriad' is often effective also in the treatment of the symptomless carrier.

Presentations: Tablets, and as granules sufficient to prepare $1\frac{1}{2}$ fl. oz. of suspension. Each tablet or 3.6 ml. (approx. one teaspoonful) of the prepared suspension contains:

STREPTOMYCIN SUL	PHATE	65	mg
SULPHATHIAZOLE		100	mg
SULPHADIAZINE		100	mg
SULPHAMERAZINE		65	mg.

Detailed information is available on request



MANUFACTURED BY
MAY & BAKER LTD

An M&B brand Medical Product

MA7310/133

an improved





antacid preparation





'ALMACARB' tablets are prepared from an aluminium hydroxide-magnesium carbonate co-dried gel manufactured by an entirely new process (Brit. Pat. 812503). The result is a product of exceptional stability with a more rapid and prolonged acid-neutralising action than that of dried aluminium hydroxide gel, either alone or in mechanical admixture with magnesium carbonate.

'ALMACARB' tablets are unique because there is no loss of neutralising action either on processing or tabletting.
'ALMACARB' tablets sucked at regular intervals will ensure effective control of gastric acidity. Used in this way they are more convenient than liquid antacids.

Dosage

One tablet to be chewed as an initial dose and thereafter one tablet to be sucked slowly at regular intervals.

> Basic N.H.S. Prices Individually foil-wrapped tablets. Cartons of 40...3/-

Cartons of 200 . . . 12/-(Subject to P.T.)





THE BRITISH DRUG HOUSES LTD. (Medical Department) LONDON N.1

Pro-Banthine

SEARLE

GUT JUNE, 1960

NULACIN "has revolutionized the treatment of duodenal ulcer"

"This new method of the administration of antacids (Nulacin) has revolutionized the treatment of duodenal ulcer . . . milk drip is no longer part of our therapeutic stock in trade." PROC. ROY. SOC. MED., December, 1958, 51, 1063

N ULACIN tablets, when allowed to dissolve slowly in the mouth, provide intragastric milk-alkali drip therapy safely and effectively for both the bedridden and ambulant patient. They may be relied on to provide a treatment for peptic ulcer as well as other conditions associated with gastric hyperacidity.











The safety and effectiveness of Nulacin therapy has been proven by "in vivo" clinical work carried out in many countries. References to this work include:-

Current Status of the Medical Treatment of Peptic Ulcer, Med. Times, January, 1958, 1, 74

Antacids, The Practitioner, January, 1957, 178, 43

Antacids in Peptic Ulcer, The Practitioner, January, 1956, 176, 103

Recent Advances in the Ulcerative Diseases of the Gastrointestinal Tract, Amer. 7. Gastro., December, 1956, 26, 665

Ambulatory Continuous Drip Method in the Treatment of Peptic Ulcer, Amer. J. Dig. Dis., March, 1955, 22, 67-71


Management of Peptic Ulceration in General Practice, Med. World., December, 1954, 81, 591-601

Clinical Investigation into the Action of Antacids, The Practitioner, July, 1954, 173, 46

> Further Studies on the Reduction of Gastric Acidity, Brit. Med. J., 23rd January, 1954, I, 183-184

Control of Gastric Acidity by a New Way of Antacid Administration, J. Lab. & Clin. Med., 1953, 42, 955

The Effect on Gastric Acidity of "Nulacin" Tablets. Med. 7. Aust., 28th November, 1953, 2, 823-824




Nulacin tablets have no B.P. equivalent. The Basic N.H.S. price of the 25 tablet tube is 2/-.



Further information is available from:

HORLICKS LIMITED · PHARMACEUTICAL DIVISION · SLOUGH · BUCKS

freedom from pain



In peptic ulcer, freedom from acid means freedom from pain. With single, intermittent doses of antacids, peaks of acidity and bouts of pain are inevitable.

But when Prodexin tablets are sucked, continuous flow and efficient buffering give not merely relief, but constant protection from acid pain and a safe level of acidity to encourage the ulcer to heal.

Prodexin

the continuous-flow buffer antacid

TABLET CONTAINS: Light magnesium carbonate 0'l gramme


EACH PRODEXIN | Aluminium glycinate 0.9 gramme

PACKAGES: 30 and 240 tablets Basic N.H.S. cost 3/1d. and 23/4d.



BRENTFORD, MIDDLESEX Tel: Isleworth 4111

GUT



Over a number of years I.C.I. have established a comprehensive library of 16 mm. films covering a wide range of medical subjects. Made in collaboration with leading authorities, these films are available, free of charge, to interested borrowers. Two subjects of special interest in the field of gastroenterology are "GASTRIC SECRETION", which demonstrates the factors responsible for the secretion of the gastric juice, and "PANCREATIC SECRETION" which describes the

chief mechanisms concerned in the stimulation of pancreatic secretion by a meal.

Both films were made in collaboration with the Department of Physiology, University of Liverpool, and are in colour.

For further details and particulars of how to borrow these, and other I.C.I. medical films, please write to the

I.C.I. Film Library, Imperial Chemical House, Millbank, London, S.W.I.

ASPIRIN IS SAFER WHEN IT IS SOLUBLE

Soluble aspirin is safer when it is stable



N.H.S. Basic Price 12/6 for 500 tablets in foils.

Solderin provides soluble aspirin in its most stable form and dissolves rapidly in water to form a solution of calcium acetylsalicylate.

SOLPRIN is easy to take and palatable. It has proved rapid and reliable in its analgesic effect and is extremely well tolerated, even in prolonged and massive dosage.

Solprin is not advertised to the public and is available only on prescription. Clinical samples and literature will be supplied on request.

SOLPRIN

Soluble and substantially neutral


RECKITT & SONS LIMITED, PHARMACEUTICAL DEPARTMENT, HULL.

IN ULCER THERAPY



Bislumina was specifically developed to embody in a single preparation all those properties considered therapeutically desirable in an agent for the local treatment of peptic ulcer and allied dyspeptic disorders.

Bislumina combines inactivation of pepsin with a sustained antacid action and protection of the inflamed gastric mucosa.



BISLUMINA Tablets

each containing Bismuth Aluminate 795 mg.

Magnesium Oxide 100 mg.

Dosage: 1-2 tablets, three times daily before meals. Packs of 12, 30 and 250. Basic N.H.S. Price: 2/11

for 30 tablets.

Also available: BISLUMINA Cream 12 fl. oz. bottles.

Dose, 1-2 dessertspoonfuls.

Samples and Literature gladly sent on request.



M.C.P. Pure Drugs Ltd.
STATION WHARF, ALPERTON, MIDDLESEX



. . . unless an adequate supply of pancreatic enzymes is available to complete digestion.

Pancrex V is the most effective preparation for replacing pancreatic secretion in the gut. It is indicated in cystic fibrosis of the pancreas, chronic pancreatitis, pancreatectomy, post-cholecystectomy and in all digestive disorders where a relative deficiency of pancreatic enzymes is a contributory factor.

Pancrex V is activated, whole dried pancreas having a tryptic activity approximately equal to 5 times that of Pancreatin B. P.



Available in the form of powder or enteric-coated tablets.

GUT JUNE, 1960

"...ALL HAD VERY SEVERE ANAEMIA AND RESPONDED

EXTREMELY WELL" *

The most difficult patient and least satisfactory haemoglobin level RESPOND to treatment with

erroids

ORAL IRON THERAPY WHICH IS SAFE AND EFFECTIVE

'Ferroids' tablets contain 350 mg. iron aminoates, a partly chelated complex of iron and amino acids which

- . . . is well absorbed
- . . . is virtually free from side-effects
- ... provides unusually high utilisation of absorbed iron.

DOSAGE One tablet three times daily is usually sufficient. Up to eight tablets daily-and more-are well-tolerated even if taken between meals, the time for maximum absorption of ingested iron.

PACKS In containers of 100 and 1000 tablets.

"...ALL THE PATIENTS RESPONDED SATISFACTORILY TO CHELATED IRON AT A RATE COMPARABLE THOSE TREATED WITH INTRAMUSCULAR IRON"*



'Ferroids' is a registered trade mark

* I. Obstet. Gynaeco, Brit. Emp., (1959) 66:627.

RIKER LABORATORIES LIMITED LOUGHBOROUGH, ENGLAND

EMERGENCIES IN GENERAL PRACTICE

This book comprises 57 specially commissioned articles on medical emergencies which appeared originally in the *British Medical Journal*.

Contents include haemorrhage and thrombosis, cardio-respiratory crises, anaesthesia and hazards of therapy, acute abdominal cases, obstetrics, fits and coma, electrocution, burns and gassing, poisoning, acute psychiatric states, and special and minor surgery.

Invaluable to all doctors everywhere.

470 pages, cloth bound with full index. 25s. net (by post—inland and overseas 26s. 9d.).

CLINICAL PATHOLOGY IN GENERÁL PRACTICE

This handbook on clinical pathology meets the needs of the general practitioner, the houseman, and the senior student. It contains thirty-nine articles comprising a series specially written for the *British Medical Journal*. Each article has been revised and brought up-to-date by its author. The book gives authoritative information on available laboratory facilities; reliable tests and which to usc; techniques for collecting and preserving specimens; interpretation of results and significance of abnormal findings.

321 pages. Price 21s. (by post—inland and overseas 22s. 3d.).

REFRESHER COURSE FOR GENERAL PRACTITIONERS

Third Collection—with Classified Contents List to volumes 1, 2 and 3

Each book contains a collection of sixty Refresher Course articles from the *British Medical Journal*. Each article, written by an acknowledged expert, has been revised by its author. The articles cover a wide range. Their object is to refresh the reader's memory of what is old and well-established and to tell him about new developments.

They are invaluable to general practitioners and residents as practical reference books, to specialists as a means of keeping in touch with developments outside their own field, and to final-year students as up-to-date and reliable textbooks.

25s. (by post—inland and overseas 26s. 9d.).

"ANY QUESTIONS?"

Third Series—with Cumulative Index

These pocket-size volumes each contain some 200 questions and expert answers from the "Any Questions?" pages of the *British Medical Journal*. Each answer has been chosen for its practical value to doctors in their day-to-day work. Many deal with subjects not covered in the standard textbooks.

The Third volume has a cumulative index to all the answers appearing in the three books in the series.

7s, 6d. (by post 8s. 3d.).

Obtainable from booksellers or by post from

BRITISH MEDICAL ASSOCIATION

B.M.A. HOUSE, TAVISTOCK SQUARE, LONDON, W.C.1

PEPTIC ULCERATION

A Symposium for Surgeons

BY CHARLES WELLS, M.B., F.R.C.S., and JAMES KYLE, M.Ch., F.R.C.S.(Eng.), F.R.C.S.I.

272 pages

69 illustrations

"This symposium presents in considerable detail recently acquired knowledge and changes in practice within the more lightly sketched framework of what is generally agreed concerning gastric and duodenal ulceration. No attempt has been made to describe in detail techniques which can only be mastered in the operating theatre but space is devoted to methods of dealing with unexpected difficulties which may arise in the course of everyday procedures. Little is said about routine diagnosis and such like but exceptions from the general pattern are discussed at some length. By contrast, actiological factors and the post-gastrectomy syndromes are described very fully."—Extract from Preface.

PEPTIC ULCER

By C. F. W. ILLINGWORTH, C.B.E., M.D., Ch.M., F.R.C.S.E., Hon. F.R.C.S.(Eng.). 95 illustrations

296 pages

A CONTRIBUTION TO THE STUDY OF PORTAL HYPERTENSION By ALAN H. HUNT, M.A., D.M., M.Ch., F.R.C.S.

244 pages

205 illustrations

SURGERY OF THE ANUS, ANAL CANAL AND RECTUM

By E. S. R. HUGHES, M.D.(Melb.), (Melb.), F.R.C.S.(Eng.), F.R.A.C.S. 316 pages 398 illustrations M.S. 50s.

SURGERY OF THE COLON


By E. S. R. HUGHES

339 illustrations 424 pages

63s.

E. & S. LIVINGSTONE LTD. TEVIOT PLACE -**EDINBURGH**

ABSTRACTS OF WORLD MEDICINE



A monthly journal of informative abstractscovering every important article published in over 1,600 medical periodicals throughout the worlddealing in each issue with a comprehensive range of subjects.

FOR THE SPECIALIST to keep him informed of work in progress in other fields as well as his own.

FOR THE GENERAL PRACTITIONER

to enable him to keep abreast of new developments in every branch of medicine.

FOR THE POSTGRADUATE STUDENT

to give him a synoptic view of current progress and a guide to further reading.

FOR THE MEDICAL LIBRARY

to provide an

invaluable supplement to its existing resources.

A specimen copy may be obtained free from the Publishing Manager

MEDICAL ASSOCIATION BRITISH

ANNUAL SUBSCRIPTION (12 issues) £4 4s. U.S.A. AND CANADA \$13.50

Lignocaine?

SPECIFY



LIGNOCAINE Local anaesthetics

RAPID · DEEP · CERTAIN · LASTING · SAFE



XYLOTOX Solutions

For major and minor surgery. ½%, 1% and 2% solutions, plain or with adrenaline, in 20, 50, 250, and 500ml. bottles.

For surface anaesthesia of the mucosa. 4% solution, plain or with adrenaline, in 25ml. bottles



XYLOTOX Jelly

For effective urethral and vaginal anaesthesia.

For surgery use:— 15G. and 30G. tubes with applicators.

For instrument lubrication:—
Boxes 12 x 1.5G. Tubes.



XYLOTOX Pastes

For truly efficient surface anaesthesia of the mucosa.

XYLOTOX-NORMAL (without

amethocaine).

XYLOTOX-EXTRA (with amethocaine). $\frac{1}{2}$ -oz. tubes.



XYLOTOX Oral

For quick, simple surface anaesthesia of the upper digestive tract.

A viscous, pleasantly flavoured product. 200ml. bottles.



A division of

Full details upon request from:—

PHARMACEUTICAL MANUFACTURING COMPANY

The Local Anaesthetic Specialists

EPSOM . SURREY

WILLOWS FRANCIS LIMITED . EPSOM AND LONDON

