Progress report

Alkaline phosphatase

This review is selective and reflects the author's personal interest in the application of laboratory techniques to clinical problems; other aspects of the subject have been well covered in recent reviews.^{1,2,3,4,5,6}

Structure

The alkaline phosphatases are zinc metallo enzymes^{7,8} and it is known that zinc has a functional role in the catalysis;⁹ magnesium or cobalt are also required for enzyme activity.¹⁰

Purified placental¹¹ and intestinal¹² alkaline phosphatases contain approximately 15% nitrogen, and the amino-acid compositions of both enzymes are known.^{13,14} Hydrolysis of bacterial alkaline phosphatases has yielded amino-acid sequences around the active centre suggesting a structural as well as a functional analogy with other esterases; the sequence found near the reactive serine residue in the alkaline phosphatase of *E. coli* is similar to the sequences found at the active centres of trypsin and chymotrypsin.^{15,16} The covalent binding of inorganic phosphate to the serine in the active site is a reaction common to the alkaline phosphatases and an intermediate phosphoryl-enzyme is formed during their action.¹⁷

Like a number of enzymes, including caeruloplasmin¹⁸ and pancreatic ribonuclease B,¹⁹ alkaline phosphatase (AP) is a glycoprotein^{20,21} but the carbohydrate content of alkaline phosphatase has been the subject of conflicting reports. Ahmed and King found little or no carbohydrate in human placental AP,¹¹ whilst Ghosh reported appreciable amounts of carbohydrate in the purified enzyme.²² Purified intestinal AP contains substantial amounts of hexose and hexosamine²³ but no sialic acid.²⁴ Alkaline phosphatases from other tissues have, however, been shown to contain bound sialic acid; for instance, the structurally different but functionally similar isoenzymes of human kidney arise as a result of variations in the amount of bound sialic acid.²¹ The neuraminic-acid-containing nature of human placental AP has been demonstrated by chemical means and the sialic acid residues shown to occupy terminal positions away from the active centre of the enzyme.²⁵ Removal of sialic acid residues from AP, as from other glycoprotein enzymes, affects the electrophoretic mobilities of the protein but does not impair enzymatic activity. The role of the carbohydrate units in glycoprotein enzymes is at present not understood but they are known to exist with a heterogeneous complexity which may represent specific genetic characteristics;²⁶ there is no evidence that carbohydrate units participate in catalysis.

Many studies have reported differences in the chromatographic or electrophoretic properties of alkaline phosphatases in serum or in crude tissue homogenates, rather than in purified preparations; in such studies, specific as well as non-specific binding can occur which may account for the degree of

heterogeneity observed. However, studies on highly purified preparations of human placental AP¹³ have shown two major molecular weight variants on Sephadex G-200: the electrophoretically slower was obtained in crystalline form and had a molecular weight of over 200 000, whilst the faster moving variant had a molecular weight of 70 000. The high molecular weight variant could be converted into the lower molecular weight variant by storage for about four months, whilst the process could be reversed by equilibrium dialysis; kinetic studies indicated no difference between the two variants. *E. coli* and human placental alkaline phosphatases are dimers, each monomeric subunit of placental AP having a molecular weight of 58 000.²⁷ Between pH 4.7 and pH 10.3 human placental alkaline phosphatase appears as a homogeneous entity with the molecular weight of the native enzyme; however, above pH 10.3, a monomer-dimer equilibrium appears to be present, and at pH 2.3 the enzyme was also heterogeneous, with the formation of both low molecular weight compounds and high molecular weight aggregates.²⁸

Functions

Robison, the first to suggest an association between AP and calcification in cartilage and bone,^{29,30,31} originally explained calcification as a hydrolysis of a phosphoric acid ester, the calcium salt of which is easily soluble in water, releasing phosphate ions. He assumed that blood is normally in equilibrium with the inorganic constituents of bone and that the supersaturation necessary for the precipitation of bone salt is produced locally in cartilage and osteoid tissue by the alkaline phosphatase at these sites from the phosphoric esters present in blood. This simple hypothesis failed to explain why some tissues calcified and others, which contained large amounts of the enzyme, did not. Robison later conceived of two factors operating in calcification: first the phosphatase mechanism causing a supersaturation of phosphate ions in the matrix fluid of calcifying cartilage and, secondly, a 'local' factor favouring the deposition of calcium phosphate from supersaturated solutions, whether this state of supersaturation in the cartilage matrix be caused by enzymatic hydrolysis or by a supersaturated solution diffusing into the cartilage from outside.^{32,33} The role of alkaline phosphatase in calcification was considered to be of little significance by some later workers³⁴ but was re-emphasized by others;³⁵ more recent work has suggested that alkaline phosphatase plays a key role not only in calcification but also in bone resorption, by removing the layer of pyrophosphate which covers the mineral surface of bone and which has been shown to retard the rate of deposition of calcium and phosphate onto this surface.³⁶

These contributions throw no light on the possible functions of alkaline phosphatase at other sites in the body. The histochemical localization of the enzyme to the absorptive surfaces of the proximal convoluted tubule of the kidney,^{37,38} the small intestinal mucosa,³⁹ the syncytiotrophoblast of the placenta,⁴⁰ and the cell wall of *Escherichia coli*⁴¹ has suggested a connexion between these enzymes and transport, possibly of phosphate across cell walls, and there is some, albeit slender, evidence to support this contention,^{42,43,44,45} including the demonstration of the ability of alkaline phosphatase to bind inorganic phosphate.^{17,46}

Alkaline phosphatase may be connected with protein synthesis in the cell.^{47,48} The enzyme is thought to be concerned with nucleoprotein and nucleotide

metabolism.^{49,50,51} A functional relationship between alkaline phosphatase and RNA has been demonstrated,⁵² and AP may play a role in controlling DNA synthesis and growth;⁵³ the enzyme is capable of hydrolyzing both DNA⁵⁴ and RNA.⁵⁵ Alkaline phosphatases from many tissues possess both pyrophosphatase^{56,57} and phosphotransferase^{58,59} activities, and, in the case of intestinal, kidney, and bone phosphatases the two catalytic activities are the result of the action of the same enzyme. A wide range of compounds have been shown to act simultaneously as substrates of the hydrolase and the phosphotransferase activities of alkaline phosphatase,⁶⁰ and ATP is a good substrate for the enzyme.⁶¹

Clubb and his colleagues raised the serum AP of human volunteers to many times the normal activity by means of enzyme infusions and found no detectable effect on any parameter of calcium or phosphate metabolism, concluding that AP in the circulation appeared to be metabolically inert.⁶² Recent work suggests that intestinal AP is involved in calcium absorption⁶³ and that it may, in fact, be the same enzyme as calcium-activated adenosine triphosphatase.⁶⁴

Infusions of micellar solutions of oleic or octanoic acid into the cannulated rat duodenum result in increased synthesis of AP in the intestinal cell,⁶⁵ whilst similar infusion experiments in man suggest that intestinal alkaline phosphatase plays an active role in the absorption of oleic acid.⁶⁶ The enzyme may also be implicated in the hydrolysis of dietary sphingomyelin to ceramide and water-soluble compounds.⁶⁷ Although there is a good deal of information available about the relationship between fat absorption and intestinal alkaline phosphatase,⁶⁸⁻⁷³ the exact function of the enzyme in this context remains to be elucidated.

Alkaline phosphatases have their optimum reaction in an alkaline medium only at high substrate concentrations; at substrate concentrations approaching the physiological levels of phosphoric esters in the cell, the pH optimum of enzyme activity is close to the physiological pH values⁷⁴ and the enzyme possesses considerable activity even at a pH of 7.2. Hence the so-called 'alkaline' phosphatases can display almost their maximum enzyme activity under the conditions of pH and concentration of phosphate esters existing in the cell. It is clear that the function of the alkaline phosphatases *in vivo* remains unknown, a state of affairs more likely to be due to laboratory ineptitude than to enzymatic impotence.

Alkaline Phosphatase as an Isoenzyme

No definition is universally acceptable for the term isoenzyme; probably a broad definition such as 'different proteins with similar enzymatic activity' is at present the most suitable.⁵ The existence of serum alkaline phosphatase in more than one form was first indicated by paper electrophoresis when serum alkaline phosphatase was reported to migrate mainly in the α 2-globulin position, with a second fainter zone migrating in the position of β -globulin.⁷⁵ Following this, numerous attempts have been made to determine the organ source of serum alkaline phosphatase in health and disease by means of electrophoresis on a variety of media, including starch gel,^{76,77,78} cellulose acetate,⁷⁹ agar gel,⁸⁰ Pevikon C-870⁸¹ and, more recently, acrylamide gel.^{82,83} The results obtained have often been conflicting; in particular, the results obtained on one electrophoretic medium are difficult to relate to those

obtained on another. Even using the same medium, however, conflicting results have been obtained by different workers. Thus it has been claimed that using starch-gel electrophoresis the differentiation between bone and liver alkaline phosphatase can be made on the basis of the minor bands.⁸⁴ Chiandussi *et al* found that whilst the main band in neonates and in patients with bone and liver disease all gave an identical band in the β -globulin region, there were additional minor bands consistently present in the γ -globulin, β -lipoprotein and slow α_2 -globulin regions in cases of biliary obstruction, whilst in patients with bone disease, additional minor bands were seen in the β -lipoprotein and slow α_2 -globulin regions. In contrast, other workers have claimed to make the distinction between bone and liver enzymes by the position of the major bands, the liver band running slightly ahead of the bone band.⁸⁵ Finally, starch-gel electrophoresis has been found by some to be unhelpful in the differential diagnosis of disease.⁷⁸

This confusing and unsatisfactory state of affairs has now been largely resolved by three main factors. First, it is now realized that the different methods of extraction and preparation of the enzyme from tissue or blood will produce variations in electrophoretic pattern.⁸⁶ Secondly, it is now clear that attention should be focused not only on the position of the main band, but also on its shape; whilst the liver band on starch-gel tends to be slightly faster than bone, the difference is small and there is some overlap.⁸⁷ However, as Newton emphasized, the bone band is more diffuse than the compact liver band. Finally, the recent advent of polyacrylamide gel disc electrophoresis, which is known to give a superior resolution of proteins,⁸⁸ has demonstrated a clear and consistent difference in both mobility and shape between the slower, more diffuse bone band and the faster and more compact liver band.^{82, 83} On polyacrylamide gel slab electrophoresis⁸⁸ human serum alkaline phosphatase can be separated into four distinct isoenzymes. Apart from the bone and liver components which comprise the main band, a third isoenzyme, which is identical to intestinal alkaline phosphatase, is found in some individuals, whilst a fourth isoenzyme which remains at the origin is frequently found in patients with liver disease. Earlier claims that the presence of this origin band implied either biliary obstruction or hepatic metastases rather than hepatocellular disease⁸⁷ have not been substantiated. However, extrahepatic obstruction causing a raised alkaline phosphatase without jaundice is usually partial or intermittent and this may occur in chronic pancreatitis or carcinoma of the pancreas. The isoenzyme characteristics of both normal pancreatic alkaline phosphatase and a tumour variant have been recently described;⁸⁹ in chronic pancreatitis, on the other hand, hyperparathyroid bone disease or osteomalacia may coexist and isoenzyme studies will indicate that the raised alkaline phosphatase is not of hepatic, but of bone origin.

Other methods which have been used to study the isoenzyme characteristics of the alkaline phosphatases include immunological techniques, gel chromatography, and examination of heat stability and urea sensitivity. Immunologically, three antigenic classes of alkaline phosphatase exist. The first includes liver, bone, spleen, and kidney; intestinal alkaline phosphatase constitutes the second class and placental alkaline phosphatase the third.⁹⁰ The second and third classes partially cross-react with each other, and in turn cross-react with a minor kidney component. More recently, it has been shown that within the non-placental, non-intestinal group, an antiserum to liver phosphatase did not react with the phosphatases from bone or kidney.⁹¹ It appears that at

least three genetic loci operate in the production of human alkaline phosphatase isoenzymes.

A difference in relative thermostability between human bone and liver alkaline phosphatases was noted by Moss and King,⁹² and similar differences were later found when serum alkaline phosphatase from groups of subjects with skeletal and hepatobiliary disease was subjected to heat inactivation.⁹³ The clinical value of heat inactivation as a simple laboratory technique to differentiate between elevated serum alkaline phosphatase of bone or liver origin has been confirmed;^{94,95} the liver enzyme is relatively more heat stable than bone alkaline phosphatase, although both enzymes are vastly less heat stable than the placental isoenzyme.^{96,97}

Considerable differences have been found in the susceptibility of various human alkaline phosphatases to urea inhibition.^{98,99} The activation of hepatic alkaline phosphatase by low molarities of urea¹⁰⁰ has not proved to be of a sufficient magnitude to provide a clinically useful tool, but the greater urea stability of the liver enzyme compared to that of bone enables the distinction between the two to be readily made, and provides a technique of a similar order of precision and reliability to that obtained with heat inactivation.^{95,101}

It is also possible to differentiate between bone and liver alkaline phosphatase on Sephadex gel filtration; serum phosphatase from controls and patients with osteoblastic bone disease shows one main peak of enzyme activity moving in the 7S gamma-globulin region, whilst patients with raised alkaline phosphatase of hepatic origin show in addition another peak in the 19S macromolecular region (MW > 200 000), which comprises about 30% of the total activity.^{95,102} Techniques depending upon observations of varying molecular size are, however, too cumbersome for routine clinical use.

Intestinal Alkaline Phosphatase

In 1941 Gomori noted a positive reaction for AP on the surface of human intestinal epithelial cells;¹⁰³ this finding was soon confirmed and the reaction localized to the brush border and Golgi area.¹⁰⁴ The alkaline phosphatase of the small bowel brush border of the mouse is not a single entity¹⁰⁵ and the alkaline phosphatase activity of extracts of human small intestine is heterogeneous on DEAE-cellulose column chromatography¹⁰⁶ and on starch-gel electrophoresis.¹⁰⁷ Much information is available about the distribution and development of alkaline phosphatase in the small intestine.^{108,109,110} Intestinal alkaline phosphatase differs from non-intestinal phosphatases in substrate specificity^{111,112} and in electrophoretic mobility.⁸⁴ A significant advance came with the discovery by Fishman's group that intestinal alkaline phosphatase was strongly inhibited by the stereo-specific inhibitor L-phenylalanine, which inhibited the bone and liver isoenzymes to an insignificant extent,^{113,114} and shortly after this Robinson and Pierce demonstrated that human small intestinal AP differed from the other isoenzymes in being unaltered in electrophoretic mobility by neuraminidase,¹¹⁵ and this was quickly confirmed.²⁰ Up to 60% of normal human sera contain a slow-moving small-intestinal band on electrophoresis,¹¹⁶ and the presence of this isoenzyme in serum is known to be genetically controlled. Thus the small-intestinal band is seen more frequently in the serum of subjects who are blood group O or B than in subjects who are blood group A, and for any given blood group it is seen more fre-

quently in secretors than in non-secretors.^{117,118,119,120,121} The appearance of small-intestinal alkaline phosphatase in serum is, furthermore, known to be related to the ingestion of fat.^{71,116}

Intestinal Alkaline Phosphatase in Duodenal Juice

Duodenal juice contains high concentrations of alkaline phosphatase and isoenzyme studies have indicated that this is partly small intestinal in origin and partly derived from bile.¹²² Following the intravenous injection of both secretin and pancreozymin in man, there is a large increase in the output of alkaline phosphatase in duodenal juice, which is only partly due to increased bile flow since isoenzyme studies have shown large increases in concentration and output of intestinal AP after both secretin and pancreozymin.^{122,123} Both hormones have also been shown to release intestinal AP into duodenal juice in the cat.¹²⁴ Secretion of intestinal AP into the bowel lumen can also be produced by perfusion of long-chain fatty acids,^{125,126} and also by glucose.¹²⁷

Causes of an Elevated Serum Alkaline Phosphatase

The recognized causes of a high serum AP include skeletal disease, hepatic disorders, intestinal disease, tumours, pregnancy,^{3,128} thyrotoxicosis,¹²⁹ the ingestion of anticonvulsant drugs,¹³⁰ and rheumatoid arthritis.¹³¹

SKELETAL DISEASE

Robison first described alkaline phosphatase in the ossifying cartilage of rats and rabbits.^{30,31} Histochemically, the enzyme is found in association with osteoblasts,¹³² which are felt to be the source of the elevated levels of alkaline phosphatase found in growing children, as well as in a variety of bone disorders including rickets,¹²⁹ Paget's disease,¹²⁹ hyperparathyroidism,¹³³ familial osteoectasia,¹³⁴ and osteoblastic bone secondaries.¹³⁵ Estimations of serum alkaline phosphatase have been found more helpful in the diagnosis of osteomalacia following gastrectomy by some workers¹³⁶ than by others,¹³⁷ a disagreement probably based on the fact that neither group employed isoenzyme techniques, which should clearly indicate the presence of 'bone' type AP in the serum of patients with osteomalacia, even when the total serum AP level is only slightly elevated. Of the various techniques available for this purpose, heat inactivation studies are the simplest to employ but recent claims of the ability to quantitate roughly the amount of bone enzyme in serum by electrophoretic means are promising.¹³⁸

HEPATIC DISORDERS

The association between high levels of serum alkaline phosphatase and liver disease has been known for many years,¹³⁹ as has the classical concept that high serum levels of AP are found in 'obstructive' jaundice, whilst in 'toxic' or 'catarrhal' jaundice only slight elevations are found.¹⁴⁰ It is now realized that a clear separation of the various types of jaundice into 'obstructive' and 'non-obstructive' is not feasible.¹⁴¹ Thus, whilst it is true that more than 80% of patients with extrahepatic obstruction have a serum alkaline phosphatase of more than 30 KA units per 100 ml, up to 40% of patients with hepatitis also have enzyme levels exceeding this figure.¹⁴² There are two main theories to explain the elevations of serum alkaline phosphatase in liver disease. The

'retention' theory postulates that in biliary obstruction there is a failure of excretion through the biliary tract of alkaline phosphatase, derived from bone, whilst the 'regurgitation' theory considers the increase in serum AP in hepatic disorders to be due to the passage into the circulation of enzyme derived from bile by way of communications between the bile canaliculi and the sinusoids.¹⁴³ The conflicting experimental evidence in favour of each theory has been previously summarized^{1,3} but recent work has come down strongly on the side of the regurgitation theory. Thus, it has been shown that ligation of one hepatic duct in the experimental animal results in increased production of alkaline phosphatase by the liver rather than decreased excretion of the enzyme,¹⁴⁴ and experiments with isolated perfused liver preparations have shown that obstruction to biliary outflow leads to a rise in the AP activity of the perfusate, suggesting increased formation of the enzyme in the liver.¹⁴⁵

In man, experiments in which heat-stable human placental AP was infused into normal subjects and into patients with biliary obstruction suggest that it is unlikely that alkaline phosphatase is excreted via the biliary tract, since patients with biliary obstruction catabolize the infused enzyme at rates similar to those found in normal subjects and infused enzyme cannot be detected in the urine, faeces, or bile of recipients.⁶² Clubb *et al*, therefore, suggest that the reason for the elevation of serum alkaline phosphatase in patients with biliary obstruction is the increased delivery into the circulation of enzyme derived from the cholangiolitic epithelium. Further evidence against the excretion theory comes from isoenzyme studies since, as already mentioned, qualitative differences can be demonstrated between the alkaline phosphatase circulating in patients with hepatic disorders and that found in the serum of patients with osteoblastic skeletal disease. Finally, recent work has thrown light on the mechanism of the rise in serum AP following bile duct ligation.¹⁴⁶ Within twelve hours of bile duct ligation in rats, there is a sevenfold increase in the concentration of AP in the liver, whilst the enzyme concentration in serum increases by two and a half-fold due to an increase in an isoenzyme originating in liver. Since both rises were inhibited by cycloheximide, the rise in hepatic AP activity appeared to be due to protein synthesis *de novo*, in contrast to the rise in serum glutamic pyruvic transaminase which followed bile duct ligation which was due to simple leakage of preformed enzyme from damaged cells.

The poor correlation between serum AP and bilirubin levels in liver disorders is thus a reflection of the differing mechanisms responsible for the accumulation of bilirubin and alkaline phosphatase. In the diagnosis of liver disorders, AP determinations play an important part, particularly when performed in conjunction with isoenzyme studies; from the point of view of prognosis, however, they are of much less help.¹⁴⁷

INTESTINAL DISEASE

Observations of elevated levels of intestinal AP in disease have been few, and to a large extent uncontrolled. Unfortunately, until a healthy control population of differing age groups and of known blood groups and secretor status has been studied, both fasting and also after a standard fat meal, the normal range of intestinal alkaline phosphatase in serum will not be available.

Fishman *et al*¹⁴⁸ first reported that some cirrhotics had increased amounts of intestinal alkaline phosphatase in their serum; though the average value in cirrhosis did not differ from the normals, there was a wider range of values

amongst the cirrhotics (10-80% intestinal component). About one third of cirrhotics had high levels of intestinal AP and in one of these grossly elevated levels were seen. Most of this group with elevated intestinal AP levels had portal hypertension.

Other workers have assayed raised levels of intestinal AP in fasting blood electrophoretically in order to diagnose cirrhosis; it appears that the levels of intestinal AP reached in serum following a fatty meal are genetically controlled, being linked to the ABH system and the ABO blood groups in patients with cirrhosis, as in normals.¹⁴⁹

There is one report¹⁵⁰ of a case of steatorrhoea of unknown cause associated with a raised serum alkaline phosphatase of mainly intestinal origin. However, no definitive diagnosis was possible during life, and no necropsy carried out.

TUMOURS

Alkaline phosphatase has been reported in a variety of tumours and there is evidence that tumours may secrete variant forms of alkaline phosphatase into the circulation.¹⁵¹ Recently, a tumour isoenzyme, named the Regan isoenzyme after the patient in whom it was first found, has been isolated from the serum and tumour tissue of patients with a wide variety of tumours.^{152,153} This tumour isoenzyme can be recognized in serum since it has properties virtually identical to placental AP in being L-phenylalanine sensitive and very heat-stable. It appears to represent another example of the ectopic synthesis by tumour cells of an enzyme protein not normally manufactured in the tissue from which the tumour originated. The ectopic production of hormones such as ADH, and the synthesis by tumour cells of proteins normally found only in foetal tissue, is well recognized. The carcinoembryonic antigen,¹⁵⁴ and alpha-fetoglobulin¹⁵⁵ in the serum of certain patients with hepatoma may, like the Regan isoenzyme, result from the derepression of specific genomes within the tumour cell.¹⁵⁶ The Regan isoenzyme, however, probably represents a special case, since many tumours produce isoenzymes of alkaline phosphatase which are heat-sensitive and phenylalanine-resistant.^{89,157,158}

T. W. WARNES

University College Hospital, London

References

- ¹Gutman, A. B. (1959). Serum alkaline phosphatase activity in diseases of the skeletal and hepatobiliary systems. A consideration of the current status. *Amer. J. Med.*, 27, 875-901.
- ²Wilkinson, J. H. (1966). *Isoenzymes*, pp. 104-120. Lippincott, Philadelphia.
- ³Posen, S. (1967). Alkaline phosphatase. *Ann. intern. Med.*, 67, 183-203.
- ⁴Fishman, W. H., and Ghosh, N. K. (1967). Isoenzymes of human alkaline phosphatase. *Advanc. clin. Chem.*, 10, 255-370.
- ⁵Latner, A. L., and Skillen, A. W. (1968). *Isoenzymes in Biology and Medicine*, pp. 56-63, 158-161. Academic Press, London and New York.
- ⁶Moss, D. W. (1970). The heterogeneity of human alkaline phosphatase. In *Enzymes and Isoenzymes: Structure, Properties and Function* (5th FEBS Meeting, vol. 18), edited by D. Shugar, pp. 227-239. Academic Press, London and New York.
- ⁷Mathies, J. C. (1958). Preparation and properties of highly purified alkaline phosphatase from swine kidneys. *J. biol. Chem.*, 233, 1121-1127.
- ⁸Plocke, D. J., Levinthal, C., and Vallee, B. L. (1962). Alkaline phosphatase of *Escherichia coli*; a zinc metallo-enzyme. *Biochemistry (Wash.)*, 1, 373-378.
- ⁹Fishman, W. H., and Ghosh, N. K. (1967). Influence of reagents reacting with metal, thiol, and amino sites on catalytic activity and L-phenylalanine inhibition of rat intestinal alkaline phosphatase. *Biochem. J.*, 105, 1163-1170.
- ¹⁰Clark, B., and Porteous, J. W. (1965). The metal ion activation of the alkaline β -glycerophosphatase of rabbit small intestine. *Biochem. J.*, 95, 475-482.
- ¹¹Ahmed, Z., and King, E. J. (1960). Kinetics of placental alkaline phosphatase. *Biochim. biophys. Acta (Amst.)*, 45, 581-592.
- ¹²Morton, R. K. (1955). Some properties of alkaline phosphatase of cow's milk and calf intestinal mucosa. *Biochem. J.*, 60, 573-582.

- ¹³Ghosh, N. K., and Fishman, W. H. (1968). Purification and properties of molecular-weight variants of human placental alkaline phosphatase. *Biochem. J.*, **108**, 779-792.
- ¹⁴Engström, L. (1964). The amino-acid sequence around the reactive serine in calf-intestinal alkaline phosphatase. *Biochim. biophys. Acta (Amst.)*, **92**, 79-84.
- ¹⁵Lazdunski, C., and Lazdunski, M. (1966). Étude cinétique du mécanisme d'action catalytique de la phosphatase alcaline d'*Escherichia coli*. *Biochim. biophys. Acta (Amst.)*, **113**, 551-566.
- ¹⁶Schwartz, J. H., Crestfield, A. M., and Lipmann, F. (1963). The amino acid sequence of a tetradecapeptide containing the reactive serine in *E. coli* alkaline phosphatase. *Proc. nat. Acad. Sci. (Wash.)*, **49**, 722-728.
- ¹⁷Engström, L. (1964). Studies on bovine-liver alkaline phosphatase. Purification, phosphate incorporation. *Biochim. biophys. Acta (Amst.)*, **92**, 71-78.
- ¹⁸Schultze, H. E. (1957). Über klinisch interessante körpereigene Polysaccharidverbindungen. *Scand. J. clin. Lab. Invest.*, **10**, Suppl. 31, 135-165.
- ¹⁹Plummer, T. H., Jr., and Hirs, C. H. W. (1964). On the structure of bovine pancreatic ribonuclease B. Isolation of a glycopeptide. *J. biol. Chem.*, **239**, 2530-2538.
- ²⁰Moss, D. W., Eaton, R. H., Smith, J. K., and Whitby, L. G. (1966). Alteration in the electrophoretic mobility of alkaline phosphatases after treatment with neuraminidase. *Biochem. J.*, **98**, 32c-33c.
- ²¹Butterworth, P. J., and Moss, D. W. (1966). Action of neuraminidase on human kidney alkaline phosphatase. *Nature (Lond.)*, **209**, 805-806.
- ²²Ghosh, N. K. (1969). Purification and molecular properties of placental and intestinal alkaline phosphatases. *Ann. N.Y. Acad. Sci.*, **166**, 604-640.
- ²³Portmann, P., Rossier, R., and Chardonnens, H. (1960). Zur Kenntnis der alkalischen Darmphosphatase. II. Untersuchungen über die Homogenität und den chemischen Aufbau der reinen alkalischen Phosphomonoesterase. *Helv. physiol. pharmacol. Acta*, **18**, 414-427.
- ²⁴Engström, L. (1961). Studies on calf-intestinal alkaline phosphatase I. Chromatographic purification, microheterogeneity and some other properties of the purified enzyme. *Biochim. biophys. Acta (Amst.)*, **52**, 36-48.
- ²⁵Ghosh, N. K., Goldman, S. S., and Fishman, W. H. (1967). Human placental alkaline phosphatase; a sialoprotein. *Enzymologia*, **33**, 113-124.
- ²⁶Berenson, G. S., Radhakrishnamurthy, B., Fishkin, A. F., Dessauer, H., and Arquembourg, P. (1966). Individuality of glycoproteins in human aorta. *J. Atheroscler. Res.*, **6**, 214-223.
- ²⁷Gottlieb, A. J., and Sussman, H. H. (1968). Human placental alkaline phosphatase; molecular weight and subunit structure. *Biochim. biophys. Acta (Amst.)*, **160**, 167-171.
- ²⁸Sussman, H. H., and Gottlieb, A. J. (1968). Subunit properties of human placental alkaline phosphatase. *Fed. Proc.*, **27**, 779.
- ²⁹Robison, R. (1923). The possible significance of hexosephosphoric esters in ossification. *Biochem. J.*, **17**, 286-293.
- ³⁰Robison, R., and Soames, K. M. (1924). The possible significance of hexosephosphoric esters in ossification. Part II. The phosphoric esterase of ossifying cartilage. *Biochem. J.*, **18**, 740-754.
- ³¹Martland, M., and Robison, R. (1924). The possible significance of hexosephosphoric esters in ossification. Part V. The enzyme in the early stages of bone development. *Biochem. J.*, **18**, 1354-1357.
- ³²Robison, R., Macleod, M., and Rosenheim, A. H. (1930). The possible significance of hexosephosphoric esters in ossification. IX. Calcification in vitro. *Biochem. J.*, **24**, 1927-1941.
- ³³Robison, R., and Rosenheim, A. H. (1934). Calcification of hypertrophic cartilage in vitro. *Biochem. J.*, **28**, 684-698.
- ³⁴McLean, F. C., Lipton, M. A., Bloom, W., and Barron, E. S. G. (1946). Biological factors in calcification in bone. *Trans. Macy Conf. Metabolic Aspects of Convalescence*, **14**, 9-19.
- ³⁵Henrichsen, E. (1958). Alkaline phosphatase and calcification: histochemical investigations on the relationship between alkaline phosphatase and calcification. *Acta orthopaed. scand., Suppl.* **34**.
- ³⁶Russell, R. G., Bisaz, S., and Fleisch, H. (1969). Pyrophosphate and diphosphonates in calcium metabolism and their possible role in renal failure. *Arch. intern. Med.*, **124**, 571-577.
- ³⁷Wachstein, M., and Bradshaw, M. (1965). Histochemical localization of enzyme activity in the kidneys of three mammalian species during their postnatal development. *J. Histochem. Cytochem.*, **13**, 44-56.
- ³⁸Mizutani, A., and Barnett, R. J. (1965). Fine structural demonstration of phosphatase at pH 9. *Nature (Lond.)*, **206**, 1001-1003.
- ³⁹Chase, W. H. (1963). The demonstration of alkaline phosphatase activity in frozen-dried mouse gut in the electron microscope. *J. Histochem. Cytochem.*, **11**, 96-101.
- ⁴⁰Lobel, B. L., Deane, H. W., and Romney, S. L. (1962). Enzymic histochemistry of the villous portion of the human placenta from six weeks of gestation to term. *Amer. J. Obstet. Gynec.*, **83**, 295-299.
- ⁴¹Done, J., Shorey, C. D., Loke, J. P., and Pollak, J. K. (1965). The cytochemical localization of alkaline phosphatase in *Escherichia coli* at the electron-microscope level. *Biochem. J.*, **96**, 27c-28c.
- ⁴²Kuo, M. H., and Blumenthal, H. J. (1961). Absence of phosphatase repression by inorganic phosphate in some micro-organisms. *Nature (Lond.)*, **190**, 29-31.
- ⁴³Horiuchi, T., Horiuchi, S., and Mizuno, D. (1959). A possible negative feedback phenomenon controlling formation of alkaline phosphomonoesterase in *Escherichia coli*. *Nature (Lond.)*, **183**, 1529-1530.
- ⁴⁴Torriani, A. (1960). Influence of inorganic phosphate in the formation of phosphatases by *Escherichia coli*. *Biochim. biophys. Acta (Amst.)*, **38**, 460-469.
- ⁴⁵Garen, A. (1960). Genetic control of the specificity of the bacterial enzyme, alkaline phosphatase. *Symp. Soc. gen. Microbiol.*, **10**, 239-247.
- ⁴⁶Engström, L. (1961). Studies on calf-intestinal alkaline phosphatase. II. Incorporation of inorganic phosphate into a highly purified enzyme preparation. *Biochim. biophys. Acta (Amst.)*, **52**, 49-59.
- ⁴⁷Bradfield, J. R. G. (1946). Alkaline phosphatase in invertebrate sites of protein secretion. *Nature (Lond.)*, **157**, 876-877.
- ⁴⁸Gold, N. I., and Gould, B. S. (1951). Collagen fiber formation and alkaline phosphatase. *Arch. Biochem.*, **33**, 155-164.
- ⁴⁹Brachet, J., and Jeener, R. (1946). Phosphatase alcaline des noyaux et vitesse de remplacement du phosphore de l'acide thymonucleique. *C.R. Soc. Biol. (Paris)*, **140**, 1121-1123.
- ⁵⁰Jeener, R. (1946). Sur les liens de la phosphatase alcaline avec les nucléoprotéides du noyau cellulaire et des granules cytoplasmiques. *Experientia (Basel)*, **2**, 458-459.
- ⁵¹Dempsey, E. W., and Wislocki, G. B. (1946). Histochemical contributions to physiology. *Physiol. Rev.*, **26**, 1-27.

- ⁵²Gavosto, F., and Pileri, A. (1958). The effect of administration of 6-mercaptopurine on nucleic acids and alkaline phosphatase of regenerating rat liver. *Cancer (Philad.)*, 11, 222-225.
- ⁵³Rubini, J. R. (1963). A role for alkaline phosphatase in controlling D.N.A. synthesis. (Abstr.) *J. clin. Invest.*, 42, 974.
- ⁵⁴Weiss, B., Live, T. R., and Richardson, C. C. (1968). Enzymatic breakage and joining of deoxyribonucleic acid V. End group labelling and analysis of deoxyribonucleic acid containing single strand breaks. *J. biol. Chem.*, 243, 4530-4542.
- ⁵⁵Pflogner, N., Bradley, A., and Fraenkel-Conrat, H. (1967). Action of cyclic phosphodiesterase from *Escherichia coli* on tobacco mosaic virus ribonucleic acid. *Biochim. biophys. Acta (Amst.)*, 142, 105-110.
- ⁵⁶Cox, R. P., and Griffin, M. J. (1965). Pyrophosphatase activity of mammalian alkaline phosphatase. (Letter). *Lancet*, 2, 1018-1019.
- ⁵⁷Fernley, H. N., and Walker, P. G. (1967). Studies on alkaline phosphatase. Inhibition by phosphate derivatives and the substrate specificity. *Biochem. J.*, 104, 1011-1018.
- ⁵⁸Axelrod, B. (1948). A new mode of enzymatic phosphate transfer. *J. biol. Chem.*, 172, 1-13.
- ⁵⁹Morton, R. K. (1958). The phosphotransferase activity of phosphatases. 2. Studies with purified alkaline phosphomonoesterases and some substrate-specific phosphatases. *Biochem. J.*, 70, 139-150.
- ⁶⁰Saev, G. K. (1963). Some concepts of the intracellular function of alkaline phosphatases, based on investigations of the mechanism of their action. *Enzymologia*, 26, 169-175.
- ⁶¹Schlesinger, M. J., and Coon, M. J. (1960). Hydrolysis of nucleoside di- and triphosphates by crystalline preparations of yeast inorganic pyrophosphatase. *Biochim. biophys. Acta (Amst.)*, 41, 30-36.
- ⁶²Clubb, J. S., Neale, F. C., and Posen, S. (1965). The behaviour of infused human placental alkaline phosphatase in human subjects. *J. Lab. clin. Med.*, 66, 493-507.
- ⁶³Norman, A. W., Mircheff, A. K., Adams, T. H., and Spielvogel, A. (1970). Studies on the mechanism of action of calciferol. III. Vitamin D-mediated increase of intestinal brush border alkaline phosphatase activity. *Biochim. biophys. Acta (Amst.)*, 215, 348-359.
- ⁶⁴Hausler, M. R., Nagode, L. A., and Rasmussen, H. (1970). Induction of intestinal brush border alkaline phosphatase by vitamin D and identity with Ca-ATPase. *Nature (Lond.)*, 228, 1199-1201.
- ⁶⁵Glickman, R. M., Alpers, D. H., Drummey, G. D., and Isselbacher, K. J. (1970). Increased lymph alkaline phosphatase after fat feeding: effects of medium chain triglycerides and inhibition of protein synthesis. *Biochim. biophys. Acta (Amst.)*, 201, 226-235.
- ⁶⁶Linscheer, W. G., Malagelada, J. R., and Fishman, W. H. (1971). Diminished oleic acid absorption in man by L-phenylalanine inhibition of an intestinal phosphohydrolase. *Nature (new Biol.)*, 231, 116-117.
- ⁶⁷Nilsson, A. (1969). The presence of sphingomyelin- and ceramide-cleaving enzymes in the small intestinal tract. *Biochim. biophys. Acta (Amst.)*, 176, 339-347.
- ⁶⁸Keiding, N. R. (1964). The alkaline phosphatase fractions of human lymph. *Clin. Sci.*, 26, 291-297.
- ⁶⁹Blomstrand, R., and Werner, B. (1965). Alkaline phosphatase activity in human thoracic duct lymph. *Acta chir. scand.*, 129, 177-191.
- ⁷⁰Inglis, N. I., Krant, M. J., and Fishman, W. H. (1967). Influence of a fat-enriched meal on human serum (L-phenylalanine-sensitive) 'intestinal' alkaline phosphatase. *Proc. Soc. exp. Biol. (N.Y.)*, 124, 699-702.
- ⁷¹Warnock, M. L. (1968). Intestinal phosphatase and fat absorption. *Proc. Soc. exp. Biol. (N.Y.)*, 129, 768-772.
- ⁷²Kleerekoper, M., Horne, M., Cornish, C. J., and Posen, S. (1970). Serum alkaline phosphatase after fat ingestion; an immunological study. *Clin. Sci.*, 38, 339-345.
- ⁷³Malagelada, J. R., Linscheer, W. G., Rufo, M., and Fishman, W. H. (1971). Two directional release of brush border alkaline phosphatase during fat absorption. (Abstr.) *Gastroenterology*, 60, 693.
- ⁷⁴Morton, R. K. (1957). The kinetics of hydrolysis of phenyl phosphate by alkaline phosphatases. *Biochem. J.*, 65, 674-682.
- ⁷⁵Baker, R. W. R., and Pellegrino, C. (1954). The separation and detection of serum enzymes by paper electrophoresis. *Scand. J. clin. Lab. Invest.*, 6, 94-101.
- ⁷⁶Rosenberg, I. N. (1959). Zone electrophoretic studies of serum alkaline phosphatase. *J. clin. Invest.*, 38, 630-644.
- ⁷⁷Keiding, N. R. (1959). Differentiation into three fractions of the serum alkaline phosphatase and the behavior of the fractions in diseases of bone and liver. *Scand. J. clin. Lab. Invest.*, 11, 106-112.
- ⁷⁸Boyer, S. H. (1961). Alkaline phosphatase in human sera and placentae. *Science*, 134, 1002-1004.
- ⁷⁹Korner, N. H. (1962). Distribution of alkaline phosphatase in serum protein fractions. *J. clin. Path.*, 15, 195-199.
- ⁸⁰Haije, W. G., and De Jong, M. (1963). Iso-enzyme patterns of serum alkaline phosphatase in agar-gel electrophoresis and their clinical significance. *Clin. chim. Acta*, 8, 620-623.
- ⁸¹Nordentoft-Jensen, B. (1964). Differentiation between the electrophoretic pattern of serum alkaline phosphatase in bone disease and in liver disease. *Clin. Sci.*, 26, 299-305.
- ⁸²Smith, I., Lightstone, P. J., and Perry, J. D. (1968). Separation of human tissue alkaline phosphatases by electrophoresis on acrylamide disc gels. *Clin. chim. Acta*, 19, 499-505.
- ⁸³Kaplan, M. M., and Rogers, L. (1969). Separation of human serum-alkaline-phosphatase isoenzymes by polyacrylamide gel electrophoresis. *Lancet*, 2, 1029-1031.
- ⁸⁴Chiandussi, L., Greene, S. F., and Sherlock, S. (1962). Serum alkaline phosphatase fractions in hepato-biliary and bone diseases. *Clin. Sci.*, 22, 425-434.
- ⁸⁵Hodson, A. W., Latner, A. L., and Raine, L. (1962). Iso-enzymes of alkaline phosphatase. *Clin. chim. Acta*, 7, 255-261.
- ⁸⁶Moss, D. W. (1962). Iso-enzymes of alkaline phosphatase in autolysed and butanol-extracted liver preparations. *Nature (Lond.)*, 193, 981-982.
- ⁸⁷Newton, M. A. (1967). The clinical application of alkaline phosphatase electrophoresis. *Quart. J. Med.*, 36, 17-28.
- ⁸⁸Raymond, S. (1964). Acrylamide gel electrophoresis. *Ann. N.Y. Acad. Sci.*, 121, 350-365.
- ⁸⁹Warnes, T. W., Timperley, W. R., Hine, P., and Kay, G. (1972). Pancreatic alkaline phosphatase and a tumour variant. *Gut*, 13, 513-519.
- ⁹⁰Boyer, S. H. (1963). Human organ alkaline phosphatases; discrimination by several means including starch gel electrophoresis of antienzyme-enzyme supernatant fluids. *Ann. N.Y. Acad. Sci.*, 103, 938-951.
- ⁹¹Sussman, H. H., Small, P. A., Jr., and Cotlove, E. (1968). Human alkaline phosphatase. Immunochemical identification of organ-specific isoenzymes. *J. biol. Chem.*, 243, 160-166.
- ⁹²Moss, D. W., and King, E. J. (1962). Properties of alkaline-phosphatase fractions separated by starch-gel electrophoresis. *Biochem. J.*, 84, 192-195.
- ⁹³Posen, S., Neale, F. C., and Clubb, J. S. (1965). Heat inactivation in the study of human alkaline phosphatases. *Ann. intern. Med.*, 62, 1234-1243.

- ⁹⁴Fitzgerald, M. X. M., Fennelly, J. J., and McGeeney, K. (1969). The value of differential alkaline phosphatase thermostability in clinical diagnosis. *Amer. J. clin. Path.*, 51, 194-201.
- ⁹⁵Fennelly, J. J., Fitzgerald, M. X., and McGeeney, K. (1969). Value of differential thermostability, urea inhibition and gel filtration of alkaline phosphatase in the identification of disease states. *Gut*, 10, 45-51.
- ⁹⁶McMaster, Y., Tennant, R., Clubb, J. S., Neale, F. C., and Posen, S. (1964). The mechanism of the elevation of serum alkaline phosphatase in pregnancy. *J. Obstet. Gynec. Brit. Cwlt.*, 71, 735-739.
- ⁹⁷Neale, F. C., Clubb, J. S., Hotchkis, D., and Posen, S. (1965). Heat stability of human placental alkaline phosphatase. *J. clin. Path.*, 18, 359-363.
- ⁹⁸Butterworth, P. J., and Moss, D. W. (1966). The effect of urea on human kidney alkaline phosphatase. *Biochem. J.*, 99, 9P-10.
- ⁹⁹Birkett, D. J., Conyers, R. A. J., Neale, F. C., Posen, S., and Brudenell-Woods, J. (1967). Action of urea on human alkaline phosphatases; with a description of some automated techniques for the study of enzyme kinetics. *Arch. Biochem.*, 121, 470-479.
- ¹⁰⁰Bahr, M., and Wilkinson, J. H. (1967). Urea as a selective inhibitor of human tissue alkaline phosphatases. *Clin. chim. Acta*, 17, 367-370.
- ¹⁰¹Horne, M., Cornish, C. J., and Posen, S. (1968). Use of urea denaturation in the identification of human alkaline phosphatases. *J. Lab. clin. Med.*, 72, 905-915.
- ¹⁰²Dunne, J., Fennelly, J. J., and McGeeney, K. (1967). Separation of alkaline phosphatase enzymes in human serum using gel-filtration (Sephadex G-200) techniques. *Cancer (Philad.)*, 20, 71-76.
- ¹⁰³Gomori, G. (1941). Distribution of phosphatase in normal organs and tissues. *J. cell. comp. Physiol.*, 17, 71-83.
- ¹⁰⁴Bourne, G. (1943). The distribution of alkaline phosphatase in various tissues. *Quart. J. exp. Physiol.*, 32, 1-19.
- ¹⁰⁵Moog, F., and Angeletti, P. U. (1962). Immunological difference between duodenal and jejunal phosphatase of the mouse. *Biochim. biophys. Acta (Amst.)*, 60, 440-442.
- ¹⁰⁶Grossberg, A. L., Harris, E. H., and Schlamowitz, M. (1961). Enrichment and separation of alkaline phosphatase activities of human tissues by chromatography on cellulose ion-exchange adsorbents. *Arch. Biochem.*, 93, 267-277.
- ¹⁰⁷Moss, D. W. (1963). Heterogeneity of human intestinal alkaline phosphatase. *Nature (Lond.)*, 200, 1206-1207.
- ¹⁰⁸Moog, F. (1944). Localizations of alkaline and acid phosphatases in the early embryogenesis of the chick. *Biol. Bull.*, 86, 51-80.
- ¹⁰⁹Moog, F. (1961). The functional differentiation of the small intestine. VIII. Regional differences in the alkaline phosphatases of the small intestine of the mouse from birth to one year. *Develop. Biol.*, 3, 153-174.
- ¹¹⁰Moog, F., Etzler, M. E., and Grey, R. D. (1969). The differentiation of alkaline phosphatase in the small intestine. *Ann. N.Y. Acad. Sci.*, 166, 447-465.
- ¹¹¹Nisselbaum, J. S., Schlamowitz, M., and Bodansky, O. (1961). Immunochemical studies of functionally similar enzymes. *Ann. N.Y. Acad. Sci.*, 94, 970-987.
- ¹¹²Keiding, N. R. (1966). Intestinal alkaline phosphatase in human lymph and serum. *Scand. J. clin. Lab. Invest.*, 18, 134-140.
- ¹¹³Fishman, W. H., Green, S., and Inglis, N. I. (1963). L-phenylalanine; an organ specific, stereo-specific inhibitor of human intestinal alkaline phosphatase. *Nature (Lond.)*, 198, 685-686.
- ¹¹⁴Fishman, W. H., and Kreisher, J. H. (1963). Stereospecific, organ-specific inhibition of intestinal alkaline phosphatase. *Ann. N.Y. Acad. Sci.*, 103, 951-955.
- ¹¹⁵Robinson, J. C., and Pierce, J. E. (1964). Differential action of neuraminidase on human serum alkaline phosphatases. *Nature (Lond.)*, 204, 472-473.
- ¹¹⁶Langman, M. J. S., Leuthold, E., Robson, E. B., Harris, J., Luffman, J. E., and Harris, H. (1966). Influence of diet on the 'intestinal' component of serum alkaline phosphatase in people of different A.B.O. blood groups and secretor status. *Nature (Lond.)*, 212, 41-44.
- ¹¹⁷Arfors, K. E., Beckman, L., and Lundin, L. G. (1963). Genetic variations of human serum phosphatases. *Acta genet. (Basel)*, 13, 89-94.
- ¹¹⁸Arfors, K. E., Beckman, L., and Lundin, L. G. (1963). Further studies on the association between human serum phosphatases and blood groups. *Acta genet. (Basel)*, 13, 366-368.
- ¹¹⁹Beckman, L. (1964). Associations between human serum alkaline phosphatases and blood groups. *Acta genet. (Basel)*, 14, 286-297.
- ¹²⁰Shreffler, D. C. (1965). Genetic studies of blood group-associated variations in a human serum alkaline phosphatase. *Amer. J. hum. Genet.*, 17, 71-86.
- ¹²¹Bamford, K. F., Harris, H., Luffman, J. E., Robson, E. B., and Cleghorn, T. E. (1965). Serum-alkaline-phosphatase and the ABO blood-groups. *Lancet*, 1, 530-531.
- ¹²²Warnes, T. W., Hine, P., and Kay, G. (1969). Alkaline phosphatase in duodenal juice following secretin and pancreozymin. (Abstr.) *Gut*, 10, 1049.
- ¹²³Warnes, T. W., Hine, P., and Kay, G. (1970). The action of secretin and pancreozymin on small intestinal alkaline phosphatase (Abstr.) In *4th World Congress of Gastroenterology, Copenhagen*, edited by P. Riis, P. Anthonisen, and H. Baden. p. 131, published by DIS Congress Service, DK-1159 Copenhagen K.
- ¹²⁴Warnes, T. W., Hine, P., and Kay, G. (to be published).
- ¹²⁵Linscheer, W. G., Pihl, O., Wehr, H., and Fishman, W. H. (1970). Enzyme release from human duodenal mucosa; effects of chain length of infused fatty acids. (Abstr.) *Gastroenterology*, 58, 970.
- ¹²⁶Bluestein, M., Malagelada, J., Linscheer, W., and Fishman, W. H. (1970). Enzymomorphology of alkaline phosphatase of rat intestine following fatty acid infusion. (Abstr.) *J. Histochem. Cytochem.*, 18, 679.
- ¹²⁷Linscheer, W. G., Blum, A. L., Inglis, N. I., and Fishman, W. H. (1967). Intestinal alkaline phosphatase during glucose absorption in man. (Abstr.) *Clin. Res.*, 15, 238.
- ¹²⁸Cayla, J., and Fabre, F. (1935). La phosphatase sérique pendant la gestation. *C.R. Soc. Biol.*, 120, 748-750.
- ¹²⁹Kay, H. D. (1932). Phosphatase in growth and disease of bone. *Physiol. Rev.*, 12, 384-422.
- ¹³⁰Richens, A., and Rowe, D. J. F. (1970). Disturbance of calcium metabolism by anticonvulsant drugs. *Brit. med. J.*, 4, 73-76.
- ¹³¹Cockell, R., Kendall, M. J., Becker, J. F., and Hawkins, C. F. (1971). Serum biochemical values in rheumatoid disease. *Ann. rheum. Dis.*, 30, 166-170.
- ¹³²Teaford, M. E., and White, A. A. (1964). Alkaline phosphatase and osteogenesis in vitro. *Proc. Soc. exp. Biol. (N.Y.)*, 117, 541-549.
- ¹³³Dent, C. E., and Harper, C. M. (1962). Plasma-alkaline-phosphatase in normal adults and in patients with primary hyperparathyroidism. *Lancet*, 1, 559-563.

- ¹³⁶Stemmermann, G. N. (1966). An histologic and histochemical study of familial osteoectasia. *Amer. J. Pathol.*, **48**, 641-651.
- ¹³⁷Woodard, H. Q. (1953). Changes in blood chemistry associated with carcinoma metastatic to bone. *Cancer (Philad.)*, **6**, 1219-1227.
- ¹³⁸Thompson, G. R., Neale, G., Watts, J. M., and Booth, C. C. (1966). Detection of vitamin-D deficiency after partial gastrectomy. *Lancet*, **1**, 623-626.
- ¹³⁹Morgan, D. B., Paterson, C. R., Woods, C. G., Pulvertaft, C. N., and Fourman, P. (1965). Search for osteomalacia in 1228 patients after gastrectomy and other operations on the stomach. *Lancet*, **2**, 1085-1088.
- ¹⁴⁰Canapa-Anson, R., and Rowe, D. J. F. (1970). Electrophoretic separation of tissue-specific serum alkaline phosphatases. *J. clin. Path.*, **23**, 499-508.
- ¹⁴¹Roberts, W. M. (1930). Variations in the phosphatase activity of the blood in disease. *Brit. J. exp. Path.*, **11**, 90-95.
- ¹⁴²Roberts, W. M. (1933). Blood phosphatase and the Van den Bergh reaction in the differentiation of the several types of jaundice. *Brit. med. J.*, **1**, 734-738.
- ¹⁴³Sherlock, S. (1962). Jaundice. *Brit. med. J.*, **1**, 1359-1366.
- ¹⁴⁴Becker, K. L., and Stauffer, M. H. (1962). Evaluation of concentrations of serum alkaline phosphatase in hepatitis and obstructive jaundice. *Amer. J. med. Sci.*, **243**, 222-227.
- ¹⁴⁵Birns, M., Masek, B., and Auerbach, O. (1962). The effects of experimental acute biliary obstruction and release on the rat liver. *Amer. J. Path.*, **40**, 95-111.
- ¹⁴⁶Polin, S. G., Spellberg, M. A., Teitelman, L., and Okumura, M. (1962). The origin of elevation of serum alkaline phosphatase in hepatic disease. *Gastroenterology*, **42**, 431-438.
- ¹⁴⁷Sebesta, D. G., Bradshaw, F. J., and Prockop, D. J. (1964). Source of the elevated serum alkaline phosphatase activity in biliary obstruction; studies utilizing isolated liver perfusion. *Gastroenterology*, **47**, 166-170.
- ¹⁴⁸Kaplan, M. M., and Righetti, A. (1970). Induction of rat liver alkaline phosphatase; the mechanism of the serum elevation in bile duct obstruction. *J. clin. Invest.*, **49**, 508-516.
- ¹⁴⁹West, M., and Zimmerman, H. J. (1959). Serum enzymes in hepatic disease. *Med. Clin. N. Amer.*, **43**, 371-400.
- ¹⁵⁰Fishman, W. H., Inglis, N. I., and Krant, M. J. (1965). Serum alkaline phosphatase of intestinal origin in patients with cancer and with cirrhosis of the liver. *Clin. Chim. Acta*, **12**, 298-303.
- ¹⁵¹Klein, U. E., Drube, H. C., and Hansen, H. T. (1967). Nahrungsabhängige Verschiebung und klinisch diagnostische Bedeutung der alkalischen Dünndarm-phosphatase bei Lebercirrhosen. *Verh. dtsch. Ges. inn. Med.*, **73**, 241-244.
- ¹⁵²Dent, C. E., Norris, T. S. M., Smith, R., Sutton, R. A. L., and Temperley, J. M. (1968). Steatorrhea with striking increase of plasma-alkaline-phosphatase of intestinal origin. *Lancet*, **1**, 1333-1336.
- ¹⁵³Lancet (1969). Ectopic alkaline-phosphatase production by tumours (annotation). *Lancet*, **2**, 1236-1237.
- ¹⁵⁴Fishman, W. H., Inglis, N. I., Green, S., Anstiss, C. L., Ghosh, N. K., Reit, A. E., Rustigian, R., Krant, M. J., and Stolbach, L. L. (1968). Immunology and biochemistry of Regan isoenzyme of alkaline phosphatase in human cancer. *Nature (Lond.)*, **219**, 697-699.
- ¹⁵⁵Fishman, W. H., Inglis, N. I., Stolbach, L. L., and Krant, M. J. (1968). A serum alkaline phosphatase isoenzyme of human neoplastic cell origin. *Cancer Res.*, **28**, 150-154.
- ¹⁵⁶Gold, P., and Freedman, S. O. (1965). Specific carcinoembryonic antigens of the human digestive system. *J. exp. Med.*, **122**, 467-481.
- ¹⁵⁷Tatarinov, I. S. (1964). Presence of embryonal alpha-globulin in the serum of patients with hepatocellular carcinoma. (Russian.) *Vop. med. Khim.*, **10**, 90-91.
- ¹⁵⁸Stolbach, L. L., Krant, M. J., and Fishman, W. H. (1969). Ectopic production of an alkaline phosphatase isoenzyme in patients with cancer. *New Engl. J. Med.*, **281**, 757-762.
- ¹⁵⁹Timperley, W. R. (1968). Alkaline-phosphatase-secreting tumour of lung. (Letter.) *Lancet*, **2**, 356.
- ¹⁶⁰Timperley, W. R., and Warnes, T. W. (1970). Alkaline phosphatase in meningiomas. *Cancer (Philad.)*, **26**, 100-103.