

Converts to Noxyflex are legion among concerned clinicians and senior sisters throughout the country. To them we do not preach. To others we speak of the singular effectiveness of Noxyflex against the intrusion of bacteria. When Noxyflex is applied directly to the site of existing or threatened infection (irrespective of sensitivity to any other anti-bacterial agent) its performance is impressive. Our literature carries the weight of extensive clinical evaluation. Have you copies?

Its uses, and users, increase.

**Noxyflex**

**Geistlich Chester**

250. NoxyHolln and 10 mgm. Amethocaine HCl

NEW EDITION OF  
A SURGERY CLASSIC...

# ABDOMINAL OPERATIONS

Sixth Edition

Edited by **RODNEY MAINGOT, F.R.C.S. (Eng.)**, Honorary Consultant Surgeon, Royal Free Hospital, Royal Waterloo Hospital, London; Southend General Hospital, Essex; Sydney Body Gold Medalist; Former Member, Council, and President, Section of Surgery, Royal Society of Medicine; Editor-in-Chief, *British Journal of Clinical Practice*. With 97 contributors.

1974 2,240 pages (two-volume set), illus.

For over 30 years, Maingot's **ABDOMINAL OPERATIONS** has been the outstanding work in its field. It has received widespread critical acclaim:

"... an authoritative and comprehensive discussion of abdominal surgery. The book will be useful to all physicians who see patients with intraabdominal disease..." *(Gastroenterology)*

"... a great surgical treatise, ideally suited for all surgeons, young and old, interested in the diseases of the abdominal cavity. The surgical profession can be greatly thankful to the author for this new work..." *(New England Journal of Medicine)*


"This work, which is comprehensive, factual, and profusely illustrated, is a great credit to British and American surgery and should be widely read by surgeons all over the world... It is by far the best work which has so far been published on abdominal operations." *(British Journal of Surgery)*

The new Sixth Edition has been considerably revised and up-dated to keep pace with the advances made in abdominal surgery during the five years since the publication of the Fifth Edition. More than 20 new chapters and chapter sections and more than 600 new illustrations have been added, including several new color plates.

New material covers such subjects as: Fibre-Optic Endoscopy; Total Pancreatectomy; Sphincteroplasty in the Prophylaxis and Treatment of Residual Common Duct Stones; Neoplasms of Liver; Transplantation of the Liver in Human Subjects; Mesh Prosthesis in the Repair of Large Incisional Hernias; Hirschsprung's Disease: Choice of Operation; Endoscopic and Biopsy Examination of the Colon.

With 111 chapters grouped into 15 sections according to the site of the surgery, this new edition of **ABDOMINAL OPERATIONS** will maintain the same high standards of quality as its predecessors.

Price: (2-vol. set) £42.90 SBN:8385-0041-2



#### Outstanding features:

- An editor of the stature of Rodney Maingot, who not only organizes the entire work and carefully edits each chapter, but contributes a great deal of his own writing.
- Ninety-seven of the most prominent abdominal surgeons as contributors.
- Over 1,500 illustrations, with drawings by the outstanding medical artists in the world.
- A complete and explicit account of the technique of all the approved abdominal operations practiced today in Great Britain and the United States.
- A thorough discussion of choice of operation, pre- and post-operative care, difficulties and dangers, complications, and recent progress.

**SECTIONS:** Abdominal Incisions. Stomach and Duodenum. Spleen. Pancreas. Gallbladder and Bile Ducts. Liver. Abdominal Vascular Surgery. Appendix. Peritoneum: Retroperitoneum: Mesentery: Omenta. Hernia. Small Intestine and Colon. Cancer of the Colon. Tumors of the Rectosigmoid, Rectum and Anal Canal. Some Nonmalignant Lesions of the Ano-rectal Region. Special Subjects.

## APPLETON-CENTURY-CROFTS

MEDICAL/NURSING PUBLISHERS






A DIVISION OF



## Prentice/Hall International

Available through your bookseller or send payment with order, for direct postage-free delivery, from:

Dept. PP1,  
INTERNATIONAL BOOK DISTRIBUTORS, LTD.,  
66 Wood Lane End,  
Hemel Hempstead, Herts. HP2 4RG,  
England.



# *ACM endoscopes look good from every angle*

There's a whole world of very good reasons why you should pick the ACM F8 Panendoscope. Below are just a few. If you would like more and would like to take a good look at an ACM Panendoscope yourself, please use the coupon. Or telephone us.

- \* The ACM Panendoscope will handle more than 90% of your routine upper G.I. investigations.
- \* Ultra-sharp image is circular, three times the size of competitors' instruments.
- \* 12X magnification covers a full, 75° field making those "difficult" areas easy to find and examine.
- \* The distal tip will retroflex in excess of 180° – and even at maximum deflection you can still easily manoeuvre biopsy forceps.
- \* ACM makes endoscopy easier on the patient too – the small, smooth distal tip has the same 12.5mm diameter as the shaft.

- \* And it's very rugged and easy to use – a single 'joystick' manoeuvres the poly-directional tip to any position.
- \* Plus, of course, ACM endoscopes are supported by a truly fast and efficient service network.

Please tell me more about ACM endoscopes.	<input type="checkbox"/>
I would like literature only _____	
Please arrange a visit _____	<input type="checkbox"/>
My telephone number is _____	
Name _____	
Address _____	
_____	G/12/74



**ACM Endoscopes – clearly the best**  
 Wappler Endoscopy Limited,  
 Park House, 22 Park Street, Croydon, Surrey, CRO OYH  
 Telephone: 01-686 4241



ACM endoscopes for gastro-enterology, bronchoscopy, urology, gynaecology and teaching.

# Positive progress against gram-negative infection



**A new and potent  
bactericidal agent against  
many of the more difficult  
gram-negative organisms**

**OBRACIN<sup>TM</sup>**  
**tobramycin sulphate**

**a new parenteral antibiotic for hospital use**

OBRACIN is indicated in the treatment of serious infections caused by susceptible organisms, including entero-bacteria, staphylococci and *Pseudomonas spp.* Effectiveness extends to some strains resistant to other antibiotics. Side-effects are few, although unduly high plasma levels may adversely affect kidney or eighth nerve function. Dosage should be adjusted to the needs of patients with impaired renal function.

## OBRACIN tobramycin sulphate

a new parenteral antibiotic for hospital use

### Prescribing Information

#### Dosage and Administration

**In Normal Renal Function:** All age groups 3 to 5 mg/kg per day in divided dose. The higher dose will only be required in patients with life-threatening infections and should be reduced as soon as possible. Intravenous infusions are usually given in a concentration of 1 mg/ml. For dosage in impaired renal function see packaging leaflet and data sheet.

Appropriate culture and susceptibility tests are recommended to determine the susceptibility of the causative organism to OBRACIN. OBRACIN is *not* for intrathecal use.

#### Contra-indications:

- Hypersensitivity to the drug.
- Should not be used in pregnancy except in life-threatening situations.

#### Warnings:

- OBRACIN is not for intrathecal or intracisternal use.
- Serum concentrations above 12 mcg/ml introduce risk of vestibular, auditory and renal toxicity.
- Concurrent administration of other potentially nephrotoxic or ototoxic drugs may result in additional risk to the eighth nerve and to renal functions.  
e.g. Potent diuretics or other aminoglycoside antibiotics.
- Neuromuscular blockade and respiratory paralysis can result from administration of aminoglycosides, especially when administered to patients who have received muscle relaxants during anaesthesia.

#### Side-effects:

The risk of side-effects is low in patients with normal renal function when the duration of therapy is not prolonged and recommended dosage is not exceeded. Elderly patients, patients with diminished renal function and those treated with larger than recommended doses are more at risk.

Reactions typical of the aminoglycoside antibiotics have been reported on the vestibular and auditory branches of the eighth nerve, such as vertigo, nystagmus, tinnitus and diminution of hearing. Increased azotaemia and oliguria have been reported. Other repeated side-effects possibly related to therapy with OBRACIN include urticaria, maculopapular rash, eosinophilia, altered liver function tests, and decreased platelet and white blood cell counts.

#### Presentation:

OBRACIN is supplied as ready-prepared injection in the following strengths:

**40 mg base in 1 ml**

Each ml contains 40,000 units tobramycin (as sulphate) equivalent to 40 mg base.

**80 mg base in 2 ml**

Each ml contains 40,000 units tobramycin (as sulphate) equivalent to 40 mg base.

**20 mg base in 2 ml**

Each ml contains 10,000 units tobramycin (as sulphate) equivalent to 10 mg base.

Boxes of 5 vials

#### Product Licence Number:

40 mg per 1 ml 0006/0084  
10 mg per 1 ml 0006/0085.

OBRACIN is the trade mark for Tobramycin sulphate, Dist.




Further information is available from  
Dist. Products Limited, Liverpool L24 9LN.  
PB74/JULY 74

**“It is concluded that maintenance treatment of ulcerative colitis with sulphasalazine (salazopyrin) should be continued indefinitely unless contraindicated by side effects”<sup>1</sup>**

The results of the above controlled trial carried out at the Nuffield Department of Clinical Medicine, Radcliffe Infirmary, Oxford, are all the more welcome as earlier trials of cortisone<sup>2</sup> and prednisone<sup>3</sup> at standard dosages have shown them to be totally ineffective in reducing the number of recurrences of ulcerative colitis.

“Fortunately, Sulphasalazine tablets, 0.5 grams, 4 times a day will prevent relapses in the majority of patients with colitis, and only a few patients cannot tolerate this relatively small dose, which can be continued indefinitely, since we do not know when, if ever, it can be safely stopped”<sup>4</sup>.

### “A striking difference in the relapse rate”.



“The patients who received dummy tablets had more than four times the relapse rate of those receiving sulphasalazine”<sup>1</sup> (Salazopyrin).

Salazopyrin (sulphasalazine) is available as the plain 0.5 g. tablet, 0.5 g. EN-tab and as an 0.5 g. suppository. Literature and detailed information on Salazopyrin are available on request.

#### References

- Gut (1973) 14 923-926
- Brit. med. J. (1959) 1 387-394
- Lancet (1965) i 188-189
- General Practitioner (1972) April 7 p11.

## Salazopyrin—

to be continued indefinitely

Salazopyrin is a registered trade mark.

Further information on request from:—

Pharmacia (Great Britain) Ltd.,  
Paramount House,  
75 Uxbridge Road,  
London W5 5SS.  
Telephone: 01-579 0102/7



Pharmacia



Every  
picture  
tells a  
story...

Baritop 100 gives the quality and accuracy expected of Double Contrast X Rays.

"...Small gastric carcinomas of the order of 2-3 cm can be demonstrated, benign and malignant disease can be distinguished and the surface changes in such conditions as chronic erosive gastritis and Menetriere's disease can be visualised as well as the small 'ulcers' following endoscopic biopsies".

KREEL et al, Clin, Radiol. (1973) 24 307.

# Baritop 100

Baritop 100 is an elegant, palatable radiological barium suspension at 100% w/v in convenient 300 ml. cans.\*Baritop Effervescent tablets for use with Baritop 100 are available.

Full information is available on request from:

 Concept Pharmaceuticals Ltd.  
Rickmansworth WD3 1EZ

\*Baritop is a registered trade mark.

# The single solution

for parenteral nutrition



## Aminoplex 5

1 litre three times per day provides 15g. of utilisable nitrogen and 3000K calories; the nutrient content being derived from synthetic L-Form amino acids, sorbitol and alcohol.

### Geistlich Chester

*In co-operation with Salvia-Werk  
G.m.b.H., Homburg/Saar, W. Germany.*

Medical literature  
available from:  
**Geistlich Sons Ltd.,**  
Newton Bank,  
Long Lane,  
Chester CH2 3QZ.  
Telephone  
Chester 47534-7





No fear of a relapse with this patient. When it's gastric or duodenal ulcer, prompt initiation of Caved-S therapy gets them back to normal without delay.

Caved-S contains deglycyrrhizinated liquorice which has become established as a first line treatment for the symptomatic relief of peptic ulcer pain, because it heals.

The unique formulation of Caved-S promotes the healing of gastric and duodenal mucosa in a positive way, therefore dispensing with the need for an additional antacid regimen.

Caved-S also allows flexibility of dosage, enabling you to treat the chronic or severe ulcer patient without the risk of untoward effects.

Full prescribing information is available on request.

*Thanks to*

**CAVED-S**  
**the ulcer fighter**





# Digestion

International Journal of Gastroenterology

## Contents Vol. 10, No. 2, 1974

<i>Hansky, J.; Soveny, C., and Korman, M.G.</i> (Melbourne): Studies with Two Gastrin Antisera of Different Specificity for Gastrins I and II .....	97
<i>Hemmati, M.; Abtahi, F.; Farrokhsiar, M., and Djahanguiri, B.</i> (Tehran): Prevention of Restraint and Indomethacin-Induced Gastric Ulceration by Bile Duct or Pylorus Ligation in Rats .....	108
<i>Okabe, S.; Nakamura, K.; Takeuchi, K., and Takagi, K.</i> (Tokyo): Effects of Antacid, Antipeptic and Anticholinergic Agents on the Development and Healing of Duodenal Ulcers in the Rat .....	113
<i>Gasbarrini, G.; Miglio, F.; Serra, M.A., and Bernardi, M.</i> (Bologna): Immunological Studies of the Jejunal Mucosa in Normal Subjects and Adult Celiac Patients .....	122
<i>Maillard, J.N.; Clot, P., and Coste, T.</i> (Colombes): Preoperative Parameters Influencing Survival in Patients with Elective Portacaval Shunts .....	129
<i>Konturek, S.J.; Tasler, J., and Obtutowicz, W.</i> (Cracow): Characteristics of Inhibition of Pancreatic Secretion by Glucagon .....	138
<i>Danielsson, Å.</i> (Umeå): Effects of Nutritional State and of Administration of Glucose, Glibenclamide or Diazoxide on the Storage of Amylase in Mouse Pancreas .....	150
Recent Advances	
<i>Feldmann, G.; Bignon, J., and Chahinian, P.</i> (Clinchy/Paris): The Liver in $\alpha_1$ -Antitrypsin Deficiency .....	162
Book Reviews .....	175



S. Karger  
Basel · München  
Paris · London  
New York · Sydney

Please complete the attached order form and

return it to:

S. Karger AG, Arnold-Böcklin-Strasse 11

CH-4002, Basel, Switzerland

- Please enter my subscription
- Please send me a specimen copy

Name \_\_\_\_\_

Address \_\_\_\_\_

SI 74007