Progress report

Radioimmunoassay of intestinal hormones

Alimentary diseases are common yet often their aetiology is unknown. This ignorance reflects a basic lack of understanding of normal gut function. One area hitherto little studied is intestinal endocrinology although the bowel is known to be an important endocrine organ. Histologists have agreed that there are at least 11 well defined endocrine cell types in the mucosa¹, presumably all producing different hormones. Why is progress in this field so slow? The answer lies in the great technical difficulty in extracting labile peptide hormones from cells scattered sparsely in the gut mucosa and surrounded by highly destructive proteolytic enzymes. The problems of low concentration and rapid degradation are so formidable that nearly 60 years of continuous work were required to purify the first hormone, secretin², and even then many thousands of hogs had to be sacrificed to yield a few milligrams of hormone. The more classical endocrine tissues of the body are gathered together as glands and extracts can easily be made to test hormone actions. In the case of gut hormones their actions have to be elucidated first. Only then is it possible to purify a single peptide from the mass of material extractable from the gut, using its assayable biological property as the reference. If a hormone lacks a well defined biological action it cannot be purified. Thus histologists can still point to several endocrine cells in the gut unassociated as yet with any known hormone.

Great progress has been made in the last decade. As well as the success in isolating secretin², gastrin³, and cholecystokinin-pancreozymin (CCK-PZ)⁴, three new substances,—motilin⁵, gastric inhibitory peptide (GIP)⁶, and vasoactive intestinal peptide (VIP)⁷—have been purified. Once a hormone has been isolated rapid advances can be made in its study. One is no longer dependent on bioassays for information. The bioassay is confusing because it measures only effects which may be produced by more than one hormone. It is laborious to perform and often too insensitive to measure plasma levels. A purified hormone allows measurement by radioimmunoassay. This technique is specific for the particular hormone and is so easy to perform that hundreds of specimens can be analysed in a day.

Radioimmunoassay Technique

When the plasma concentration of any hormone is measured by radioimmunoassay in several different laboratories the answers usually differ widely. This reflects the fact that many technical problems still occur with this type of assay. The requirements to set up an assay are basically simple. A quantity of semipure hormone is coupled to a carrier protein mixed with Freund's adjuvant and injected into rabbits. After about three months a few animals will produce the required antibodies. Iodine 125 is attached to a tyrosine

in the hormone to make the radioactive hormone tracer. Unknown samples are then read off against the hormone standards by their ability to block the binding of the radioactive hormone tracer to a small fixed quantity of antibody.

There are three important causes of error in the current radioimmunoassay: methodological errors, factors affecting antibody binding, and factors of antibody specificity. The methodological errors include failure to prevent hormone degradation in blood samples and during assay incubation. Another cause of error is the use of bad radioactive tracer hormone. This can be because the hormone contains an impurity which is iodinated or because the act of iodinating the hormone itself causes significant chemical alteration. Secondly, factors affecting antibody binding may cause error. The answer in a radioimmunoassay is given by the percentage of radioactive hormone bound to antibody; thus anything that affects antibody binding in an unknown sample is read as if it were the hormone. For example, urea depresses antibody binding and so a uraemic patient's plasma may wrongly appear to have a high concentration of hormone. Other substances acting in this way are plasma proteins and haemoglobin, for example, in a haemolysed sample. In addition to these, there are unknown factors present in plasma which also very significantly alter antibody binding. It is such unknown factors which bedevil the measurement of low hormone values. The third main cause of error in radioimmunoassay is antibody specificity. Antibodies may have too broad a specificity. For example, an antigastrin antibody may cross react with, and therefore measure, cholecystokinin-pancreozymin. Antibodies may also have too narrow a specificity and measure biologically inactive fragments more potently than the whole hormone. Antisera are often raised to hormone preparations containing degraded material and then react avidly with degraded hormone.

Thus, while much useful new data can be provided by radioimmunoassay of intestinal hormones, care is needed in its interpretation. There is often an element of error in the measurement and only a single antigenic part of the hormone is being assessed and not necessarily the biological potency.

Individual Hormones

INTESTINAL GASTRIN

The duodenal mucosa of man has been found to contain almost as much radioimmunoassayable gastrin as the gastric antral mucosa⁸. Gastrin has been found to exist in two major forms⁹. The classical form, whose sequence¹⁰ and synthesis¹¹ was achieved in 1964, has 17 amino acids and a molecular weight of 2200 and is now known as little gastrin or G 17. Synthetic little gastrin is freely available from ICI Ltd, so that many centres have been able to set up radioimmunoassays and several commercial assay kits are available. The second major form of gastrin has 34 amino acids and a molecular weight of about 4000 and is known as big gastrin or G34¹². Big gastrin contains the entire sequence of little gastrin¹³ so that it is usually fully measured by assays for little gastrin¹⁴. About half of the duodenal gastrin is little gastrin and half big, whereas the gastric antrum contains nearly four-fifths little gastrin⁸. Significant quantities of gastrin are also present in the jejunum where the proportion of big gastrin is very large^{8,15}. Recent assay data have demonstrated the presence of three other minor components of gastrin plasma, an even larger gastrin

(big big gastrin)¹⁵, an intermediate big gastrin, and a little little gastrin¹⁶. The 34 amino acid big gastrin has been found to have approximately equal molar potency with little gastrin for acid production, but has a slower onset and longer duration of action¹⁷. It has also been found that big gastrin has a longer half life in the circulation¹⁷ and thus forms a large proportion of the fasting plasma immunoreactive gastrin¹⁴. As most of big gastrin is found in the small intestine it is perhaps not surprising that patients after antrectomy still have significant levels of gastrin in the circulation and that levels are higher when food passes through the duodenum, as in a Billroth I gastrectomy, than when it is bypassed, as in a Billroth II or Polya gastrectomy¹⁸. The mode of release and the physiological role of the intestinal gastrins, however, are still incompletely understood.

SECRETIN

Secretin is a member of the secretin, glucagon, enteroglucagon, VIP, and GIP family of peptides. These hormones have markedly similar amino acid sequences¹⁹ and it is likely that they have all been evolved from a single ancestral hormone. Their actions overlap and, for example, at various dose levels they inhibit gastric acid^{20,21,22,23}; several members stimulate intestinal secretion²² and several stimulate insulin release^{24,25,26,27}. Fortunately an antiserum raised to one shows little tendency to cross react with other members of the group, which may be partly because of their very different charge configurations. The exception to this is, of course, enteroglucagon whose discovery was entirely due to cross reaction with pancreatic glucagon antibodies²⁸.

Although pure secretin has been available since 1961² and synthetic secretin since 1964²⁹, its radioimmunoassay has been hindered by the lack of tyrosine, an amino acid necessary for coupling to ¹²⁵I to make the hormone radioactive for use as a tracer. The problem has been overcome either by use of synthetic secretin with a tyrosine put in³⁰, or by the iodination of the N terminal histidine in natural secretin^{31,32}, relatively straightforward procedure.

The first published radioimmunoassay reported extremely high plasma secretin concentrations after stimulation, and, in particular, secretin release after oral glucose³³. Several biological studies, however, have failed to demonstrate any increase in the flow of pancreatic juice after glucose^{34,35,36}, and subsequent radioimmunoassays have not recorded any plasma secretin rise with glucose^{37,38,39}. The fasting plasma levels of secretin reported are variable, eg, 467 ± 150 (SD) pg/ml³¹ or well below 100 pg/ml³⁰, but it has been demonstrated that intraduodenal acid provokes a moderate secretin rise^{30,31}. Secretin assay of primate bowel extracts show the greatest total quantity to be in the jejunum but that the duodenal mucosa has the highest secretin concentration⁴⁰, which parallels bioassay results^{41,42}. Preliminary studies show that intraduodenal infusions of HCl in patients with duodenal ulcer releases only about half as much secretin as in control subjects⁴³. This raises the possibility that in duodenal ulceration there may be a primary or secondary failure to release the upper intestinal hormones which inhibit gastric acid production and stimulate alkalinization of the duodenum. Infusion of highly purified porcine secretin, just sufficient to inhibit pentagastrin-stimulated gastric acid (62 ng/kg/10 min), produces a plasma secretin concentration much higher than that resulting from endogenous secretin release induced by intraduodenal infusion of 40 ml 0.1 NHCl, which markedly inhibits gastric acid. Thus if por-

cine and human secretin are equivalent, this would suggest that endogenous secretin may not be an important factor in gastric acid inhibition.

CHOLECYSTOKININ-PANCREOZYMIN

A number of radioimmunoassays have now been reported for cholecystokinin-pancreozymin^{40,44,45,46,47,48} (CCK-PZ) but few of these appear to give reasonable values for human plasma concentrations. The lack of success is partly due to difficulty in iodinating CCK-PZ's sulphated tyrosine and partly due to its poor antigenicity. A major problem is the difficulty in obtaining enough purified CCK-PZ for an adequate immunization programme. This is due to the poor yield of CCK-PZ from hog intestine which is only just over half that of secretin because many more steps are required for purification⁴⁹. Great difficulties have also been encountered in the attempted synthesis of the whole hormone, although the C terminal octapeptide, which has the full range of biological activity⁵⁰, was synthesized quite early.

There has been a tendency for initial assays of a hormone to indicate higher plasma concentrations than those published subsequently. In retrospect, when considerable correlative data have been reviewed, the higher values have usually been seen to be in error. It is thus helpful to have criteria to assess what plasma concentrations could be expected, and to treat with reserve values reported greatly in excess of these. In the case of CCK-PZ (and indeed of gastrin and secretin) a great deal is known of the biological effect produced by the administration of various amounts of hormone. A 30-minute infusion of either 0.5 or 1.0 Ivy dog units CCK-PZ/kg produces contraction of the human gallbladder similar to that observed radiographically after a large fat meal⁵¹. If a 10% distribution space and half life in the circulation of five minutes is assumed the steady state plasma increment achieved by the above infusion can be calculated to be approximately 278 and 556 pg/ml respectively (3000 IDU per mg pure CCK-PZ⁴⁹). Under natural conditions hormone levels much lower than this might be fully effective since other factors such as vagal reflexes and the simultaneous release of facilitatory hormones may make the receptor organ more sensitive.

The earliest reported immunoassay found CCK-PZ undetectable in fasting human plasma but a peak increment of 107 to 200 pg/ml occurred after a fatty meal⁴⁴ (1280 to 2400 μ U/ml reported and assuming 12 000 CHRU/mg pure CCK-PZ). Subsequently rather higher values have been reported, for example, basal levels well over 1 ng/ml⁴⁶, or levels of 8 to 16 ng/ml after stimulation by a milk drink⁴⁸. A preliminary suggestion has been made that levels are higher in patients with pancreatic deficiency⁴⁸ and this illustrates the potential role of gut hormone assays in diagnosis, even in diseases in which hormones are not the prime cause of the pathology.

ENTEROGLUCAGON

Enteroglucagon is a convenient name for the substance found in the mucosa of the gastrointestinal tract which, although very different physiologically from pancreatic glucagon, cross reacts with its antibodies. Other less sonorous names that have been used include 'gut glucagon' and 'glucagon-like immuno-reactivity of gastrointestinal origin' (GLI). Enteroglucagon has not yet been purified in significant quantities so that it usually has to be assayed with a cross reacting pancreatic glucagon radioimmunoassay. This measures total glucagon and the enteroglucagon content is then calculated by subtraction of

pancreatic glucagon, which is measured with a second, non-cross-reacting, radioimmunoassay. Enteroglucagon is found in greatest quantity in the mucosa of the ileum and thus occurs rather lower in the bowel than the classical gut hormones⁴⁰. The cell of origin is of endocrine type and lies in the middle zone of the intestinal mucosa, closely applied to the basement membrane⁵². Extraction studies of enteroglucagon show that, like gastrin, it exists in a high and low molecular weight form⁵³. For this reason it is preferable to express concentrations in molar rather than weight units.

Enteroglucagon has been shown to be most potently released from the bowel by long-chain triglycerides⁵⁴ and glucose²⁶. Using the 100 gram oral glucose tolerance test it was found that enteroglucagon release was increased in patients after the operation of vagotomy and drainage and very much higher if the symptoms of dumping occurred⁵⁵. Enteroglucagon release also showed a very high degree of correlation with the fall in plasma volume that occurs in these circumstances⁵⁶. In the dumping syndrome there is an initial period of rapid intestinal propulsion followed by a period of hypomotility⁵⁷ and this later phase coincides in time with the very high levels of enteroglucagon. A patient with a renal tumour producing enteroglucagon⁵⁸ was found to have gross small and large bowel stasis and also a marked hypertrophy of small intestinal villi⁵⁹. After the tumour had been removed plasma enteroglucagon levels returned to normal and the abnormalities disappeared⁵⁹. From such studies as these the possible physiological role of enteroglucagon can be postulated. Enteroglucagon, which is maximally released when unabsorbed food passes into the lower small intestine, may act to produce a slowing of further food transport and also to enhance long-term growth of the absorptive intestinal mucosa.

GASTRIC INHIBITORY POLYPEPTIDE

The presence of a gastric inhibitory component in 10% pure CCK-PZ was first detected by careful comparison of the biological properties of this impure material with a more highly purified CCK-PZ preparation⁶⁰. When a gastric inhibitory polypeptide (GIP) was completely separated from CCK-PZ-like activity it was found to be a potent inhibitor of even histamine-stimulated gastric acid production²³. The sequence of GIP was published in 1971⁶¹ and seen to have striking similarities with secretin and glucagon. This led to the search for, and discovery of, insulin releasing activity²⁷. Radioimmunoassay of human plasma GIP concentrations showed a considerable rise following ordinary meals⁶², of which the glucose and fat components appeared to be the main stimulatory agents⁶³. The plasma levels achieved were of the same order of magnitude as might have been expected to occur following the exogenous administration of enough GIP to inhibit gastric acid and stimulate insulin release. Clearly further work is required to establish the exact physiological role of GIP, but there is the exciting prospect of the discovery of an important enterogastrone and possible major component of the entero-insular axis.

MOTILIN

When the duodenum of the dog is perfused with an alkaline fluid coordinated gastric contractions result⁶⁴. This effect is seen even in the denervated fundic pouch, and is also mimicked by injection of duodenal extracts⁶⁵. It seems likely, therefore, that a circulating hormone is responsible and the name

'motilin' has been proposed. In 1972 motilin was completely purified and was found to have 22 amino acids⁶⁶ and an amino acid sequence quite dissimilar from other gastrointestinal hormones⁶⁷. Preliminary results of radioimmunoassay of plasma motilin concentrations show a distinctly raised level in dogs after intraduodenal instillation of alkali⁶⁸. More sensitive assays will be required, however, before the physiological importance of this new hormonal peptide can be fully assessed.

VASOACTIVE INTESTINAL PEPTIDE

In 1970 a vasodilator activity was discovered in extracts of the upper small intestine of the pig⁶⁹ and subsequently purified⁷. The amino acid sequence of this vasoactive intestinal peptide (VIP) had considerable similarities with GIP, secretin, and glucagon¹⁹. It was also found to have secretin-like actions in stimulating alkaline pancreatic juice production, and glucagon-like actions in raising blood glucose⁷⁰. Further, VIP produced a powerful stimulation of small intestinal secretion and could inhibit histamine-stimulated gastric acid production²². Radioimmunoassay of VIP has shown it to be distributed throughout the primate gut, with the total quantity present being greater than that of gastrin, secretin, or CCK-PZ⁷¹. Human VIP and porcine VIP elute from Sephadex gel columns in an identical position, suggesting that they may differ little in their amino acid sequences⁷¹.

The physiology of VIP is still unknown but some information can be obtained by studying the pancreatic cholera syndrome. In 1958 Verner and Morrison described two patients who died from extremely profuse watery diarrhoea associated with a pancreatic neoplasm⁷². This syndrome, also called WDHA, because of the watery diarrhoea, hypokalaemia, and achlorhydria⁷³, was found to be associated with high plasma VIP levels⁷⁴. Extracts of the responsible tumours contained large quantities of VIP⁷¹, which behaved as a single substance on gel columns and had the same elution position as VIP from normal human bowel⁷¹. The finding that the main effect of chronically elevated plasma VIP levels may be to cause watery diarrhoea alters the emphasis on the known actions of this hormonal peptide. Although first purified as a general vasodilatory substance it is probable that the gastrointestinal effects of VIP are at least equally important physiologically. Vasoactive intestinal peptide has a wide distribution in the bowel⁷¹ and it may be that it is not primarily concerned with digestion but, perhaps, acts to protect against injurious substances by increasing intestinal secretions and alimentary blood flow.

Conclusions

The gut appears histologically to be a large and complex endocrine organ. Elucidation of the nature and actions of this endocrine system has proved difficult because the cells are widely scattered and the hormones labile. In recent years the purification and sequencing has been reported on six probable hormones and active investigation is proceeding on several others. When a hormone has been isolated it can be measured by the technique of radioimmunoassay which allows very rapid collection of further data. Unfortunately it is not only quite easy to set up a radioimmunoassay but also very easy to get misleading results. It is important to ensure that all the structures that are being measured by the assay reflect biological activity.

Intestinal hormones probably play a significant role in gut and metabolic diseases. Now that their plasma measurement is feasible, an increase in the current research in this field should add considerably to our understanding of the pathology. Determination of the 'gut hormone profile' will also be an important diagnostic tool in investigating diseases not primarily of endocrine origin. It is predictable that in future most hospitals will need access to a comprehensive and reliable gut hormone radioimmunoassay service.

S. R. BLOOM

*The Institute of Clinical Research,
The Middlesex Hospital Medical School,
London*

References

- ¹Pearse, A. G. E. (1973). The gut as an endocrine organ. In *Proceedings of the Symposium on Advanced Medicine*, edited by G. Walker, pp. 400-409. Pitman, London.
- ²Jorpes, E., and Mutt, V. (1961). On the biological activity and amino acid composition of secretin. *Acta chem. scand.*, **15**, 1790-1791.
- ³Gregory, R. A., and Tracy, H. J. (1964). The constitution and properties of two gastrins extracted from hog antral mucosa I. The isolation of two gastrins from hog antral mucosa. II. The properties of two gastrins isolated from hog antral mucosa. *Gut*, **5**, 103-114.
- ⁴Jorpes, E., Mutt, V., and Toczko, K. (1964). Further purification of cholecystokinin and pancreozymin. *Acta chem. scand.*, **18**, 2408-2410.
- ⁵Brown, J. C., Mutt, V., and Dryburgh, J. R. (1971). The further purification of motilin, a gastric motor activity stimulating polypeptide from the mucosa of the small intestine of hogs. *Canad. J. Physiol.*, **49**, 399-405.
- ⁶Brown, J. C., Mutt, V., and Pederson, R. A. (1970). Further purification of a polypeptide demonstrating enterogastrone activity. *J. Physiol. (Lond.)*, **209**, 57-64.
- ⁷Said, S. I., and Mutt, V. (1970). Polypeptide with broad biological activity: Isolation from small intestine. *Science (Wash.)*, **169**, 1217-1218.
- ⁸Berson, S. A., and Yalow, R. S. (1971). Nature of immunoreactive gastrin extracted from tissues of gastrointestinal tract. *Gastroenterology*, **60**, 215-222.
- ⁹Yalow, R. S., and Berson, S. A. (1970). Size and charge distinctions between endogenous human plasma gastrin in peripheral blood and heptadecapeptide gastrins. *Gastroenterology*, **58**, 609-615.
- ¹⁰Gregory, H., Hardy, P. M., Jones, D. S., Kenner, G. W., and Sheppard, R. C. (1964). The antral hormone gastrin: structure of gastrin. *Nature (Lond.)*, **204**, 931-933.
- ¹¹Anderson, J. C., Barton, M. A., Gregory, R. A., Hardy, P. M., Kenner, G. W., Macleod, J. K., Preston, J., Sheppard, R. C., and Morley, J. S. (1964). The antral hormone gastrin II. Synthesis of gastrin. *Nature (Lond.)*, **204**, 933-934.
- ¹²Gregory, R. A. Personal communication.
- ¹³Gregory, R. A., and Tracy, H. J. (1972) Isolation of two 'big gastrins' from Zollinger-Ellison tumour tissue. *Lancet*, **2**, 797-799.
- ¹⁴Yalow, R. S., and Berson, S. A. (1971). Further studies on the nature of immunoreactive gastrin in human plasma. *Gastroenterology*, **60**, 203-214.
- ¹⁵Yalow, R. S., and Berson, S. A. (1972). And now 'big, big' gastrin. *Biochem. biophys. Res. Commun.*, **48**, 391-395.
- ¹⁶Rehfeld, J. F. (1972). Three components of gastrin in human serum: gel filtration studies on the molecular size of immunoreactive serum gastrin. *Biochim. biophys. Acta (Amst.)*, **285**, 364-372.
- ¹⁷Walsh, J. H., and Grossman, M. I. (1973). Circulating gastrin in peptic ulcer disease. *Mnt. Sinai J. Med.*, **40**, 374-381.
- ¹⁸Stern, D. H., and Walsh, J. H. (1973). Gastrin release in post-operative ulcer patients: Evidence for release of duodenal gastrin. *Gastroenterology*, **64**, 363-369.
- ¹⁹Bodanszky, M., Klausner, Y. S., and Said, S. I. (1973). Biological activities of synthetic peptides corresponding to fragments of and to the entire sequence of the vasoactive intestinal peptide. *Proc. nat. Acad. Sci. (Wash.)*, **70**, 382-384.
- ²⁰Chey, W. Y., Hitanant, S., Hendricks, J., and Lorber, S. H. (1970). Effect of secretin and cholecystokinin on gastric emptying and gastric secretion in man. *Gastroenterology*, **58**, 820-827.
- ²¹Lin, T. M., and Spray, G. F. (1968). Effect of glucagon on gastrin HCl secretion. *Gastroenterology*, **54**, 1254.
- ²²Barbezat, G. O., and Grossman, M. I. (1971). Intestinal secretion: stimulation by peptides. *Science*, **174**, 422-423.
- ²³Pederson, R. A., and Brown, J. C. (1972). Inhibition of histamine, pentagastrin and insulin stimulated canine gastric secretion by pure 'gastric inhibitory polypeptide'. *Gastroenterology*, **62**, 393-400.
- ²⁴Samols, E., Marri, G., and Marks, V. (1965). Promotion of insulin secretion by glucagon. *Lancet*, **2**, 415-416.
- ²⁵Dupré, J. (1964). An intestinal hormone affecting glucose disposal in man. *Lancet*, **2**, 672-673.
- ²⁶Unger, R. H., Ohneda, A., Valverde, I., Eisentraut, A. M., and Exton, J. (1968). Characterisation of the responses of circulating glucagon-like immunoreactivity to intraduodenal and intravenous administration of glucose. *J. clin. Invest.*, **47**, 48-65.
- ²⁷Dupré, J., Ross, S. A., Watson, D., and Brown, J. C. (1973). Stimulation of insulin secretion by gastric inhibitory polypeptide in man. *J. clin. Endocr.*, **37**, 826-828.
- ²⁸Unger, R. H., Eisentraut, A. M., Sims, K., McCall, M. S., and Madison, L. L. (1961). Sites of origin of glucagon in dogs and humans. *Clin. Res.*, **9**, 53.
- ²⁹Bodansky, M., Ondetti, M. A., Levine, S. D., et al. (1966). Synthesis of a heptacosapeptide amide with the hormonal activity of secretin. *Chem. and Ind.*, **42**, 1757-1758.

- ³⁰Bloom, S. R., and Ogawa, O. (1973). Radioimmunoassay of human peripheral plasma secretin. *J. Endocr.*, **58**, 24-25.
- ³¹Boden, G., and Chey, W. Y. (1973). Preparation and specificity of antiserum to synthetic secretin and its use in a radioimmunoassay. *Endocrinology*, **92**, 1617-1624.
- ³²Holohan, K. N., Murphy, R. F., Flanagan, R. W. J., Buchanan, K. D., and Elmore, D. T. (1973). Enzymic iodination of the histidyl residue of secretin: a radioimmunoassay of the hormone. *Biochim. biophys. Acta (Amst.)*, **322**, 178-180.
- ³³Young, J. D., Lazarus, L., Chisholm, D. J., and Atkinson, F. F. V. (1968). Radioimmunoassay of secretin in human serum. *J. nucl. Med.*, **9**, 641-642.
- ³⁴Sum, P. T., and Preshaw, R. M. (1967). Intraduodenal glucose infusion and pancreatic secretion in man. *Lancet*, **2**, 340-341.
- ³⁵Hong, S. S., Chin, D. S., and Hur, K. B. (1961). Influences of hexamethonium and some dietary factors on human pancreatic and bile secretion. *J. appl. Physiol.*, **16**, 810-814.
- ³⁶Wang, C. C., and Grossman, M. I. (1951). Physiological determination of the release of secretin and pancreozymin from intestine of dogs with transplanted pancreas. *Amer. J. Physiol.*, **164**, 527-545.
- ³⁷Boehm, M., Oliai, A., and Chey, W. Y. (1973). The production of specific anti-secretin sera and its use in the radioimmunoassay of secretin (Abstr.) *Gastroenterology*, **64**, 703.
- ³⁸Bloom, S. R., Unpublished observations.
- ³⁹Buchanan, K. D. Personal communication.
- ⁴⁰Bloom, S. R., and Bryant, M. G. (1974). Distribution of radioimmunoassayable gastrin, secretin, pancreozymin and enteroglucagon in rat, dog and baboon gut. *J. Endocr.*, in press.
- ⁴¹Wormsley, K. G. (1970). Response to duodenal acidification in man. III. Comparison with the effects of secretin and pancreozymin. *Scand. J. Gastroent.*, **5**, 353-360.
- ⁴²Meyer, J. H. Way, L. W., and Grossman, M. I. (1970). Pancreatic response to acidification of various lengths of proximal intestine in the dog. *Amer. J. Physiol.*, **219**, 971-977.
- ⁴³Bloom, S. R., and Ward, A. S. (1974) Secretin release in man after intraduodenal acid (Abstr.) *Gut*, **15**, 338.
- ⁴⁴Young, J. D., Lazarus, L., and Chisholm, D. J. (1969). Radioimmunoassay of pancreozymin cholecystokinin in human serum. *J. nucl. Med.*, **10**, 743-745.
- ⁴⁵Go, V. L. W., Ryan, R. J., and Summerskill, W. H. J. (1971). Radioimmunoassay of porcine cholecystokinin-pancreozymin. *J. Lab. clin. Med.*, **77**, 684-689.
- ⁴⁶Reeder, D. D., Becker, H. D., Smith, N. J., Rayford, P. L., and Thompson, J. C. (1972). Radioimmunoassay of cholecystokinin. *Surg. Forum*, **23**, 261-362.
- ⁴⁷Englert, E., Jr. (1973). Radioimmunoassay (RIA) of cholecystokinin. *Clin. Res.*, **21**, 207.
- ⁴⁸Harvey, R. F., Dowsett, L., Hartog, M., and Read, A. E. (1973). A radioimmunoassay for cholecystokinin-pancreozymin. *Lancet*, **2**, 826-828.
- ⁴⁹Jorpes, J. E., and Mutt, V. (1973). Secretin and cholecystokinin. In *Secretin, Cholecystokinin, Pancreozymin and Gastrin*, edited by J. E. Jorpes and V. Mutt, pp. 1 to 179. Springer, Berlin.
- ⁵⁰Rubin, B., Engel, S. L., Drungis, A. M., Dzelzkalns, M., Grigas, E. O., Waugh, M. H., and Yiacas, E. (1969). Cholecystokinin-like activities in guinea pigs and in dogs of the C-terminal octopeptide (SQ 19,844) of cholecystokinin. *J. pharm. Sci.*, **58**, 955-959.
- ⁵¹Torsali, A., Ramorino, M. L., and Carratu, R. (1973). On the use of cholecystokinin in the roentgenological examination of the extrahepatic biliary tract and intestines. In *Secretin, Cholecystokinin, Pancreozymin and Gastrin*, edited by J. E. Jorpes and V. Mutt, pp. 247-258. Springer, Berlin.
- ⁵²Polak, J. M., Bloom, S. R., Coulling, I., and Pearse, A. G. E. (1971). Immunofluorescent localisation of enteroglucagon cells in the gastrointestinal tract of the dog. *Gut*, **12**, 311-318.
- ⁵³Valverde I., Rigopoulou, D., Exton, J., Ohneda, A., Eisentraut, A., and Unger, R. H. (1968). Demonstration and characterization of a second fraction of glucagon-like immunoreactivity in jejunal extracts. *Amer. J. med. Sci.*, **255**, 415-420.
- ⁵⁴Botting, I., Faloon, G., and Unger, R. H. (1972). Response of islet cell and gut hormones to fat absorption: an 'enteroinsular axis' for fat. *Clin. Res.*, **20**, 542.
- ⁵⁵Bloom, S. R., Royston, C. M. S., and Thomson, J. P. S. (1972). Enteroglucagon release in the dumping syndrome. *Lancet*, **2**, 789-791.
- ⁵⁶Thomson, J. P. S., Bloom, S. R., Haynes, S., and Ogawa, O. (1973). Plasma enteroglucagon and plasma volume changes after oral hypertonic glucose: their relationship to the dumping syndrome. (Abstr.) *Brit. J. Surg.*, **60**, 308.
- ⁵⁷Christofferson, E., Kewenter, J., and Kock, N. G. (1962). Intestinal motility during provoked dumping reaction. *Acta chir. scand.*, **123**, 405-414.
- ⁵⁸Bloom, S. R. (1972). An enteroglucagon tumour. *Gut*, **13**, 520-523.
- ⁵⁹Gleeson, M. H., Bloom, S. R., Polak, J. M., Henry, K., and Dowling, R. H. (1971). Endocrine tumour in kidney affecting small bowel structure, motility, and absorptive function. *Gut*, **12**, 773-782.
- ⁶⁰Brown, J. C., and Pederson, R. A. (1970). A multiparameter study on the action of preparations containing cholecystokinin-pancreozymin. *Scand. J. Gastroent.*, **5**, 537-541.
- ⁶¹Brown, J. C., and Dryburgh, J. R. (1971). A gastric inhibitory polypeptide. II. The complete amino acid sequence. *Canad. J. Biochem.*, **49**, 867-872.
- ⁶²Kuzio, M., Dryburgh, J. R., Mallow, K., and Brown, J. C. (1973). Radioimmunoassay for gastric inhibitory polypeptide. *Gastroenterology*, in press.
- ⁶³Brown, J. C. (1973). Gastric inhibitory polypeptide. In *Endocrinology*, 1973, edited by S. Taylor. Heinemann, London. (In press).
- ⁶⁴Brown, J. C., Johnston, L. P., and Magee, D. F. (1966). Effect of duodenal alkalization on gastric motility. *Gastroenterology*, **50**, 333-339.
- ⁶⁵Brown, J. C. (1967). The presence of a gastric motor stimulating property in duodenal extracts. *Gastroenterology*, **52**, 225-229.
- ⁶⁶Brown, J. C., Cook, M. A., and Dryburgh, J. R. (1972). Motilin, a gastric motor activity—stimulating polypeptide: final purification, amino acid composition, and C terminal residues. *Gastroenterology*, **62**, 401-404.
- ⁶⁷Brown, J. C., Cook, M. A., and Dryburgh, J. R. (1973). Motilin, a gastric motor activity stimulating polypeptide: the complete amino acid sequence. *Canad. J. Biochem.*, **51**, 533-537.
- ⁶⁸Brown, J. C., and Dryburgh, J. R. (1973). Personal communication.
- ⁶⁹Said, S. I., and Mutt, V. (1970). Potent peripheral and splanchnic vasodilator peptide from normal gut. *Nature (Lond.)*, **225**, 863-864.

- ⁷⁰Said, S. I., and Mutt, V. (1972). Isolation from porcine-intestinal wall of a vasoactive octacosapeptide related to secretin and to glucagon. *Europ. J. Biochem.*, **28**, 199-204.
- ⁷¹Bloom, S. R., and Bryant, M. G. (1973). The distribution of vasoactive intestinal peptide (VIP) in the primate gastrointestinal tract and characterisation of VIP from human tumours. (Abstr.) *Gut*, **14**, 823.
- ⁷²Verner, J. V., and Morrison, A. B. (1958). Islet cell tumor and syndrome of refractory watery diarrhoea and hypokalemia. *Amer. J. Med.*, **25**, 374-380.
- ⁷³Marks, I. N., Bank, S., and Louw, J. H. (1967). Islet cell tumor of the pancreas with reversible watery diarrhoea and achlorhydria. *Gastroenterology*, **52**, 695-708.
- ⁷⁴Bloom, S. R., Polak, J. M., and Pearse, A. G. E. (1973). Vasoactive intestinal peptide and watery-diarrhoea syndrome. *Lancet*, **2**, 14-16.