

Leading article

Fibre, fermentation, flora, and flatus

Research into flatogenesis has undergone several, apparently unrelated, phases. Early studies of the volume of gas in the gastrointestinal tract^{1,2} were prompted by the abdominal pain and discomfort experienced by sailors and airmen who had been subjected to explosive decompression. The second phase was aided by the realisation that human gas production could pose an explosion hazard during longterm space flight.³ Since that time, there has been little attempt to quantitate gas production in man although there have been many studies on concentrations of H₂ and CH₄ in end expired breath. There have been remarkable advances during the last 15 years in our understanding of the nature of non-absorbed sugars and dietary fibre and of their fermentation in the large intestine. It is now possible to construct a clearer picture of the overall stoichiometry of colonic fermentation and gas production in man.


The link between large intestine fermentation of non-absorbed carbohydrate and gas production has been exploited in numerous breath tests for malabsorption because a portion of the gas produced (H₂ and CH₄) is absorbed from the lumen of the large intestine into the blood stream and excreted in the lungs. As much as 13% of H₂ gas may be lost through this route.⁴ Whilst H₂ appears to be the major unique gas produced by intestinal fermentation, a proportion of individuals also excrete appreciable quantities of CH₄.^{5–14} In patients with unresected colon cancer the proportion of methane-producers is high (86–92%), falling to normal levels (43–47%) after resection.¹³ Methanogenic bacteria are also present in large quantities in patients with diverticulosis.¹⁵ In contrast with these data, a recent study of South African populations has shown that although there was a wide variation in the proportion of methane producers in different ethnic groups, this did not relate to the relative risk of developing colon cancer.¹⁶ The picture is confusing and the report of Weaver and colleagues,¹⁷ in this issue of *Gut* is therefore timely. In this study, an attempt has been made to determine the stoichiometry of fermentation of defined substrates by incubation of faecal samples from a H₂ and a CH₄ producer. It addresses the essential questions of 'How much', 'Why?', 'Where?', 'What?', and 'Why not?'.

How much?


Most studies have measured only the small proportion of H₂ and CH₄ which is excreted in the breath. Because H₂ and CH₄ are expressed as concentrations, it is not possible to calculate total output unless respiratory volumes are known. End expiratory gas measurements are straightforward in patients or volunteers, but rectal intubation for measurement of flatus volumes is not. A non-invasive alternative to rectal intubation is suggested

by the report of Garlick and colleagues,¹⁸ in which total ¹⁴C output was measured during infusion with ¹⁴C-labelled amino acids. Each subject was housed in a ventilated, disposable oxygen tent and measurements of gas flow and gas composition made with simple equipment. This technique may be useful in quantitating intestinal gas production in man. There is evidence that humans are considerably less flatogenic than theory would suggest and the reason for this may relate to the health of the large intestine.

A reason for believing that man passes less wind than he should is suggested by the elegant study of Weaver and colleagues.¹⁷ In an earlier review of the subject,¹⁹ the stoichiometry of colonic fermentation in man was proposed to be:



Alternatively, if no CH₄ were produced by CO₂ reduction, the equation could be rewritten:



In Weaver's¹⁷ study faecal suspensions from both subjects fermented glucose and corn starch to produce VFA's and gas, although the relative proportions of each product varied. The VFA, CH₄, and H₂ yields per glucose molecule were considerably lower than would be expected by either equation.

A second reason is suggested by considering the amounts of carbohydrate which are fermented in the large intestine. It is thought that an average value should be approximately 70 g, calculated on the basis of the amount required to produce the observed bacterial mass of human faeces.²⁰ In other words, to the quantity of fermentable non-starch polysaccharide normally ingested^{21, 22} should be added the significant quantities of starch which may be present in the diet in a form which resists small intestinal amyolysis and brush border assimilation of constituent glucose moieties.²³⁻²⁵ Fermentation of 70 g of carbohydrate would produce, by either scheme described above, either 6 litres/d of CH₄ or a truly Zeppelinogenic 24 litres/d of H₂. Flatus volume, however is rarely more than 1 litre/day.^{3, 4, 26}

Where, therefore does all of the gas go and what may be the significance of submaximal flatogenesis? The pattern of carbohydrate fermentation by anaerobic bacteria in the large intestine may explain part of this difference.

Why?

Metabolic events in the lumen of the large bowel occur under anaerobic conditions such that complete oxidative microbial metabolism of carbohydrate by intestinal flora cannot occur through the Krebs's Cycle. Limited oxidation of sugars and their intermediary metabolites can only proceed if reduced equivalents are removed by a 'metabolic sink'. H₂ and CH₄ production therefore provides a simple means of removing reduced equivalents, in gaseous form. The energy yield of carbohydrate to the bacteria is rather low and further oxidative metabolism of the bulk of

volatile fatty acids (VFA's) produced has to occur within the colonocyte, or in other tissues after uptake and transmucosal transport. An elegant summary of the metabolic pathways by which organic acids, and in particular the VFA's, are produced by different classes of colonic bacteria, is given elsewhere.^{19 20 27} The major feature of these pathways is that hexose sugars are cleaved to form pyruvate, through the Emden-Meyerhoff Pathway. Two pathways are then available for disposal of pyruvate. First, it may be reductively converted to propionate (through succinate). The second pathway converts pyruvate to acetyl-CoA which may then either be hydrolysed (to release acetate), or reduced to form butyrate. Conversion of pyruvate to acetyl-CoA produces reducing equivalents as H₂, whereas formation of propionate and butyrate consume H₂. If, however, anaerobic degradation of sugars were allowed to proceed to completion, the end products would be CO₂ and CH₄.

It is therefore clear that the extent of methanogenesis will determine the pattern of VFA's formed in the intestine. The major species responsible for methane production in man is *Methanobrevibacter smithii*.^{6 7} CH₄ is formed by this organism from CO₂ and H₂ produced by other bacteria. It seems that in man (as in the ruminant), the time required by acetate using methanogens to oxidise all fatty acids and H₂ to CO₂ and CH₄ is longer than large bowel transit time.²⁰ Production of methane from CO₂ and hydrogen results in decreased gas volumes, because one volume of methane is produced from four volumes of H₂. This may explain in part, submaximal flatogenesis in man.

Different bacterial species are responsible for H₂ and CH₄ generation and there are variations in the dynamics of their production in man. For the readily fermentable carbohydrate, lactose, one half of H₂ production occurs in the caecum and right colon, compared to less than 10% of methane production.⁴ Ingestion of lactulose and breath H₂ also correlate well²⁸ and the evidence suggests that malabsorption of a test carbohydrate should be 'calibrated' by normalising the breath H₂ response to that of a known amount of lactulose.²⁹ In methane producers, however, the breath CH₄ response to oral lactulose is idiosyncratic¹² and may relate to the rapid fermentation of this substrate in the caecum and right colon such that little is available for further fermentation at more distal, methanogenic sites. Despite the usefulness of breath H₂ tests, there are several caveats. First, preexisting treatment (antibiotics, enema or colonoscopy) markedly affects H₂ production.^{30 31} More subtly, mood dramatically alters breath H₂ output, which is increased during periods of acute^{3 32} or chronic³ tension or when gas measurements are made in public places.³² This may be caused either by a sudden increase in the amount of fermentable substrate reaching the caecum and right colon (during periods of stress), or to retention of gas (and an increase in its excretion in breath). It is therefore clear that care should be exercised in designing diagnostic H₂ breath tests and that attention should be paid to the comfort and privacy of the subjects.

Interestingly, methane production is unresponsive to mood,³² or pre-existing treatment,³¹ again suggesting that H₂ and CH₄ production are partially separated by anatomy and microbiology in the human large intestine and may be independently altered by transit phenomena. The nature of the non-absorbed carbohydrate may therefore be crucial to the quantities of gas produced.

Where?

The flatogenic potential of all non-absorbed carbohydrates should be the same, if all saccharide bonds are available for hydrolysis and constituent monosaccharides liberated for subsequent bacterial fermentation. This is clearly not the case, however, as shown by the variation in fermentation of non-starch polysaccharides in man³³ and in the pig.³⁴ Variability seems to be determined not only by the type of saccharide linkage involved, but also by the physicochemical characteristics of the non-absorbed polysaccharide – such as the ‘resistant starches’.^{35–37} For non-starch polysaccharides, availability relates more to physical, rather than chemical factors.³⁸ The access of bacterial cell wall polysaccharidases²⁷ to bonds buried deep within highly ordered, relatively dehydrated polysaccharide structures may be limited. Decreased fibre particle size,^{38–40} increased hydration (due to removal of lignin during digestion⁴¹), increased fibre porosity and solubility and decreased viscosity⁴² all tend therefore, to increase the fermentability of non-absorbed carbohydrates. Fermentation of soluble, non-absorbed sugars (Lactulose, Lactitol, Palmitit, Isomalt) in the caecum and right colon is limited by none of these factors and they all express their full fermentative and flatogenic potential when administered orally.^{43–46}

The possibility therefore exists that hydrolysis and fermentation of different non-absorbed polysaccharides may occur at different sites in the large intestine. Levitt and Ingelfinger⁴ observed that methane production occurred primarily in the left colon. Dietary fibre which is poorly hydrolysed and fermented in the right colon may provide a good supply for H₂ and CH₄ production in the left colon. At present this concept is speculative but would be supported by the observation that not only is CH₄ production by methanogenic bacteria pH dependent⁴⁷ but that these strains have increased growth rates at slightly alkaline pH.⁴⁸ Thus their growth would be most favoured by conditions in the left colon. The decrease in large bowel transit time produced by a high fibre diet may, however, lower bacterial contact time and CH₄ production. This may be inferred from the observation that large bowel transit time and faecal pH correlate significantly with breath methane levels⁴⁹ although in the recent study of Segal *et al.*¹⁶ this was not the apparent cause of high levels of methane production in rural black South Africans who had rapid transit consequent on high fibre consumption.

What?

The proportions of gases produced during colonic fermentation may provide a clue to the pattern of VFA production. Weaver and colleagues¹⁷ in this issue of *Gut* provide some data on this point. Volatile fatty acids and gas production were measured, using glucose or starch as the substrate. In the hydrogen producer, glucose and starch disappearance were similar, indicating that hydrolysis of starch was not rate limiting for fermentation. In contrast, starch appeared to be less well fermented than glucose, in the methane producer. The pattern of VFA's also differed. Methanogenesis was accompanied by less propionate and more acetate production. Similar observations have been made in ruminants when ciliate protozoa were established in the rumen: a significant negative correlation between propionate and CH₄ production, was reported.⁵⁰ These changes may have an

impact on the health of the large bowel. It has been suggested⁵¹ that VFA's may have a protective role in relation to colon cancer. Butyrate is a major substrate for colonocyte energy metabolism⁵² and has cytostatic effects on colorectal cell lines.⁵¹ It has recently been shown that VFA's stimulation of crypt cell production rate in the small and large intestine of rats occurs *in vivo* (butyrate > propionate > acetate), but this may be mediated systemically, as all VFA's inhibit epithelial cell proliferation *in vitro*.⁵³ Factors which reduce VFA production may therefore be of importance. Methane production represents a 'sink' into which bacteria may dump reducing equivalent, thus releasing the coenzymes NAD, NADP, and FAD for further bacterial oxidative metabolism of VFA's. In ruminants and to a lesser extent in pigs, methanogenesis represents a loss of VFA's to the host, which would otherwise be absorbed for further metabolism.^{50,54}

The possibility therefore exists that excess methane production may be associated with two, possibly deleterious, phenomena. First, it may tend to decrease the supply of readily metabolised substrate (VFA's) to the colon. It may also alter the production of specific VFA's with growth regulatory properties within the large bowel lumen. The experimental evidence for trophic effects of VFA's on small and large bowel function during enteral nutrition are reviewed elsewhere.⁵⁵ This is clearly an important area and warrants further study.

Why not?

The relationship between H₂ and CH₄ production, the type of fermentable non-absorbed polysaccharide and sites of fermentation is not clearly known. Perfusion and representative sampling of these sites may not be possible without disturbing the balance of microflora within these regions. Attention has therefore been given to incubation systems which mimic the anatomy of the large bowel by use of sequential fermentation chambers.^{56,57} This strategy has been used to study rumen fermentation and the digestion of dietary proteins in the stomach and small intestine.⁵⁸ Site and time related aspects of carbohydrate assimilation may thus be studied in a continuous culture system which is not dissimilar to the human colon. Using this approach, the impact of sulphate reducing bacteria (SRB) on gas and VFA production was determined.⁵⁷ The three chambers were pH controlled to simulate the pH gradient between the caecum and left colon and after initial inoculation with a faecal slurry the system was allowed to reach equilibrium. During infusion of porcine gastric mucin (containing sulphated mucopolysaccharides) CH₄ and H₂ production were completely suppressed in the two distal chambers operated at higher pH. This was accompanied by sulphate reduction (by H₂) and generation of sulphide, catalysed by increasing numbers of SRB. After the mucin infusion ceased, the system returned to its steady-state. In separate experiments with incubated faecal slurries from methane producers and non-methane producers, significant quantities of CH₄ were only produced in the absence of sulphate reduction. Mixtures of slurries from both sources showed the ability of SRB to outcompete methanogenic bacteria (MB) for H₂.⁵⁹ Addition of nitrate,⁵⁹ which can be reduced to nitrite by other classes of faecal anaerobes⁶⁰ variably suppressed CH₄ and H₂S production in both types of slurries. The suggestion that establishment of either MB or SRB colonies (but not both)

occurs in the human colon is supported by the report that SRB in faecal samples from British and South African populations correlated inversely with breath CH₄ production.⁶¹ The supply of sulphate containing mucin to the colon may be the major determinant of SRB colonisation.⁵⁹ Whatever the reason for establishment of SRB in the human colon, it will result in diversion of H₂, away from CH₄ (and thus flatogenesis), and towards non-volatile S²⁻ production.

Conclusion

Research into fibre, fermentation, flora, and flatus has reached an exciting point. There is now a clearer understanding of the stoichiometry of large intestinal fermentation, the ways in which the colonic microflora utilise different types of non-absorbed polysaccharides and also the effects of VFA's on growth of normal and transformed colocytes. What is particularly interesting is the extent to which man is an underperformer in the flatogenic stakes. Secondary utilisation of H₂ by MB, SRB, and nitrate reducing bacteria may account for all of this difference, but quantitative data are lacking. The author would suggest that interest in whole body flatometry be revived. It would not be unreasonable to determine the overall stoichiometry of fermentation and H₂ disposal from the extent to which H₂ and CH₄ production fall short of theoretical yields for a particular dietary fibre, whose digestibility can be defined with the methods of analysis which are now available.

The observation that methanogenic or sulphate reducing bacteria are established in the colon, to the exclusion of the other, is particularly fascinating. Does this represent an underlying difference in the intrinsic sulphation pattern of epithelial mucopolysaccharides which may ultimately be fermented in the large intestine? It is hard not to consider this a good kite to fly in future research.

GEORGE GRIMBLE

*Department of Gastroenterology & Nutrition,
Central Middlesex Hospital,
Acton Lane,
London NW10 7NS*

References

- 1 Blair HA, Dern RJ, Bates PL. The measurement of volume of gas in the digestive tract. *Am J Physiol* 1947; **149**: 688-707.
- 2 Bedell GN, Marshall R, DuBois AB, Harris JH. Measurement of the volume of gas in the gastrointestinal tract. Values in normal subjects and ambulatory patients. *J Clin Invest* 1956; **35**: 336-45.
- 3 Calloway DH, Murphy EL. Intestinal hydrogen and methane of men fed space diet. *Life sciences and space research* 1969; **VII**: 102-9.
- 4 Levitt MD, Ingelfinger FJ. Hydrogen and methane production in man. *Ann NY Acad Sci* 1968; **150**: 75-81.
- 5 Bond JH, Engel RR, Levitt MD. Factors influencing pulmonary methane excretion in man. An indirect method of studying the in situ metabolism of the methane-producing colonic bacteria. *J Exp Med* 1971; **133**: 572-88.
- 6 Miller TL, Wolin MJ, Macario EC de, Macario AJL. Isolation of *Methanobrevibacter smithii* from human feces. *Appl Env Microbiol* 1982; **43**: 227-32.
- 7 Miller TL, Wolin MJ. Enumeration of *Methanobrevibacter smithii* in human species. *Arch Microbiol* 1982; **131**: 14-8.

- 8 Miller TL, Wolin MJ. Stability of *Methanobrevibacter smithii* populations in the microbial flora excreted from the human large bowel. *Appl Env Microbiol* 1983; **45**: 317–8.
- 9 McKay LF, Brydon WG, Eastwood MA, Smith JH. The influence of pentose on breath methane. *Am J Clin Nutr* 1981; **34**: 2728–33.
- 10 Bjorneklepp A, Jenssen E. Relationships between hydrogen (H₂) and methane (CH₄) production in man. *Scand J Gastroenterol* 1982; **17**: 985–92.
- 11 Haines A, Metz G, Dilwari J, Blendis L, Wiggins H. Breath-methane in patients with cancer of the large bowel. *Lancet* 1977; **ii**: 481–3.
- 12 Pitt P, De Bruijn KM, Beeching MF, Goldberg E, Blendis LM. Studies on breath methane: the effect of ethnic origins and lactulose. *Gut* 1980; **21**: 951–9.
- 13 Pique JM, Pallares M, Cuso E, Vilar-Bonet J, Gassull MA. Methane production and colon cancer. *Gastroenterology* 1984; **87**: 601–5.
- 14 McKay LF, Eastwood MA, Brydon WG. Methane excretion in man – a study of breath, flatus and faeces. *Gut* 1985; **26**: 69–74.
- 15 Weaver GA, Krause JA, Miller TL, Wolin MJ. Incidence of methanogenic bacteria in a sigmoidoscopy population: an association of methanogenic bacteria and diverticulosis. *Gut* 1986; **27**: 698–704.
- 16 Segal I, Walker ARP, Lord S, Cummings JH. Breath methane and large bowel cancer risk in contrasting African populations. *Gut* 1988; **29**: 608–13.
- 17 Weaver GA, Krause JA, Miller TL, Wolin MJ. Constancy of glucose and starch fermentations by two different human faecal microbial communities. *Gut* 1989; **30**: 19–25.
- 18 Garlick PJ, Clugston GA, Swick RW, Waterlow JC. Diurnal patterns of protein and energy metabolism in man. *Am J Clin Nutr* 1980; **33**: 1983–6.
- 19 Miller TL, Wolin MJ. Fermentations by saccharolytic intestinal bacteria. *Am J Clin Nutr* 1979; **32**: 164–72.
- 20 Smith CJ, Bryant MP. Introduction to metabolic activities of intestinal bacteria. *Am J Clin Nutr* 1979; **32**: 149–57.
- 21 Cummings JH. Short chain fatty acids in the human colon. *Gut* 1981; **22**: 763–79.
- 22 McNeill NI. The contribution of the large intestine to energy supplies in man. *Am J Clin Nutr* 1984; **39**: 338–42.
- 23 Southgate D, Englyst H. Dietary fibre: chemistry, physical properties and analysis. In: Trowell H, Burkitt D, Heaton K, eds. *Dietary fibre, fibre-depleted foods and disease*. London: Academic Press, 1985: 31–55.
- 24 Englyst HN, Macfarlane GT. Breakdown of resistant and readily digestible starch by human gut bacteria. *J Sci Food Agric* 1986; **37**: 699–706.
- 25 Englyst HN, Macfarlane GT, Cummings JH. New concepts in starch digestion in man [Abstract]. *Proc Nutr Soc* 1988; **47**: 64A.
- 26 Calloway DH, Burroughs SE. Effect of dried beans and silicone on intestinal hydrogen and methane production in man. *Gut* 1969; **10**: 180–4.
- 27 Salyers AA, Leedle JAZ. Carbohydrate metabolism in the human colon. In: Hentges DJ, ed. *Human intestinal microflora in health and disease*. New York, London: Academic Press, 1983: 129–46.
- 28 LaBrooy SJ, Male P-J, Beavis AK, Misiewicz JJ. Assessment of the reproducibility of the lactulose breath test as a measure of mouth to caecum transit time. *Gut* 1983; **24**: 893–6.
- 29 Bond JH, Levitt MD. Fate of soluble carbohydrate in the colon of rats and man. *J Clin Invest* 1975; **57**: 1158–64.
- 30 Gilat T, Ben-Hur H, Gelman-Malaghi E, Terdiman R, Peled Y. Alterations of the colonic flora and their effect on the hydrogen breath test. *Gut* 1978; **19**: 602–5.
- 31 Murphy EL, Calloway DH. The effect of antibiotic drugs on the volume and composition of intestinal gas from beans. *Am J Dig Dis* 1972; **17**: 639–42.
- 32 Calloway DH, Murphy EL. The use of expired air to measure intestinal gas formation. *Ann NY Acad Sci* 1968; **150**: 82–95.
- 33 Nyman M, Asp N-G, Cummings J, Wiggins H. Fermentation of dietary fibre in the intestinal tract: comparison between man and rat. *Br J Nutr* 1986; **55**: 487–96.
- 34 Stanogias G, Pearce GR. The digestion of fibre by pigs. 1. The effects of amount and type of fibre on apparent digestibility, nitrogen balance and rate of passage. *Br J Nutr* 1985; **53**: 513–30.
- 35 Englyst HN, Cummings JH. Digestion of the polysaccharides of some cereal foods in the human small intestine. *Am J Clin Nutr* 1985; **42**: 778–87.
- 36 Englyst HN, Cummings GT. Breakdown of resistant and readily digestible starch by human gut bacteria. *J Sci Food Agric* 1986; **37**: 699–706.
- 37 Englyst HN, Cummings JH. Digestion of polysaccharides of potato in the small intestine of man. *Am J Clin Nutr* 1987; **45**: 423–31.

- 38 Robertson JA. Physicochemical characteristics of food and the digestion of starch and dietary fibre during gut transit. *Proc Nutr Soc* 1988; **47**: 143–52.
- 39 Ehle FR, Murphy MR, Clark JH. In situ particle size reduction and the effect of particle size on degradation of crude protein and dry matter in the rumen of dairy steers. *J Dairy Sci* 1982; **65**: 963–71.
- 40 Heller SN, Hackler LR, Rivers JM, *et al.* Dietary fiber: the effect of particle size of wheat bran on colonic function in young adult men. *Am J Clin Nutr* 1980; **33**: 1734–44.
- 41 Millard P, Chesson A. Modifications to swede (*Brassica napus* L.) anterior to the terminal ileum of pigs: some implications for analysis of dietary fibre. *Br J Nutr* 1984; **52**: 583–94.
- 42 Tomlin J, Read NW, Edwards CA, Duerden BI. The degradation of guar gum by a faecal incubation system. *Br J Nutr* 1986; **55**: 481–6.
- 43 Patil DH, Grimble GK, Silk DBA. Lactitol, a new hydrogenated lactose derivative: intestinal absorption and laxative threshold in normal human subjects. *Br J Nutr* 1987; **57**: 195–9.
- 44 Grimble GK, Patil DH, Silk DBA. Assimilation of lactitol, an ‘unabsorbed’ disaccharide in the normal human colon. *Gut* 1988; **29**: 1666–71.
- 45 Grenby TH, Desai T. The dental effects of lactitol replacing sucrose in confectionery examined in a clinical trial [Abstract]. *Proc Nutr Soc* 1988; **47**: 2A.
- 46 Wursch P, Schweizer T. Breath hydrogen after ingestion of three disaccharide alcohols and lactulose. *Proceedings of Zuckersymposium ‘Karies und Zucker’*. 12–13th June 1986, Wurzburg, FRG.
- 47 Bryant MP. Microbial methane production – theoretical aspects. *J Anim Sci* 1979; **48**: 193–201.
- 48 Schauer NL, Ferry JG. Metabolism of formate in *Methanobacter formicum*. *J Bacteriol* 1980; **142**: 800–7.
- 49 Stephen AM, Wiggins HS, Englyst HN, Cole TJ, Wayman BJ, Cummings JH. The effect of age, sex and level of intake of dietary fibre from wheat on large-bowel function in thirty healthy subjects. *Br J Nutr* 1986; **56**: 349–61.
- 50 Whitelaw FG, Eadie JM, Bruce LA, Shand WJ. Methane formation in faunated and ciliate-free cattle and its relationship with rumen volatile fatty acid proportions. *Br J Nutr* 1984; **52**: 261–75.
- 51 Cummings J. Cancer of the large bowel. In: Trowell H, Burkitt D, Heaton K, eds. *Dietary fibre, fibre-depleted foods and disease*. London: Academic Press, 1985: 161–89.
- 52 Roediger WEW. Role of anaerobic bacteria in the metabolic welfare of the colonic mucosa in man. *Gut* 1980; **21**: 793–8.
- 53 Sakata T. Stimulatory effect of short-chain fatty acids on epithelial cell proliferation in the rat intestine: a possible explanation for trophic effects of fermentable fibre, gut microbes and luminal trophic factors. *Br J Nutr* 1987; **58**: 95–103.
- 54 Christensen K, Thorbeck G. Methane excretion in the growing pig. *Br J Nutr* 1987; **57**: 355–61.
- 55 Silk DBA. Progress report: fibre and enteral nutrition. *Gut* 1989. (In press).
- 56 MacFarlane GT, Englyst HN. Use of multichamber continuous culture system to investigate the degradation of non-starch polysaccharides by human gut bacteria [Abstract]. Proceedings of 4th International Symposium on Microbial Ecology. Slovene Society for Microbiology. Ljubljana. 1986; **D14-5**: 109.
- 57 Gibson GR, Cummings JH, MacFarlane GT. Use of a three-stage continuous culture system to study the effect of mucin on dissimilatory sulphate reduction and methanogenesis by mixed populations of human gut bacteria. *Appl Env Micro* 1988; **54** (In press).
- 58 Chung S-Y, Swaisgood HE, Catigani GL. Effects of alkali treatment and heat treatment in the presence of fructose on digestibility of food proteins as determined by an Immobilised digestive Enzyme Assay (IDEA). *J Agric Food Chem* 1986; **34**: 579–84.
- 59 Gibson GR, Cummings JH, MacFarlane GT. Competition for hydrogen between sulphate reducing bacteria and methanogenic bacteria from the human large intestine. *J Appl Bact* 1988; **65** (In press).
- 60 Allison C, MacFarlane GT. Effect of nitrate upon methane production. *J Gen Microbiol* 1988; **134**: 1397–405.
- 61 Gibson GR, MacFarlane GT, Cummings JH. Occurrence of sulphate-reducing bacteria in human faeces and the relationship of dissimilatory sulphate reduction to methanogenesis in the large gut. *J Appl Bact* 1988; **65**: 103–11.