Regulation of the oncodevelopmental expression of type 1 chain ABH and Lewis^b blood group antigens in human colon by α -2-L-fucosylation

T F Ørntoft, P Greenwell, H Clausen, W M Watkins

Abstract

Blood group antigen expression in the distal human colon is related to the development of the organ and is modified by malignant transformation. To elucidate the biochemical basis for these changes, we have (a) analysed the activity of glycosyltransferases coded for by the H, Se, Le, X, and A genes, in tissue biopsy specimens from normal and malignant proximal and distal human colon; (b) characterised the glycosphingolipids expressed in the various regions of normal and malignant colon by immunostaining of high performance thin layer chromatography plates; and (c) located the antigens on tissue sections from the same subjects by immunohistochemistry. In both secretors and non-secretors we found a significantly higher activity of α -2-L-fucosyltransferases in carcinomatous rectal tissue than in tissue from normal subjects, whereas the other transferase activities studied showed no significant differences. The acceptor substrate specificity suggested that both the Se and the H gene dependent α -2-L-fucosyltransferases are increased in carcinomas. In non-malignant tissue the only enzyme which showed appreciably higher activity in caecum than in rectum was α -2-L-fucosyltransferase. Immunohistochemistry and immunohistochemistry showed α -2-L-fucosylated structures in normal caecum from secretors and in tumour tissue from both secretors and non-secretors. We conclude that the α -2-L-fucosyltransferases control the expression of ABH, and Lewis^b structures in normal and malignant colon.

The antigens that constitute the ABO blood group system are oligosaccharides. Their biosynthesis is regulated by genes coding for specific glycosyltransferases that add monosaccharides to precursors in a sequential fashion (Fig 1). In this way the product of one transferase becomes the substrate of the next, and the loss of activity of one transferase leads to the disappearance of the antigen it would have formed and all subsequent antigens, which would have been synthesised with this structure or its derivatives, as acceptor substrates. This complicated interrelation leads to a differential expression of blood group antigens in people with different genotypes. The role of the ABO, Se, H, X, and Le gene loci and their glycosyltransferase products in determining these antigens has been reviewed in detail elsewhere.¹ These antigens are not only found in blood cells but are expressed in most epithelia and body fluids. In epithelia they have been shown to be differentiation depend-

ent,^{2,3} and several carbohydrate antigens have been suggested as tissue specific oncodevelopmental antigens.⁴

A well examined epithelium is the colorectal mucosa.⁵⁻⁸ Both chemical analysis of carbohydrate structures and immunohistological analysis with monoclonal antibodies have shown that ABH related carbohydrate antigens carried by type 1 chain (Gal β 1-3GlcNAc-R) core structures, as well as those carried by type 2 chain (Gal β 1-4GlcNAc-R) core structures are expressed in the fetal colon and rectum, and that there is a gradual disappearance or appreciable reduction in expression of blood group antigens apart from Lewis^a in the distal colon during fetal growth.^{9,10} In contrast, all type 1, and to a less extent some type 2, chain antigens are expressed in the proximal colon in adults.⁶⁻⁸ In carcinomas of the distal colon, the ABH, Le^b, Le^x, and Le^y antigens⁶⁻⁸ and extended,⁸⁻¹¹ polyfucosylated,^{8,12} or sialylated^{7,13} derivatives of these antigens have been shown by immunochemical and immunohistological means to reappear or to be synthesised de novo in large amounts. This has made them suitable as tumour markers,¹⁴⁻¹⁶ as more than two thirds of all colonic carcinomas are located in the distal colon.

The purpose of the present work was to examine the enzyme and structural basis for the differential expression of blood group carbohydrate antigens in the proximal and distal normal colon and in colorectal carcinomas. The enzyme findings have been correlated with immunochemical characterisation and immunohistochemical localisation of blood group antigens in tissues of the same subjects by the use of monoclonal antibodies with established carbohydrate specificities.

We report that control of the expression of type 1 chain, ABH, and Le^b structures in both normal and malignant colon is probably through α -2-L-fucosyltransferase, whereas type 2 chain structures are, in addition, subject to other regulatory mechanisms.

Methods

REAGENTS

GDP-L-[U-¹⁴C]-fucose (210-292 mCi/mmol), UDP-N-acetyl-D-[1-¹⁴C]-galactosamine (54 mCi/mmol), and UDP-D-[U-¹⁴C]-galactose (300 mCi/mmol) were obtained from Amersham International Ltd UK. N-Acetylglucosamine (Gal β 1-4GlcNAc), lacto-N-biose I (Gal β 1-3GlcNAc), and 2'-fucosyllactose (Fuc α 1-2Gal β 1-4Glc) were kindly supplied by Dr A S R Donald, MRC Clinical Research Centre, Harrow, UK.

Department of
Experimental Clinical
Oncology, Danish Cancer
Society and Department
of Clinical Chemistry,
Aarhus, Denmark
T F Ørntoft

Biomembrane Institute,
Seattle and University of
Washington, USA
H Clausen

Department of
Immunochemical
Genetics, Clinical
Research Centre,
Harrow, Middlesex, UK
P Greenwell
W M Watkins

Correspondence to:
Dr T F Ørntoft, Department
of Experimental Clinical
Oncology, Danish Cancer
Society, Nørrebrogade 44,
DK-8000 Aarhus C,
Denmark.

Accepted for publication
29 May 1990

Figure 1: Biosynthetic pathways for blood group ABO related antigen determinants carried by type 1 (Gal β 1-3GlcNAc-R) and type 2 (Gal β 1-4GlcNAc-R) carbohydrate chains. The gene responsible for each biosynthetic step is underlined, and the pathway that seems to control the expression of structures in normal and malignant colon is shown by two lines. Fuc=fucose; Gal=galactose; GalNAc=N-acetylgalactosamine; GlcNAc=N-acetylglucosamine.

Phenyl- β -D-galactoside was obtained from Koch-Light, Colnbrook; Triton X-100 was purchased from BDH Chemicals, UK. Diethylaminoethyl cellulose (DE 81), and No 40 chromatography papers were obtained from Whatman, UK. Mouse monoclonal antibodies against blood group H and biotinylated swine antirabbit immunoglobulins were purchased from Dakopatts, Copenhagen, Denmark. Avidin-Biotin-Peroxidase complex was obtained from Vector Lab, Burlingame, California, USA. Mouse monoclonal antibodies against Le^a and Le^b antigens were kindly supplied by Dr L Messeter, the Bloodbank, Lund, Sweden. The specificity of anti-Le^a was tested against Le^a, Le^b, Le^c, and Le^d glycolipids, that of anti-Le^b against Le^b, Le^d, H type 2 chain, Le^a, Le^c, and ALe^b glycolipids (L Messeter, personal communication). Mouse monoclonal antibodies against Le^a, clone AH₆, IgM¹⁷; Le^a, clone FH₃,¹² IgG₃; A type 1 chain, clones AH21 (monofucosylated), IgM¹⁸; and HH₃ (difucosylated), IgG_{2A}¹⁹; A type 2 chain, clones HH₂ (mono- and difucosylated), IgG₃²⁰; and HH₄ (difucosylated), IgG₃,²¹ were from the Biomembrane Institute, Seattle, Washington. 3-Amino-9-ethylcarbazole and Tris were purchased from Sigma Chemical Co, St Louis, USA.

SPECIMENS

Biopsy specimens of normal tissue were

obtained from the rectum, sigmoid, left flexure, right flexure, and caecum from four A₁ Le(a-b+), two O Le(a+b-), and one A₁ Le(a+b-) subjects. Tumour tissue was obtained from the rectum of 15 subjects. The tumours were graded as well differentiated two, moderate 10, and poor three; two were Dukes's A, 10 were Dukes's B, and three were Dukes's C. Tumour donors were blood grouped as five A₁ Le(a-b+); one A₁ Le(a-b-); one A₂ Le(a+b-); five O Le(a-b+); and three O Le(a+b-). Le(a-b+) subjects were regarded as ABH secretors, Le(a+b-) subjects as ABH non-secretors. In the Le(a-b-) subject, saliva was used for determination of secretor state by haemagglutination inhibition.

GLYCOSYLTRANSFERASE ASSAYS

A modification of previously described^{22,23} assays was used. One gram of sample was solubilised in 1 ml 0.15 M NaCl, 2% Triton X-100 by sonication (Soniprobe, Dawes Instruments Ltd, UK) at 4°C, the resulting suspension centrifuged at 10 000 rpm for 30 min (Sorvall RC2-B. Rotor SS-34), and the supernatant used for the assays. For the assays of α -2-, α -3-, and α -4-L-fucosyltransferases, 20 μ l of the enzyme source was added to GDP-L-[¹⁴C]-fucose (0.56 nmol, 120 000 cpm), MnCl₂ (1 μ mol), acceptor (0.5 μ mol), adenosine triphosphate (0.5 μ mol), Triton X-100 500 μ g, and Tris-HCl, pH 7.2 (5 μ mol) in a total volume

of 95 μ l. The mixture was incubated at 37°C for 4 hours. When phenyl- β -D-galactoside was used as the acceptor substrate the mixture was spotted on Whatman No 40 paper and developed with ethylacetate/pyridine/water, 10:4:3 (by vol, solvent 1) for 4 hours. When N-acetyllactosamine, lacto-N-biose I or 2'-fucosyllactose were used as acceptors the incubation mixtures were chromatographed on Whatman No 40 paper in solvent 1 for 48 hours. The papers were dried and scanned for radioactivity in a Packard radiochromatogram scanner. The mobilities of the radioactive peaks were measured relative to known compounds and these areas were then cut out and counted by liquid scintillation spectrometry.

When α -2-, α -3-, and α -4-L-fucosyltransferase activities occur together in an enzyme source, the substrate N-acetyllactosamine functions as an acceptor for both the α -2- and the α -3-transferases, and lacto-N-biose I as an acceptor for both α -2- and α -4-transferases. Difucosyl products are seldom observed and the products of transfer to the 0-2 position of the terminal galactose residue (2'-fucosyllactosamine and 2'-fucosyllacto-N-biose I, R_{Lac} 1.0 and 0.94, respectively, in solvent 1) are separable from the product of transfer to the 0-3 (3-fucosyllactosamine, R_{Lac} 0.75 in solvent 1) or the 0-4 (4-fucosyllacto-N-biose I, R_{Lac} 0.68 in solvent 1) position of the subterminal N-acetylglucosamine unit. In contrast to these two acceptor substrates the phenyl- β -D-galactoside acceptor cannot be difucosylated.

α -3-N-acetyl-D-galactosaminyl transferase (UDP-N-acetylgalactosamine:2-fucosylgalactoside α -3-N-acetylgalactosaminyltransferase) activity was measured with incubation mixtures consisting of enzyme source 20 μ l, UDP-N-acetyl-D-[¹⁴C]-galactosamine (1.3 nmol, 124 000 cpm), adenosine triphosphate (0.5 μ mol), $MnCl_2$ (2.0 μ mol), 2'-fucosyllactose (0.25 μ mol), Triton X-100 (500 μ g), and N-acacodylate buffer pH 6.0 (5.0 μ mol) in a total volume of 95 μ l. The mixture was incubated for four hours at 37°C. The products were separated on Whatman DE 81 paper in propan-1-ol/ethylacetate/pyridine/water, 5:1:1:3 (by vol) for 16 hours. Duplicate determinations of glycosyltransferase activities varied by less than 10%.

GLYCOLIPID EXTRACTION

A modification of the method described by Hakomori and Kannagi was used.²⁴ The tissues

were homogenised in isopropanol-hexane-water (55:25:20), sonicated, and centrifuged. The glycolipids in the supernatant were acetylated and subjected to florisil chromatography,²⁵ deacetylated, and the neutral glycolipid fraction was isolated by diethyl-aminoethyl-Sephadex chromatography.²⁶

THIN LAYER CHROMATOGRAPHY IMMUNOSTAINING

A modification of the method described by Magnani *et al*²⁷ was used. Briefly, approximately 20 μ g of total neutral glycolipids were applied on a high performance thin layer chromatography (HPTLC) plate, and the plate was developed in a mixture of chloroform/methanol/water (50:40:10). Subsequently, the plate was blocked in 5% bovine serum albumin in phosphate buffered saline and incubated with monoclonal antibody, followed by incubation with rabbit antimouse IgG or IgM and ¹²⁵I-protein A solution, successively. The dried plate was subjected to autoradiography for 18 to 24 hours by use of an intensifying screen and Kodak X-omat AR5 film.

IMMUNOHISTOCHEMISTRY

Formalin fixed 4 μ m sections were deparaffinised and rehydrated. Endogenous peroxidase was reacted with 0.03% hydrogen peroxide in methanol for 30 min. After washing, monoclonal antibodies were applied for 16 hours at 4°C. The sections were washed and sequentially incubated with rabbit antimouse immunoglobulins, biotinylated swine antirabbit immunoglobulins, and avidin-biotin-peroxidase complex, each step separated by washings in Tris-phosphate buffered saline (composition see above). Bound enzyme was visualised with 0.04% 3-amino-9-ethylcarbazole in N, N dimethylformamide 0.01% hydrogen peroxide.

Results

GLYCOSYLTRANSFERASE ACTIVITY IN NORMAL AND MALIGNANT RECTUM

α -2-L-fucosyltransferases

α -2-L-fucosyltransferase activity was assayed with phenyl- β -D-galactoside, N-acetyllactosamine, and lacto-N-biose I as acceptor substrates. In Le(a-b+) subjects we found a significantly higher level of this transferase in tumour tissue compared with normal tissue with

TABLE 1 α -2-L-fucosyltransferase activity in tumour tissue and normal tissue from rectum

Blood group	Tissue	No	Acceptor phenyl- β -D-galactoside		Lacto-N-biose I		N-acetyllactosamine	
			Median	Lower and upper quartile	Median	Lower and upper quartile	Median	Lower and upper quartile
Le(a-b+)	Normal	4	0.1	(0.1-0.3)	0.3	(0.1-0.3)	0.1	(0.1-0.1)
Le(a-b+)	Tumour	10	5.3	(2.6-8.7)	3.3	(1.6-4.2)	2.4	(1.6-3.7)
Le(a-b-)	Tumour	5	1.0	(0.8-1.3)	1.3	(1.3)	1.7	(1.6-1.8)
Le(a-b-)	Normal	3	0.3	(0.2-0.3)	0.2	(0.1-0.2)	0.2	(0.1-0.2)

NS=not significant at the $p=0.05$ level.

The activity was measured as pmol [¹⁴C] labelled fucose incorporated into the acceptor/h/mg. Mann-Whitney two tailed test was used for statistical comparison.

TABLE II α -3- and α -4-L-fucosyltransferase activity in tumour tissue and normal tissue from rectum

Blood group	Tissue	No	α -3-L-fucosyltransferase		α -4-L-fucosyltransferase	
			Median	Lower and upper quartile	Median	Lower and upper quartile
Le(a-b+) and Le(a+b-)	Tumour	14	15.5	(12.6-17.1)	8.4	(7.1-10.0)
Le(a-b+) and Le(a+b-)	Normal	7	11.9	(7.2-14.9)	10.1	(7.6-11.7)

*NS=not significant at the $p=0.05$ level (Mann-Whitney two tailed test). α -4-L-fucosyltransferase was significantly lower than α -3-L-fucosyltransferase in tumour tissue ($p<0.001$). The activity was measured as pmol [14 C] fucose incorporated into N-acetylglucosamine (α -3-L-fucosyltransferase) and lacto-N-biose I (α -4-L-fucosyltransferase)/h/mg.

all three acceptor substrates (Table I). Le(a+b-) individuals are believed to have a low or absent expression of α -2-L-fucosyltransferase in mucin secreting tissue. We therefore compared α -2-L-fucosyltransferase activity in tumour tissue from Le(a-b+) subjects with that from Le(a+b-) subjects (Table I). We found significantly lower activity in tumour tissues from Le(a+b-) subjects ($n=5$) when phenyl- β -D-galactoside and lacto-N-biose I were used as acceptor substrates. Lower activity was also observed with N-acetylglucosamine as substrate in tissues from Le(a+b-) subjects, but the difference was not significant. In Le(a+b-) subjects it was remarkable to find a significantly higher α -2-L-fucosyltransferase activity in tumour tissue compared with normal tissue with all three acceptor substrates (Table I).

α -3- and α -4-L-fucosyltransferases

No significant difference could be found between the activities in tumour tissue and normal tissue (Table II). α -3-L-fucosyltransferase was active in all samples examined, whereas α -4-L-fucosyltransferase activity was related to the Lewis type of the subject, being almost absent in Le(a-b-) subjects. In tumours, the α -3-L-fucosyltransferase was significantly higher than the α -4-L-fucosyltransferase, whereas both showed almost equal activity in normal tissue. α -2-L-fucosyltransferases compete with both the α -3- and α -4-L-fucosyltransferases for a mutual acceptor substrate (Fig 1). To evaluate alterations in this competitive action we calculated the ratios between the enzyme activities in each subject (Fig 2). The individual approach was necessary to eliminate interindividual variation. The α -4/ α -2-L-fucosyltransferase ratio measured for both enzymes with lacto-N-biose I as substrate was significantly lower ($p<0.005$) in tumour tissue

Figure 2: Ratios between fucosyltransferase activities in tumour tissue from rectum (10 subjects), and in different locations of normal colon (four subjects). Only secretors are shown. Circles indicate median; bars indicate lower and upper quartile. Open circles: α 4/ α 2 ratio measured with lacto-N-biose I as acceptor. Closed circles: α 3/ α 2 ratio measured with N-acetylglucosamine as acceptor.

from rectum compared with normal rectal tissue (Fig 2). The α -3/ α -2-L-fucosyltransferase ratio measured for both enzymes with N-acetylglucosamine as substrate was also significantly lower ($p<0.005$) in rectal tumour tissue than in normal rectal tissue.

α -3-N-acetyl-D-galactosaminyltransferase

This transferase was assayed at pH 6, which is the pH optimum for the A¹ transferase.²⁸ Activity was only found in blood group A subjects (Table III). The activity in tumour tissue was lower (though not significant) than the activity in normal tissue. The individual ratios between the A¹ transferase and the fucosyltransferases with which it competes for acceptor substrates were not significantly altered.

GLYCOSYLTRANSFERASE ACTIVITY THROUGHOUT THE NORMAL COLON

The activity of the α -2-L-fucosyltransferase was higher in biopsy tissue from the caecum than from all other parts of the normal colon (Fig 3A). The high activity in caecum was more pronounced when lacto-N-biose I and phenyl- β -D-galactoside were used as acceptors than when N-acetylglucosamine was used. The α -3-, α -4-, and A¹ transferases (data for the A¹ transferase are not shown) were similar in activity throughout the colon. Because of this the individual α -4/ α -2 ratio, and to a lesser extent the individual α -3/ α -2 ratio, were much lower in caecum than in the rest of the colon (Fig 2). None of these regional

TABLE III α -3-N-acetyl-D-galactosaminyltransferase (A¹ transferase) activity in tumour tissue and normal tissue from rectum

Blood group	Tissue	No	A ¹ transferase		Ratio		Ratio	
			Median	Lower and upper quartile	A ¹ transferase/ α -4-L-fucosyltransferase*	Lower and upper quartile	A ¹ transferase/ α -3-L-fucosyltransferase	Lower and upper quartile
0	Tumour	7	1.5	(1.0-3.3)	7.1	(4.3-12.2)	5.7	(3.9-8.1)
A ¹	Tumour	7	68.1	(48.9-86.5)				
A ¹	Normal	5	103.2	(71.2-139.1)	7.8	(6.0-11.3)	8.4	(6.4-11.9)

*Only Lewis positive subjects were included. NS=not significant. Figures indicate pmol [14 C]GalNAc incorporated into 2'-fucosyllactose/h/mg. Mann-Whitney two tailed test was used for statistical comparison.

Figure 3: (A) Glycosyltransferase activity throughout the normal colon of four secretors. Open triangles = α -2-fucosyltransferase (FT) activity with N-acetyllactosamine as acceptor. Open circles: α -2-FT activity with phenyl- β -D-galactoside as acceptor. Diamonds: α -2-FT activity with lacto-N-biose I as acceptor. Squares: α -4-FT activity with lacto-N-biose I as acceptor. Closed triangles: α -3-FT activity with N-acetyllactosamine as acceptor. Symbols indicate median value. (B) Glycosyltransferase activity throughout the normal colon of three non-secretors. Symbols as in (A).

differences was observed in the three non-secretors examined (Fig 3B).

GLYCOLIPID PATTERN IN NORMAL HUMAN COLONIC MUCOSA AND COLONIC ADENOCARCINOMAS

Extracted neutral glycolipids were subjected to immunostaining after separation on HPTLC plates. The glycolipid profile of caecum was compared with that of rectum in both normal and malignant tissue (Fig 4). Four normal subjects and two subjects with carcinomas of the rectum, as well as two with carcinomas of the caecum were examined. The staining intensity of the positive bands showed slight interindividual variation. In caecum both Le^a and Le^b active glycolipids were present in normal as well as malignant tissue. The type 2 chain structures Le^x and Le^y were only present in appreciable amounts in malignant tissue. In rectum only Le^a active glycolipids were present in both normal and malignant mucosa, whereas Le^b, Le^x, and Le^y antigens occurred only in glycolipids extracted from malignancies. The type 2 chain structures Le^x and Le^y were present in glycolipids with penta- and hexasaccharide chains as well as in slower migrating components. The type 1 chain structures Le^a and Le^b were detected only in glycolipids with penta- and hexasaccharide chains, respectively.

Figure 4: Characterisation of blood group active total neutral glycosphingolipids in normal and malignant tissue from the caecum and rectum. The tissues were extracted in organic solvents, acetylated and subjected to florisil chromatography, deacetylated, and separated by DEAE Sephadex chromatography. The total neutral glycosphingolipids obtained were separated according to sugar-chain length on high performance thin layer chromatography plates, and reacted with monoclonal antibodies against Le^a, Le^b, Le^x, and Le^y antigens. Binding of antibodies was visualised by ¹²⁵I-protein A autoradiography.

IMMUNOHISTOCHEMICAL LOCATION OF BLOOD GROUP ANTIGENS

Normal specimens from rectum of both Le(a+b-) and Le(a-b+) subjects showed staining only with monoclonal antibodies against Le^a antigens, all other antibodies were unreactive (Table IV). Normal colonic specimens from caecum were stained in Le(a-b+) subjects by all the antibodies examined that were against type 1 chain H, Le^a, Le^b, and A antigens, whereas type 2 chain specific antibodies against Le^x, Le^y, H, and A antigens reacted more weakly (Table IV). In Le(a+b-) subjects only the type 1 chain anti-Le^a specific antibody showed strong staining in caecum (Table IV). In carcinomas from the

TABLE IV Immunohistochemical localisation of carbohydrate antigen expression in tissue from A₁ secretors and non-secretors

Tissue	No	Type 1 chain (Galβ1-3GlcNAc-R)					Type 2 chain (Galβ1-4GlcNAc-R)				
		Le ^a	H	Le ^b	A	ALe ^b	Le ^a	H	Le ^b	A	ALe ^b
<i>Secretors:</i>											
Normal rectum	4	+	—	—	—	—	—	—	—	—	—
Normal caecum	4	+	+	+	+	+	(+)*	+	(+)	+	(+)
Cancerous rectum	5	+	+	+	+	+	+	+	+	+	+
<i>Non-secretors (3 blood group O, 1 blood group A†):</i>											
Normal rectum	4	+	—	—	—	—	—	—	—	—	—
Normal caecum	4	+	—	—	(+)	—	(+)	(+)	—	(+)	—
Cancerous rectum	4	+	—	+	+	(+)	+	(+)	+	+	+

ABH secretor state determined on saliva specimens. Monoclonal antibodies were applied to tissue sections from secretors and visualised as described in methods. *Symbols in parenthesis= weakly and partially positive. †A antigens were only detected in the A subject.

rectum, however, all the antibodies gave positive staining patterns in Le(a-b+) subjects, and, remarkably, all but anti-H type 1, also in Le(a-b-) subjects (Table IV).

Discussion

The major difference observed in glycosyltransferase activity between normal rectal mucosa and rectosigmoidal carcinomas was the significantly increased activity of the α-2-L-fucosyltransferases in carcinomas from both secretors and non-secretors. This is in agreement with a previous report²⁹ showing that, in comparison with normal colonic mucosa, the activity of the α-2-L-fucosyltransferases, as measured with phenyl-β-D-galactoside, was increased in colonic carcinomas of ABH secretors. The present study also showed that the α-2-L-fucosyltransferases were significantly higher in normal caecum than in normal rectum from secretors. All other transferases studied were of the same magnitude throughout the colon.

The biosynthesis of H, A and Le^b blood group determinants from their common precursor largely depends on the activity of the α-2-L-fucosyltransferases (Fig 1). These antigens could be shown in rectal carcinomas and in normal caecum of secretors, but not in normal rectum. Therefore, we suggest that the α-2-L-fucosyltransferases control the expression of these blood group antigens in both normal and malignant human colon.

Non-secretors differed from secretors in the way that the α-2-L-fucosyltransferase activity was low in all regions of the normal colon, including caecum. It was remarkable, however, that rectal carcinomas from non-secretors showed a significant increase in α-2-L-fucosyltransferase activity with all three acceptor substrates, the main difference in enzyme activity between rectal carcinomas from secretors and non-secretors being higher activity with phenyl-β-D-galactoside and lacto-N-biose I in secretors. This interesting enzymatic finding in non-secretors was found to lead to antigen synthesis, as Le^b, A, and ALe^b antigens started to appear in rectal carcinomas from these subjects.

In colon mucosa there seem to be two forms of the α-2-L-fucosyltransferase, one that is Se gene dependent and has a preference for type 1 chain acceptors, and one, the H gene dependent, that is expressed independently of the Se gene and has a low activity with both type 1 and 2 chain acceptors.³⁰⁻³² In non-secretors no activity was present in caecum, and the activity observed

in rectal carcinomas of non-secretors showed similar levels of activity with both N-acetyl-lactosamine and lacto-N-biose I as acceptor substrates. These data indicate that in non-secretors the H gene dependent α-2-L-fucosyltransferase is increased in colorectal carcinomas. In secretors the present data show higher levels of α-2-L-fucosyltransferase activity in normal caecum than in other regions of the colon with both type 1 and type 2 chain oligosaccharides and with phenyl-β-D-galactoside as acceptor. Increased activity with these acceptors was also seen in rectosigmoidal carcinomas compared with normal tissue from the same region (Table I). Phenyl-β-D-galactoside is an efficient acceptor for the α-2-L-fucosyltransferases and clearly distinguishes between secretors and non-secretors when used to assay material from submaxillary glands.³³ The high activity reported in secretors with this acceptor and with lacto-N-biose I, and the difference in activity between secretors and non-secretors in both carcinomas and normal caecum, indicate that in secretors it is the Se gene dependent transferase which is increased in caecum. In colorectal carcinomas from secretors an increase in activity of the Se gene dependent α-2-L-fucosyltransferase or an increase in both this transferase and the H gene dependent transferase could explain the present findings.

The α-2-L-fucosyltransferase competes with α-3- and α-4-L-fucosyltransferases for mutual acceptor substrates (Fig 1). If the α-2-L-fucosyltransferase acts first the H antigen and subsequently Le^a and Le^b antigens are formed, whereas if the α-3- and α-4-L-fucosyltransferases act first, Le^a and Le^b antigens are formed. The last two antigens are poor substrates for the α-2-L-fucosyltransferase and cannot be fucosylated to form Le^a and Le^b antigens. To determine the prevailing pathway of type 1 and 2 chain biosynthesis (Fig 1), the individual ratios between α-4/α-2- and α-3/α-2-fucosyltransferases measured respectively with lacto-N-biose I and N-acetyl-lactosamine were calculated for each subject. Both ratios were significantly decreased in tissue from the caecum and rectal tumour tissue compared with normal rectal tissue. Thus in normal rectum type 1 chain Le^a antigens are synthesised and no H, Le^b, and A antigens are formed; a result which can be explained by the high α-4/α-2- enzyme ratio. This ratio is lower in carcinomas and in normal caecum, and hence both Le^a and H, Le^b and A antigens are formed. Although type 2 chain blood group antigens may be synthesised by enzymes that also work on type

1 chains, their biosynthesis is obviously regulated in a different way because the positional isomer of Le^a, namely the Le^x antigen, is not detectable to any great degree in normal rectum. A lower α -3/ α -2- ratio in carcinomas will only lead to formation of H type 2 chain antigens if the appropriate precursor structures are present. The lack of Le^x antigens in normal rectum indicates that this is not the case. Regulation of type 2 chain biosynthesis by enzymes that work on the synthesis of the type 2 chain core structure,⁵ as well as by enzymes that work on terminal fucosylation, thus seems probable.

The immunochemical characterisation of blood group antigen expression obtained by immunostaining of blood group active glycolipids verified the immunohistochemical detection of antigens in this and previous studies.⁶⁻⁸ As previously shown,¹¹⁻¹³ only type 2 chain antigens are synthesised on extended chains; however, the mechanism leading to this difference between type 1 and 2 chains is at present unknown.

Reduction of α -3-N-acetyl-D-galactosaminyl-transferase activity has previously been shown in intestinal carcinoma tissue.³⁴ A reduction of this enzyme does not, however, seem to be important in the large bowel, as the present data confirm that carcinomas from A individuals can synthesise blood group A antigens when the α -2-fucosyltransferase level is increased.

The results suggest that the critical event required for blood group H, Le^b, A, and B antigen biosynthesis, in both normal mucosa from the caecum and rectal carcinomas, is the expression of α -2-L-fucosyltransferase activity.

This study was supported by the Danish Cancer Society.

- 1 Oriol R, Le Pendu J, Mollicone R. Genetics of ABO, H, Lewis, X, and related antigens. *Vox Sang* 1986; 51: 161-71.
- 2 Dabelsteen E, Buschard K, Hakomori S, Young WW. Pattern of distribution of blood group antigens on human epidermal cells during maturation. *J Invest Dermatol* 1984; 82: 13-7.
- 3 Dabelsteen E, Vedtofte P, Hakomori S, Young WW. Carbohydrate chains specific for blood group antigens in differentiation of human oral epithelium. *J Invest Dermatol* 1982; 79: 3-7.
- 4 Feizi T. Demonstration by monoclonal antibodies that carbohydrate structures of glycoproteins and glycolipids are oncodevelopmental antigens. *Nature* 1985; 314: 53-7.
- 5 Holmes EH, Ostrander GK, Clausen H, Graem N. Oncofetal expression of Le^x carbohydrate antigens in human colonic adenocarcinomas. *J Biol Chem* 1987; 262: 11331-8.
- 6 Yuan M, Itzkowitz SH, Palekar A, et al. Distribution of blood group antigens A, B, H, Lewis^a, and Lewis^b in human normal, fetal, and malignant colonic tissue. *Cancer Res* 1985; 45: 4499-511.
- 7 Itzkowitz SH, Yuan M, Fukushi Y, et al. Lewis^a- and sialylated Lewis^x-related antigen expression in human malignant and non-malignant colonic tissues. *Cancer Res* 1986; 46: 2627-32.
- 8 Kim YS, Yuan M, Itzkowitz SH, et al. Expression of Le^x and extended Le^x blood group-related antigens in human malignant, premalignant, and non-malignant colonic tissues. *Cancer Res* 1986; 46: 5985-92.
- 9 Szulman AE. The histological distribution of the blood group substances in man as disclosed by immunofluorescence III. The A, B, and H antigens in embryos and fetuses from 18 mm length. *J Exp Med* 1984; 119: 503-15.
- 10 Szulman AE, Marcus DM. The histological distribution of the blood group substances in man as disclosed by immunofluorescence VI. The Le^a and Le^b antigens during fetal development. *Lab Invest* 1973; 5: 565-73.
- 11 Kaizu T, Levery SB, Nudelman E, Kamp RES, Hakomori S. Novel fucolipids of human adenocarcinoma VI. Monoclonal antibody specific for trifucosyl Le^x (III'Fuc V'Fuc VI'Fuc L_{C6}), which does not crossreact with Le^x antigen, and a possible conformational epitope structure. *J Biol Chem* 1986; 261: 11254-8.
- 12 Fukushi Y, Hakomori S, Nudelman E, Cochran N. Novel fucolipids accumulating in human adenocarcinoma: II. Selective isolation of hybridoma antibodies that differentially recognise mono-, di-, and trifucosylated type 2 chain. *J Biol Chem* 1984; 259: 4681-5.
- 13 Magnani JL, Nilsson B, Brockhaus M, et al. A monoclonal antibody-defined antigen associated with gastro-intestinal cancer is a ganglioside containing sialylated lacto-N-fucopentaose II. *J Biol Chem* 1982; 257: 14365-9.
- 14 Kannagi R, Fukushi Y, Tachikawa T, et al. Quantitative and qualitative characterization of human cancer-associated serum glycoprotein antigens expressing fucosyl or sialyl-fucosyl type 2 chain polylactosamine. *Cancer Res* 1986; 46: 2619-26.
- 15 Fukushima K, Hirota M, Terasaki PI, et al. Characterisation of sialosylated Lewis^x as a new tumour-associated antigen. *Cancer Res* 1984; 44: 5279-85.
- 16 Sears HF, Herlyn M, DelVillano B, Steplewski Z, Koprowski H. Monoclonal antibody detection of a circulating tumor-associated antigen II. A longitudinal evaluation of patients with colorectal cancer. *J Clin Immunol* 1984; 2: 141-9.
- 17 Abe K, McKibbin JM, Hakomori S. The monoclonal antibody directed to difucosylated type 2 chain (Fucal-2GalBI-4 [Fucal-3]GlcNAc; Y determinant). *J Biol Chem* 1983; 258: 11793-7.
- 18 Abe K, Levery SB, Hakomori S. The antibody specific to type 1 chain blood group A determinant. *J Immunol* 1984; 132: 1951-4.
- 19 Clausen J, Levery SB, McKibbin JM, Hakomori S. Blood group A determinants with mono- and difucosyl type 1 chain in human erythrocyte membranes. *Biochemistry* 1985; 24: 3578-86.
- 20 Clausen H, McKibbin JM, Hakomori S. Monoclonal antibodies defining blood group A variants with difucosyl type 1 chain (ALe^b) and difucosyl type 2 chain (ALe^x). *Biochemistry* 1985; 24: 6190-4.
- 21 Clausen H, Levery SB, Nudelman E, Baldwin M, Hakomori S. Further characterization of type 2 and type 3 chain blood group A glycosphingolipids from human erythrocyte membranes. *Biochemistry* 1986; 25: 7075-85.
- 22 Betteridge A, Watkins WM. Variant forms of α -2-L-fucosyltransferase in human submaxillary glands from blood group ABH "secretor" and "non-secretor" individuals. *Glycoconjugate J* 1985; 2: 61-78.
- 23 Ørntoft TF, Wolf H, Watkins WM. Activity of the human blood group ABO, Se, H, Le, and X gene-encoded glycosyltransferases in normal and malignant bladder urothelium. *Cancer Res* 1988; 48: 4427-33.
- 24 Hakomori S, Kannagi R. Carbohydrate antigens in higher animals. In: Weir DM, Herzenberg LA, Blackwell CC, eds. *Handbook of experimental immunology*. Vol 4. Oxford: Blackwell Scientific Publications 1986: 1-39.
- 25 Saito T, Hakomori S. Quantitative isolation of total glycosphingolipids from animal cells. *J Lipid Res* 1971; 12: 257-62.
- 26 Yu RK, Ledeen RW. Gangliosides of human, bovine, and rabbit plasma. *J Lipid Res* 1972; 13: 680-6.
- 27 Magnani JL, Smith DF, Ginsburg V. Detection of gangliosides that bind cholera toxin: direct binding of ¹²⁵I-labelled toxin to thin-layer chromatograms. *Anal Biochem* 1980; 109: 399-402.
- 28 Schachter H, Michaels MA, Tilley CA, Crookston MC, Crookston J. Qualitative difference in the N-acetyl-D-galactosaminyltransferases produced by human A¹ and A² genes. *Proc Natl Acad Sci USA* 1973; 70: 220-4.
- 29 Piller F, Schenkel-Brunner H, Tuppy H. Altered biosynthesis of blood group related antigens in neoplastic colon mucosa. In: Schauer R, Kramer MF, Vliegenhart JFG, Wiegandt H, eds. *Glycoconjugates. Proceedings of the Fifth International Symposium Kiel*. [Abstract] Stuttgart. Georg Thieme, 1979: 639-40.
- 30 Kumazaki T, Yoshida A. Biochemical evidence that secretor gene, Se, is a structural gene encoding a specific fucosyltransferase. *Proc Natl Acad Sci USA* 1984; 81: 4193-7.
- 31 Watkins WM, Greenwell P. Biosynthesis of blood group ABH antigens: genetic regulation and tissue distribution of glycosyltransferases. *Transplant Proc* 1987; 19: 4415-6.
- 32 Le Pendu J, Oriol R, Juszczak G, et al. α -2-L-Fucosyltransferase activity in sera of individuals with H-deficient red cells and normal H antigen in secretions. *Vox Sang* 1983; 44: 360-5.
- 33 Chester MA, Yates AD, Watkins WM. Phenyl β -D-galactopyranoside as an acceptor substrate for the blood-group H gene-associated guanosine diphosphate L-fucose: β -D-galactosyl α -2-L-fucosyltransferase. *Eur J Biochem* 1976; 69: 583-92.
- 34 Stellner K, Hakomori S, Wagner GA. Enzymatic conversion of H₁ glycolipid and deficiency of these enzyme activities in adenocarcinoma. *Biochem Biophys Res Commun* 1973; 55: 439-45.