Glutathione S-transferase in humans in health and disease

P C Hayes, I A D Bouchier, G J Beckett

Most biological processes are catalysed by means of enzymes and study of these in health and disease is important for our understanding of both biology and disturbed function. The glutathione S-transferases (GST; E.C.2.5.1.18) are a complex multigene family of enzymes that are widely distributed in the animal kingdom.¹ These enzymes possess many biological functions, the most important of which is detoxification, conjugating reduced glutathione with a large number of electrophiles.² Such conjugation reactions result in the synthesis of mercapturic acids and represent an important excretory route for xenobiotics including carcinogens, toxins, and drugs.¹ As well as being concerned in the metabolism of xenobiotics, GST are involved in the metabolism of endogenous substances such as leukotriene A₄ and prostaglandin A₁.^{3,4} GST are also involved in non-substrate binding to such substances as bilirubin and bile acids and have an important role in hepatic anion transport. Several GST isoenzymes possess selenium-independent glutathione peroxidase activity with certain organic hydroperoxides such as linoleic and arachidonic hydroperoxide but not with hydrogen peroxide.⁵

In humans the cytosolic GST isoenzymes are dimeric and can be subdivided on the basis of their physicochemical and immunological properties into three major groups, namely pi, alpha, and mu class GST (Table⁶). In humans two immunologically distinct subunits of the alpha class GST have been named B₁ and B₂. A fourth GST isoenzyme has recently been identified in the liver and has been purified from the microsomal subcellular fraction. By contrast with the cytosolic enzymes, the microsomal GST is trimeric containing subunits with a molecular mass of 17 000 kDa.^{7,8}

In this review we concentrate on the distribution of these GST isoenzymes in various human tissues in both health and disease. We also discuss the value of measuring GST isoenzymes in plasma as a means of identifying tissue damage and neoplasia.

Tissue distribution of GST isoenzymes

Alpha class GST have been shown to be the predominant isoenzymes in human hepatocytes with the microsomal and mu class GST present in small quantities.⁹ Mu class GST is subject to polymorphism and, in the Scottish population only about 55% of individuals express this GST class.¹⁰ Pi class GST in the adult human liver is only expressed in the biliary epithelium. Why pi class GST is expressed so strongly in biliary epithelium including the gall bladder is unclear, but large quantities of this isoenzyme can be identified in human bile.¹¹ In most hepatocellular

carcinomas alpha class GST are expressed, whereas pi class GST is the predominant enzyme in cholangiocarcinoma.¹²

The heterogeneity of GST isoenzyme distribution has also been shown in small bowel epithelium. Pi and mu class GST are present in cells lining both the villi and crypts, whereas alpha class GST has only been identified in the epithelium of villi and microsomal GST only in cells lining the intestinal crypts.⁹ In the colon, both pi and mu class GST (the latter in 50% only) are present in the epithelium, whereas alpha class and microsomal GST are absent. Overexpression of pi class GST in carcinoma of the colon has been reported while expression of the other isoenzymes is variable.¹³

In the pancreas, pi class GST has been identified in the centroacinar cells and ducts but not in the exocrine acini nor in the islets of Langerhans.¹⁴ Alpha class GST are the predominant isoenzymes in the acini and are also present in pancreatic ducts and mu class GST is predominantly present in the islets of Langerhans. Microsomal GST is variably present and shows no constant pattern of distribution. Pancreatic adenocarcinoma strongly expresses pi class GST but the other isoenzymes are generally absent.¹⁴

The explanation for the heterogeneity of distribution of the various GST isoenzymes within and between tissues is unclear, but one possible reason is that they subserve different functions in different tissues, sometimes acting as carrier proteins rather than being metabolically active. The overexpression of certain isoenzymes, principally pi class GST in intestinal and pancreatic tumours and alpha class GST in hepatocellular carcinoma, may simply represent clonal expansion of those cells expressing these isoenzymes. It is particularly noteworthy that in pancreatic carcinoma, which is believed to arise either from ductular or acinar cells,¹⁵ the pi class and not alpha class GST is strongly expressed. Recent evidence by Pour¹⁶ suggests that the centroacinar cells may give rise to the pseudo-ductular complexes which are believed to be preneoplastic lesions. This distribution of GST isoenzymes would support such a hypothesis.

Nomenclatures for human GST⁶

GST isoenzyme	Family	Subunit (mol wt)	Isoelectric point
B ₁ B ₁	ε	25 900	8.9
B ₁ B ₂	δ	25 900	8.75
B ₂ B ₂	α	25 900	8.4
μ	μ	26 700	6.1
ψ	μ	26 600	5.5
π	π	24 800	4.8
Microsomal	—	17 300	Not determined

University Departments of Medicine and Clinical Chemistry, Royal Infirmary, Edinburgh EH3 9YW, Scotland
P C Hayes
I A D Bouchier
G J Beckett

Correspondence to:
Dr P C Hayes, Department of Medicine, Royal Infirmary, Edinburgh EH3 9YW.

Accepted for publication
10 September 1990

An alternative explanation for the particular expression of GST in tumours is that selective overexpression of certain GST isoenzymes confers replicative advantage to dedifferentiated cells. Cellular drug resistance is commonly accompanied by increases in GST and in the drug-binding membrane-bound P-glycoprotein, and a decrease in cytochrome P450.¹⁷ Many parallels between acquired drug resistance and preneoplasia exist and support the selective advantage hypothesis.

GST measurements in plasma as markers of disease

The GST isoenzymes catalyse the conjugation of glutathione to a wide range of compounds of which 1-chloro-2-4-dinitrobenzene is most frequently used as the substrate to measure the activity of human GST.¹⁸ There are, however, several problems associated with activity measurements. Firstly, normal plasma activity is low and a precise measurement of activity is difficult to achieve.¹⁹ Secondly, the GST binds a number of anions such as bile salts and bilirubin which inhibit enzymic activity.²⁰ In liver disease, when high plasma concentrations of these anions occur the activity of the GST released from the hepatocyte into plasma is inhibited thus decreasing the clinical sensitivity of the test. Another major problem with activity measurements is their poor organ specificity because the various substrates do not allow the isoenzymes of GST to be differentiated. Because of these problems assays of plasma or serum GST based on activity measurements have proved to be of little clinical value.^{19, 21} Immunoassays have been described that allow the precise and specific measurement of each of the three cytosolic classes of GST²²⁻²⁶ and also the measurement of individual GST subunits within a class such as the B₁ and B₂ subunits which comprise the alpha class GST.^{22, 27} Immunoassay measurements measure GST concentrations even when they are enzymatically inactive because of the presence of inhibitors such as bilirubin or bile salts. Using these immunoassays it has become apparent that GST measurements may have a clinical role in the diagnosis and monitoring of patients with hepatobiliary disease and malignancy.²⁸ At present the only methodological problem relates to the speed of the assay which takes at least 24 hours. The use, however, of labelled antibody rather than labelled antigen assays should overcome this problem and enable results to be provided in less than three hours.

GST in liver damage

The alpha class GST, which comprise the immunologically distinct subunits B₁ and B₂, constitute as much as 3% of cytosolic protein in the hepatocyte. The subunits are distributed uniformly across the liver lobule; this contrasts with the aminotransferases which are found predominantly in the periportal hepatocytes.²⁹ The physicochemical properties and hepatic distribution of the B₁ and B₂ subunits mean that they are released quickly and in large quantities from damaged hepatocytes into the plasma. The

plasma half life of the B₁ and B₂ subunits is less than 60 minutes; this contrasts with alanine aminotransferase which has a half life of approximately 48 hours.³⁰ These characteristics of the B₁ and B₂ subunits indicate that their measurement in plasma should provide an early and sensitive measure of hepatocellular damage even when only centrilobular hepatocytes are involved. The short half life of the GST should also allow early recognition of when active liver damage has stopped, and measurements of the B₁ and B₂ subunits might therefore provide a useful means of investigating mechanisms of hepatotoxicity.^{30, 31}

Homogenates of normal human liver obtained at necropsy contain extremely low concentrations of pi class GST³² and immunohistochemical examination has shown that this GST is confined to the cells of the biliary epithelium.^{13, 33} In alcoholic liver disease, however, sinusoidal macrophages, Kupffer cells, and hepatocytes express GST-pi.³⁴ There are also data to suggest that biliary epithelial cells secrete GST-pi into bile possibly as a mechanism to export potentially harmful toxins from the cell.¹¹

ACUTE LIVER DAMAGE

(i) Paracetamol poisoning

After paracetamol poisoning abnormal plasma concentrations of GST B₁ subunits (B₁) are seen in approximately 90% of patients while alanine aminotransferase becomes abnormal in about 40%. In addition, abnormal concentrations of B₁ are observed within four hours of paracetamol ingestion whereas abnormalities in alanine aminotransferase are not recognised for at least 12 hours.^{30, 35} The results of plasma B₁ measurements show two distinct phases of hepatotoxicity in paracetamol poisoning.³⁰ The early phase occurs between four and 12 hours after the overdose and is characterised by increasing B₁ concentrations; this phase may be reversed by giving intravenous N-acetylcysteine. The second phase of hepatotoxicity occurs up to 24–36 hours after overdose when concentrations of B₁ plateau at approximately 100 times the upper reference interval. In this second phase of hepatotoxicity the hepatocytes are damaged irreversibly and ultimately hepatic necrosis occurs. The measurement of plasma B₁, on admission, may provide a good indication of which patients will develop subsequent liver damage.³⁵

(ii) Birth asphyxia

Early evidence of liver damage after birth asphyxia is provided by plasma B₁ measurements.³⁶ In a study of 14 neonates with a clinical diagnosis of birth asphyxia abnormal plasma concentrations of B₁ were found within six hours of asphyxia in 11 whereas only seven had abnormal alanine aminotransferase activities at this time. In plasma sampled 24 hours after birth, values for alanine aminotransferase were abnormal in 10 neonates while B₁ remained abnormal in only six. The same study also showed that plasma B₁ concentrations were greater in normal infants than in adults despite

the fact that the concentration of the B₁ subunit in neonatal liver is significantly lower than in the adult. These data suggest that during the first few days of life there may be impaired hepatocellular integrity particularly centrilobular cells.

Although measurement of plasma B₁ or B₂ subunits in the adult seems to be equally sensitive in detecting impaired hepatocellular integrity, in the neonate B₂ measurements are of no value since this subunit seems to be poorly expressed in developing liver.³⁶

(iii) *Ethanol ingestion*

We have recently shown that plasma concentration of B₁ increased in heavy drinkers 60 minutes after 80 g of ethanol had been ingested over a 30 minute period. No increase in B₁ was observed when healthy volunteers, with a modest alcohol intake, were given the same acute alcohol load. Although as a group the heavy drinkers produced a rise in B₁ after alcohol ingestion, some showed no significant change in B₁.³⁷ These observations suggest that the measurement of plasma B₁ in response to an alcohol load may be a predictor of the sensitivity of an individual to alcohol induced liver damage.

(iv) *Halothane anaesthesia*

Halothane anaesthesia may be followed by unexplained hepatitis that may be mild or severe. The more severe form (type 2) has a 20–50% mortality but the nature of the mechanism and predisposing factors that relate to type 2 halothane hepatitis remain unclear, although hypersensitivity and familial constitutional susceptibility have been implicated.

The mild type 1 form of halothane hepatitis is associated with moderate increases in the transaminases. Production of toxic metabolites of halothane after a combination of reductive biotransformation of halothane and hepatic hypoxia have been implicated in halothane induced hepatotoxicity. Enflurane and isoflurane are associated with less severe hepatic problems than halothane.^{38–40}

Three hours after halothane anaesthesia small but significant increases in plasma B₁ concentration occur in approximately half of all patients which resolve after 12 hours. In approximately 10% of patients a secondary rise in plasma B₁ occurs 24 hours after anaesthesia.⁴¹ In contrast, isoflurane produces no significant change in plasma B₁ concentrations at three hours and the frequency of abnormal plasma B₁ concentrations at 24 hours is lower than for halothane.⁴² It seems likely that the increase in plasma B₁ concentrations three hours after halothane results from diminished hepatic blood flow, while it is possible that the secondary rise at 24 hours may be due to production of toxic halothane metabolites.^{41, 42}

Although changes in plasma B₁ concentrations may be seen after halothane anaesthesia, these studies have not shown any clinical value for measurement of the GST B₁ subunit as a predictor of patients who will subsequently develop clinical hepatic dysfunction.

CHRONIC LIVER DISEASE

(i) *Chronic active hepatitis*

An early study showed that all patients with chronic active hepatitis had raised serum concentrations of alpha class GST; serum GST concentrations correlated significantly with the activity of the disease assessed by histology, whereas aminotransferase activities showed no such correlation.³¹ We have recently produced evidence to suggest that serum B₁ may be of value in monitoring the effectiveness of treatment in patients receiving immunosuppressive drugs for autoimmune chronic active hepatitis. We studied 22 patients and found that serum B₁ concentrations were raised in 16 patients, whereas aspartate aminotransferase activities were abnormal in only six of the patients. This work led us to suggest that perhaps the aim of immunosuppressive treatment should be to normalise serum B₁ rather than aminotransferase levels.⁴³

(ii) *Cirrhosis*

Raised serum concentrations of B₁ occur in more than half of patients with cirrhosis.^{31, 44} In a study of 54 patients with biopsy proved alcoholic cirrhosis, abnormal levels of the aminotransferases and B₁ subunits were found in 28 patients, but these two measurements seemed to identify different populations.⁴⁴

An explanation for the observation of an abnormal B₁ concentration in the presence of a normal aspartate aminotransferase is that B₁ measurements are probably more sensitive at detecting alcohol induced centrilobular damage. The explanation for the converse pattern of results – that is, normal B₁ with raised aspartate aminotransferase – found in nine patients is less clear but this may be due to a change in GST expression from alpha class to pi class GST in the hepatocyte.³⁴

(iii) *Thyrotoxicosis and thyroxine replacement therapy*

Patients with thyrotoxicosis often have abnormal plasma B₁ concentrations.⁴⁵ This is not surprising as before effective treatment for hyperthyroidism was introduced serious hepatobiliary complications were commonly associated with the disease.⁴⁶ The observation, however, that many patients who are receiving longterm thyroxine replacement therapy for hypothyroidism have raised plasma B₁ concentrations⁴⁷ has fuelled the controversy regarding the necessity to monitor the dose of thyroxine by biochemical tests.⁴⁸

Patients who most commonly have abnormal plasma B₁ concentrations are those in whom plasma free thyroxine is raised and thyroid stimulating hormone suppressed to undetectable levels.⁴⁷ These observations have led to the suggestion that the aim of thyroxine replacement therapy should be to normalise plasma thyroid stimulating hormone as this reduces the prevalence of abnormal plasma B₁ concentrations.⁴⁹ Recent observations have complicated the issue for it seems that patients with spontaneous

hypothyroidism are more prone to develop impaired hepatocellular integrity when starting thyroxine replacement therapy than patients with radioiodine induced hypothyroidism.⁵⁰ The evidence for this is attributable to the observation that plasma B₁ concentrations increase significantly when the dosage of thyroxine is increased above 100 µg per day in patients with spontaneous hypothyroidism, while no significant change is found in patients with radioiodine induced hypothyroidism.

At present it is unknown whether overreplacement with thyroxine has any longterm deleterious effects on the liver because no retrospective or prospective studies have been reported. In animals, however, thyroid hormones increase the hepatotoxicity associated with administration of halogenated hydrocarbons such as halothane or carbon tetrachloride.^{51,52} In part, the increased hepatotoxicity of these compounds in thyroid hormone treated animals may be the result of a diminished hepatic GST content because the GST may be involved in the metabolism and detoxification of these compounds.⁵³

GST-mu and susceptibility to liver disease

The mu class GST have different substrate specificities than the alpha and pi class isoenzymes. A number of mu class GST occur but GST-mu is of particular interest as it is polymorphic and expressed in only approximately half of the normal population.^{23,28,54,55} Harada *et al* showed that only 25% and 20%, respectively, of patients with alcoholic liver disease and liver carcinoma expressed GST-mu, whereas the enzyme was expressed in 94% of patients with chronic hepatitis.⁵⁶ This study, however, comprised a very small number of patients; a much larger study is necessary to confirm these findings.

GST measurements in malignancy

There is considerable interest in the association between pi class GST and malignancy after the discovery that increased expression of this enzyme occurs in many tumours.^{32,57-59} In neoplasms of the lung, colon, and stomach the expression of GST-pi is increased approximately twofold when compared with matched normal tissue, but in kidney and liver no significant change in the expression of this isoenzyme occurs.³²

The expression of GST-pi may vary with the oestrogen receptor status in breast cancer; the levels of both GST-pi RNA and the protein are significantly higher in oestrogen receptor positive tumours when compared with oestrogen receptor negative tumours.⁶⁰

In contrast to GST-pi the expression of B₁ and B₂ subunits is suppressed in some tumours including stomach and kidney. In other tumour types, however, concentrations are unchanged with the exception of lung in which B₁ seems to show an increased expression.³²

The polymorphic GST-mu and its relation to malignancy is important. It has been claimed

that smokers who lack GST-mu are more susceptible to developing carcinoma of the lung compared with smokers who express the enzyme. This relation, however, only seems to hold true for adenocarcinoma of the lung.⁵⁵

The changes in GST expression that occur in malignancy have resulted in the use of GST measurements as tumour markers.

GST-pi measurements in malignancy

PLASMA GST MEASUREMENTS

Two studies suggest that the serum concentrations of GST-pi are increased in a wide range of malignant growths including gastric, oesophageal, colonic, pancreatic, and hepatobiliary cancers. In contrast, few abnormalities in serum GST-pi are found in patients with lung or breast cancer.^{25,26} Recently, however, a serious methodological problem has been identified with serum GST-pi measurements following the observation in healthy subjects that, in the clotting process, large quantities of GST-pi are released from platelets.⁶¹ These findings cast doubt on the results from previous studies that used serum and show that only plasma should be used.

Using plasma obtained under strict sampling conditions, we have shown that plasma GST-pi concentrations are raised in patients with lung cancer, particularly in adenocarcinoma of the lung.⁶¹ We have found that plasma GST-pi is raised in a high proportion of patients with gastrointestinal malignant growths, particularly if there are metastases (unpublished observation).

We conclude that serological measurements of GST-pi show promise as a tumour marker for hepatobiliary and gastrointestinal malignant tumours but earlier studies must be repeated using plasma. It is unlikely, however, that even with strict sampling conditions GST-pi measurements will find a wide role in diagnosis of malignant disease because the ubiquitous nature of the enzyme suggests that it may be raised in a variety of diseases and, indeed, raised concentrations are often found in benign liver disease.^{25,26}

GST MEASUREMENTS IN BILE

Using radioimmunoassay it has been shown that GST-B₁ and GST-pi are present in bile. Increased concentrations of GST-pi have been reported in two patients with cholangiocarcinoma but not in one patient with pancreatic carcinoma.¹¹ Further studies are needed to show whether biliary GST measurements have any value in diagnosing biliary tract cancers.

Conclusion

Study of the distribution of GST isoenzymes in healthy tissue has shown pronounced heterogeneity. The reasons for this are unclear but probably relate to different biological functions of GST isoenzymes in different tissues. Changes in the tissue expression in diseased states may

relate to metabolic changes which occur with disease progression. Overexpression of GST in neoplasia may be causal, allowing replicative advantage, or casual, accompanying clonal expansion.

The use of radioimmunoassay has enabled different GST isoenzymes to be measured in body fluids. Particularly, in situations where acute liver damage occurs, assays for GST-B₁ show high sensitivity compared with transaminase estimation. The major limitation to its widespread use is the length of time needed for performance of the assay and until this is overcome it will remain primarily a research tool.

We thank Mrs Vanessa Campbell for typing this manuscript.

- Mannervik B. The isoenzymes of glutathione transferase. *Adv Enzymol Relat Areas Mol Biol* 1985; 57: 357-417.
- Wolkoff AW. The glutathione S-transferases: their role in the transport of organic anions from blood to bile. *Int Rev Physiol* 1980; 21: 151-69.
- Bach MK, Brashler JR, Morton DR Jr. Solubilization and characterization of the leukotriene C₄ synthetase of rat basophil leukemia cells: a novel particulate glutathione S-transferase. *Arch Biochem Biophys* 1984; 230: 455-65.
- Christ-Hazelhof E, Nugteren DH, Van Dorp DA. Conversions of prostaglandin endoperoxides by glutathione S-transferases and serum albumins. *Biochim Biophys Acta* 1976; 450: 450-61.
- Meyer DJ, Beale D, Tan KH, Coles B, Ketterer B. Glutathione S-transferase in primary hepatomas: the isolation of a form with GSH peroxidase activity. *FEBS Lett* 1985; 184: 139-43.
- Hayes JD, Pickett CB, Mantle TJ, eds. Nomenclatures for GST. In: *Glutathione S-transferases and drug resistance*. London: Taylor and Francis, 1990: xi-iv.
- Morgenstern RF, Butenberg C, De Pierre JW. Microsomal glutathione S-transferase. Purification, initial characterization and demonstration that it is not identical to the cytosolic glutathione S-transferases A, B and C. *Eur J Biochem* 1982; 128: 243-7.
- Morgenstern R, De Pierre JW. Microsomal glutathione transferase. Purification in unactivated form and further characterization of the activation process, substrate specificity and amino acid composition. *Eur J Biochem* 1983; 134: 591-7.
- Hayes PC, Harrison DJ, Bouchier IAD, McLellan LI, Hayes JD. Cytosolic and microsomal glutathione S-transferase isoenzymes in normal human liver and intestinal epithelium. *Gut* 1989; 30: 854-59.
- Hussey AJ, Hayes JD, Beckett GJ. The polymorphic expression of neutral glutathione S-transferase in human mononuclear leucocytes as measured by specific radioimmunoassay. *Biochem Pharm* 1987; 36: 4013-5.
- Howie AF, Hayes PC, Bouchier IAD, Hayes JD, Beckett GJ. Glutathione S-transferase in human bile. *Clin Chim Acta* 1989; 184: 269-78.
- Hayes PC, Aldis P, Portmann B, Hayes JD. Glutathione S-transferase in human liver pathology. In: Hayes JD, Pickett CB, Mantle TJ, eds. *Glutathione S-transferases and carcinogenesis*. London: Taylor and Francis, 1987.
- Batist G, Hudson N, Michaeli Ische K, De Muys JM. Human colon carcinoma has the same biochemical phenotype as resistant carcinogen-induced pre-neoplastic nodules and as human breast cancer cells with multidrug resistance. *Pan American Association for Cancer Research* 1987; 28: 1105.
- Hayes PC, Harrison DJ. Immunohistochemical analysis of pancreas and gastrointestinal tract in man. In: Hayes JD, Pickett CB, Mantle TJ, eds. *Glutathione S-transferases and drug resistance*. London: Taylor and Francis, 1990: 441-50.
- Longnecker DS. Experimental pancreatic carcinogenesis. *Current Opinion in Gastroenterology* 1988; 4: 843-8.
- Pour PM. Mechanism of pseudoductular (tubular) formation during pancreatic carcinogenesis in the hamster model: an electron-microscopic and immunohistochemical study. *Am J Pathol* 1988; 130: 335-44.
- Hayes JD, Pickett CB, Mantle TJ. The glutathione S-transferases and their contribution to drug resistance in nature. In: Hayes JD, Pickett CB, Mantle TJ, eds. *Glutathione S-transferases and drug resistance*. London: Taylor and Francis, 1990: 3-16.
- Mannervik B, Danielson VH. Glutathione-structure and catalytic activity. *Crit Rev Biochem* 1988; 23: 283-337.
- Adachi Y, Horii K, Takahashi Y, Tanihata M, Ohba Y, Yamamoto T. Serum glutathione S-transferase activity in liver diseases. *Clin Chim Acta* 1980; 106: 243-55.
- Singh SV, Leal T, Awasthi Y. Inhibition of human glutathione S-transferase by bile acids. *Toxicol Appl Pharmacol* 1980; 95: 248-54.
- Beckett GJ, Hayes JD. Plasma glutathione S-transferase measurements and liver disease in man. *Journal of Clinical Biochemistry and Nutrition* 1987; 2: 1-24.
- Beckett GJ, Hayes JD. Development of specific radioimmunoassays for the measurement of human hepatic basic and N/A2b glutathione S-transferases. *Clin Chim Acta* 1984; 141: 267-73.
- Hussey AJ, Hayes JD, Beckett GJ. The polymorphic expression of neutral glutathione S-transferase in human mononuclear leucocytes as measured by specific radioimmunoassay. *Biochem Pharmacol* 1987; 36: 4013-5.
- Howie AF, Hayes JD, Beckett GJ. Purification of acidic glutathione S-transferases from human lung, placenta and erythrocyte and the development of a specific radioimmunoassay for their measurement. *Clin Chim Acta* 1988; 177: 65-86.
- Niitsu Y, Takahashi Y, Saito T, et al. Serum glutathione S-transferase as a tumor marker for gastrointestinal malignancies. *Cancer* 1989; 63: 317-23.
- Tsuchida S, Sekine Y, Shintcha R, Nishihira T, Sato K. Elevation of the placental glutathione S-transferase form (GST π) in tumor tissues and the levels in sera of patients with cancer. *Cancer Res* 1989; 49: 5225-9.
- Stockman PK, Beckett GJ, Hayes JD. Identification of a hybrid glutathione S-transferase from human liver. *Biochem J* 1985; 227: 457-65.
- Beckett GJ, Howie AF, Hussey AJ, Hayes PC, Miller WR, Hayes JD. Radioimmunoassay measurements of the human glutathione S-transferases. In: Hayes JD, Hayes PC, Mantle TJ, Pickett CB, eds. *Glutathione S-transferase and drug resistance*. London: Taylor and Francis, 1990: 399-409.
- Hiley C, Fryer A, Bell J, Hume R, Strange RC. The human glutathione S-transferases. Immunohistochemical studies of the developmental expression of alpha- and pi-class isoenzymes in liver. *Biochem J* 1989; 254: 255-9.
- Beckett GJ, Chapman BJ, Dyson EH, Hayes JD. Plasma glutathione S-transferase measurements following paracetamol overdose: evidence for early hepatocellular damage. *Gut* 1985; 26: 26-31.
- Sherman M, Bass NM, Campbell JAH, Kirsch RE. Radioimmunoassay of human ligandin. *Hepatology* 1983; 3: 162-9.
- Howie AF, Forrester LM, Glancy MJ et al. Glutathione S-transferase and glutathione peroxidase in human tumors. *Carcinogenesis* 1990; 11: 451-8.
- Strange RC, Fryer AA, Hiley C, et al. Developmental expression of GST in human tissues. In: Hayes JD, Hayes PC, Mantle TJ, Pickett CB, eds. *Glutathione S-transferase and drug resistance*. London: Taylor and Francis, 1990: 262-72.
- Harrison DJ, Hayes PC. Immunolocalisation of glutathione S-transferases in human renal and liver diseases. In: Hayes JD, Hayes PC, Mantle TJ, Pickett CB, eds. *Glutathione S-transferase and drug resistance*. London: Taylor and Francis, 1990: 431-41.
- Beckett GJ, Foster GR, Hussey AJ, et al. Plasma glutathione S-transferase and F protein are more sensitive than alanine aminotransferase as indicators of paracetamol (acetaminophen)-induced liver damage. *Clin Chem* 1989; 35: 2186-9.
- Beckett GJ, Hussey AJ, Laing I, et al. Measurement of glutathione S-transferase B₁ in plasma after birth asphyxia: an early indication of hepatocellular damage. *Clin Chem* 1989; 35: 995-9.
- Hayes PC, Hayes JD, Hussey AJ, Bouchier IAD, Beckett GJ. Changes in plasma glutathione S-transferase B₁ concentration after alcohol ingestion in man: a measure of hepatocellular sensitivity to chronic alcohol excess. *Clinical Chemistry and Enzymology Communications* 1990; 2: 189-94.
- Neuberger J, Williams R. Halothane anaesthesia and liver damage. *BMJ* 1984; 289: 1136-9.
- Farrel G, Prendergast D, Murray M. Halothane hepatitis: detection of a constitutional susceptibility factor. *N Engl J Med* 1985; 313: 1310-4.
- Brown BR, Gandolfi AJ. Adverse effects of volatile anaesthetics. *Br J Anaesth* 1987; 59: 14-23.
- Hussey AJ, Howie JJ, Allan LG, Drummond H, Hayes JD, Beckett GJ. Impaired hepatocellular integrity during general anaesthesia as assessed by measurement of plasma glutathione S-transferase. *Clin Chim Acta* 1986; 161: 19-28.
- Hussey AJ, Aldridge LM, Paul D, Ray DC, Beckett GJ, Allan LG. Plasma glutathione S-transferase levels as a measure of hepatocellular integrity following a single general anaesthetic with either halothane, enflurane or isoflurane. *Br J Anaesth* 1988; 60: 130-5.
- Hayes PC, Hussey AF, Keating J, et al. Glutathione S-transferase levels in autoimmune chronic active hepatitis: a more sensitive index of hepatocellular damage than aspartate aminotransferase. *Clin Chim Acta* 1988; 172: 211-6.
- Beckett GJ, Hayes PC, Hussey AJ, Bouchier IAD, Hayes JD. Plasma glutathione S-transferase measurements in patients with alcoholic cirrhosis. *Clin Chim Acta* 1987; 169: 85-90.
- Beckett GJ, Kellett HA, Gow SM, Hussey AJ, Hayes JD, Toft AD. Raised plasma glutathione S-transferase values in hyperthyroidism and in hypothyroid patients receiving thyroxine replacement: evidence for hepatic damage. *BMJ* 1985; 291: 427-31.
- Sheridan P. Thyroid hormones and the liver. *Clinics in Gastroenterology* 1983; 12: 797-818.
- Gow SM, Caldwell G, Toft AD, et al. Relationship between pituitary and other target organ responsiveness in hypothyroid patients receiving thyroxine replacement. *J Clin Endocrinol Metab* 1987; 64: 364-70.
- Toft AD. Thyroxine replacement: clinical judgment or biochemical control. *BMJ* 1985; 291: 233-4.
- Gow SM, Caldwell G, Toft AD, Sweeting VM, Beckett GJ. Restoration of normal thyrotrophin secretion reduces the abnormally high glutathione S-transferase levels found in patients receiving thyroxine replacement therapy. *Clin Endocrinol* 1988; 29: 249-56.

- 50 Gow SM, Caldwell G, Toft AD, Beckett GJ. Different hepatic responses to thyroxine replacement in spontaneous and ¹³¹I-induced primary hypothyroidism. *Clin Endocrinol* 1989; **30**: 505–12.
- 51 Wood M, Berman ML, Harbison RD. Halothane-induced hepatic necrosis in triiodothyronine-treated rats. *Anesthesiology* 1980; **50**: 470–6.
- 52 Calvert DW, Brody TM. The effects of thyroid function upon carbon tetrachloride hepatotoxicity. *J Pharmacol Exp Ther* 1961; **134**: 304–10.
- 53 Beckett GJ, Boyd R, Beddows SE, Hayes JD. Decreased hepatic glutathione S-transferases A, AA and L concentrations produced by prolonged thyroid hormone administration. *Br J Pharmacol* 1988; **37**: 3201–4.
- 54 Board PG. Biochemical genetics of glutathione S-transferase in man. *Am J Hum Genet* 1981; **33**: 36–43.
- 55 Seidegard J, Pero RW, Markowitz MM, *et al.* Isoenzymes of glutathione S-transferase (class Mu) as a marker for the susceptibility to lung cancer: a follow-up study. *Carcinogenesis* 1990; **11**: 33–6.
- 56 Harada S, Abei M, Tanaka N, Agarwai DP, Goedde HW. Liver glutathione S-transferase polymorphism in Japanese and its pharmacogenetic importance. *Hum Genet* 1987; **75**: 322–5.
- 57 Kodate C, Fukushi A, Narita T, Kudo H, Soma Y, Sato K. Human placental form of glutathione S-transferase (GST) as a new immunohistochemical marker for human colonic cancer. *Gann* 1986; **77**: 226–9.
- 58 Di Illio C, Del Boccio G, Aceto A, *et al.* Elevation of glutathione S-transferase activity in human lung tumor. *Carcinogenesis* 1988; **9**: 335–40.
- 59 Eimoto H, Tsutsumi M, Nakajima A. Expression of the glutathione S-transferase placental form in human lung carcinomas. *Carcinogenesis* 1988; **9**: 2325–7.
- 60 Howie AF, Miller WR, Hawkins RA, Hutchison AR, Beckett GJ. Expression of glutathione S-transferase B₁, B₂, mu and pi in breast cancers and their relationship to oestrogen receptor status. *Br J Cancer* 1990; **60**: 834–7.
- 61 Howie AF, Douglas JG, Ferguson RJ, Beckett GJ. Plasma glutathione S-transferase pi measurements: a possible marker for adenocarcinoma of the lung. *Clin Chem* 1990; **36**: 453–6.