

Leading article

Sulphate reducing bacteria and hydrogen metabolism in the human large intestine


One of the main functions of the large intestine is to salvage energy from dietary carbohydrate which has neither been digested nor absorbed in the small intestine. This occurs through a process known as fermentation in which anaerobic bacteria breakdown carbohydrate to short chain fatty acids (SCFA), which in turn provide energy for the host. The major carbohydrates available for fermentation are resistant starches, non-starch polysaccharides (dietary fibre), and a variety of unabsorbed sugars and oligosaccharides. Other substances can also be fermented, including dietary proteins and host derived substrates such as pancreatic enzymes, mucus, and sloughed epithelial cells.¹

The fermentation of organic matter by gut microorganisms is mediated by a number of interdependent reactions in which complex polymers are first broken down to their constituent monomers by hydrolytic enzymes synthesised by the bacteria. They are subsequently oxidised to SCFA, lactate, succinate, ethanol, and the gases H₂ and CO₂.¹ In some people, methane is also produced.² Fermentation is regulated by the amount of substrate available and its chemical composition, the substrate specificities and preferences of the bacterial species present and the metabolic pathways through which they metabolise the substrate.³ Host factors such as transit time may also play a part.


Hydrogen is an important product of fermentation and is primarily formed by bacteria as a 'sink' for the disposal of reducing power (electrons) generated during oxidation of sugars and amino acids. In mammalian cells, oxygen is used as the terminal electron acceptor and reducing power is disposed of in the form of H₂O, this is not possible, however, in anaerobic metabolism. For example, some bacteria form ethanol, lactate, or succinate and do not evolve much H₂ because they dispose of reducing energy in these electron sink products.⁴

Hydrogen formation results principally from the oxidation of pyruvate, formate, or reduced pyridine nucleotides (NADH⁺, FADH⁺). Moreover, many clostridia generate H₂ from pyruvate via ferredoxin, whereas enterobacteria evolve the gas through cleavage of pyruvate by pyruvate formate lyase. The formate produced is then converted to CO₂ and H₂. The partial pressure of H₂ has no effect on its production from pyruvate, but accumulation of the gas inhibits formation from the oxidation of pyridine nucleotides. Since the regeneration of oxidised pyridine nucleotides through H₂ production enables increased substrate level phosphorylation to occur, an efficient H₂ removal may be a contributing factor in maintaining the fermentation balance in the colon. Hydrogen partial pressures in the colon are kept low by losses in breath and flatus^{5–7} and by the activities of hydrogen utilising species such as methanogenic,⁸ aceto-genic,⁹ and sulphate reducing bacteria.¹⁰

Between 30 and 50% of people in western countries harbour methanogenic bacteria in their colons.² These bacteria grow by reducing CO₂ and H₂ according to the following equation¹¹:



Thus, 4 mol of H₂ are converted to 1 mol of CH₄. Methane production is an effective and safe pathway for H₂ disposal and occurs in most animals. In people who do not excrete CH₄, other routes for microbial H₂ disposal frequently exist. An important alternative is the reduction of sulphate by sulphate reducing bacteria.¹² We have shown that sulphate reducing bacteria occur in large numbers (ca 10⁸–10¹⁰ (g dry weight gut contents)¹) in non-methanogenic but not in methanogenic individuals.¹³ Sulphate reducing bacteria utilise H₂ according to the following equation¹¹:



As in methanogenesis, 4 mol of H₂ are consumed in the formation of 1 mol of product. Unlike methanogenesis, however, the product is not a harmless gas, but is highly toxic hydrogen sulphide, an agent that is potentially damaging to the colonic epithelium.¹³ People who harbour sulphate reducing bacteria in their large bowel have higher levels of sulphide in their faeces than methanogenic subjects.¹² The main substrates for colonic sulphate reducing bacteria are fermentation products formed by other bacteria, such as acetate, propionate, lactate, butyrate, succinate, ethanol, pyruvate, some amino acids, and H₂/CO₂.¹⁴ Species that can utilise long chain fatty acids and alcohols (up to C6) have also been isolated by chemostat enrichment.¹⁵ These strict anaerobes are the final link in the food chain that develops in the gut ecosystem and use sulphate as a terminal electron acceptor during oxidative reactions.¹⁶ The predominant sulphate reducing bacteria found in the gut belong to five genera, namely *Desulfovibrio*, *Desulfobacter*, *Desulfomonas*, *Desulfobulbus* and *Desulfotomaculum*.¹⁷ Species that use molecular H₂ as an electron donor in the gut belong to the genera *Desulfovibrio* and *Desulfobulbus*. *Desulfovibrios* account for approximately 66% of all colonic sulphate reducing bacteria, while *Desulfobulbus* spp. constitute about 16%.¹⁷

Fermentation experiments in our laboratory have shown that colonic sulphate reducing bacteria outcompete methanogenic bacteria for H₂.¹⁸ More recently, however, Strocchi *et al*¹⁹ have postulated that methanogens were able to competitively displace other H₂ consuming bacteria in mixed faecal slurries. In this study, sulphate reducing bacteria must have been absent, or their activities limited in the non-methanogenic samples tested, because it is highly unlikely that methanogenic bacteria will displace sulphate reducing bacteria for this mutual growth substrate given an adequate supply of electron acceptor (sulphate). The physiological explanation for this resides in the greater affinity of sulphate reducing bacteria for H₂ compared with methanogenic bacteria (Ks of *Desulfovibrio vulgaris*, 1 μmol.l⁻¹; Ks of *Methanobrevibacter smithii*, 6 μmol.l⁻¹).²⁰ Moreover, the oxidation of H₂ by sulphate reducing bacteria is thermodynamically more favourable (ΔG₀' = -152.2 kJ per mol) than by methanogenic bacteria (ΔG₀' = -131 kJ per mol).¹¹ The outcome of competition is largely dependent on sulphate availability, with sulphate reducing bacteria outcompeting methanogenic bacteria for the mutual growth substrate H₂

in the presence of sulphate.²¹⁻³¹ Strocchi *et al*¹⁹ did not enumerate viable sulphate reducing bacteria in their study, and the addition of 20 mM sulphate to non-methanogenic faecal slurries made little difference to sulphide production, indicating an absence of viable sulphate reducers. In contrast, we have found that in two geographically and nutritionally diverse populations, viable sulphate reducing bacteria were present and active in all non-methanogenic samples tested.¹⁴

Thus, if sufficient sulphate is available *in vivo*, our belief is that sulphate reducing bacteria will be responsible for most of the H₂ utilised by the gut microflora. When sulphate availability is limited, methanogenic bacteria may then predominate. Other lines of evidence support this hypothesis. For example, if sulphate is added to the diet of methanogenic individuals, competitive displacement of methanogenic bacteria occurs and hitherto undetectable sulphate reducing bacteria begin to appear in faeces within a few days.³² This effect is observed only in about 50% of methanogenic individuals, however, suggesting that sulphate reducing bacteria and methanogenic bacteria coexist in some, but not all, people. Furthermore, in whole body calorimetry studies when total H₂ and CH₄ excretion were measured in healthy subjects eating test meals of various fermentable carbohydrates, the amount of excreted H₂ that could be accounted for was much less in non-methanogenic subjects. For example, in response to a 15 g dose of lactulose, methanogenic subjects excreted mean (SEM) 1150 (212) ml hydrogen equivalents, while non-methanogenic volunteers excreted only 327 (88) ml.³³ These data support the existence of alternative pathways for H₂ disposal in man.

For sulphate reducing bacteria to outcompete methanogenic bacteria for H₂ in the colon, there must be an adequate supply of sulphate. Studies of sulphate absorption using ileostomists have indicated that concentrations of this anion can vary considerably in diet and that amounts reaching the colon may range from 2 to 9 mmol.day⁻¹.³⁴ Other potential sources exist, including highly sulphated polysaccharides such as chondroitin sulphate.^{35,36} Colonic mucins are also highly sulphated, more so in fact than mucins produced elsewhere in the gastrointestinal tract.^{37,38} There are many species of bacteria in the colon including clostridia, bifidobacteria, and bacteria of the *Bacteroides fragilis* group that are able to degrade these substances and thereby release free sulphate³⁹⁻⁴⁵ which would then become available for utilisation by sulphate reducing bacteria. If this is indeed the case, then the amount of sulphated glycoprotein produced by the host, its degree of sulphation, and the activities of the hydrolytic bacteria involved in releasing free sulphate from these polymers, would all influence sulphate reducing bacterial growth. Because mucin structure and production varies between individuals, there may possibly be a genetic predisposition to the acquisition of sulphate reducing bacteria.

Another pathway of H₂ disposal is by the activities of acetogenic bacteria, which also grow in the colon. Species that reduce CO₂ with H₂ do so according to the following equation¹¹:



Acetate production from CO₂ occurs in the rat caecum⁴⁶ and in the gut of termites.⁴⁷ Significant acetate production as a result of H₂ dependent CO₂ reduction has also been reported in human faeces in the absence of methanogenesis.⁴⁸ In terms of energy, however, H₂/CO₂ acetogenesis, with a free energy change of -95 kJ per mol,¹¹ is considerably less efficient than either dissimilatory sulphate reduction or methanogenesis, and in most anaerobic ecosystems, hydrogen metabolism is dominated by one or other of these processes.^{49,50} However, pH can influence the outcome of

competition, with acidic conditions selecting for acetogenesis,¹² and it is possible therefore that the luminal pH in the colon is important in this respect. Nevertheless, it is unlikely that homoacetogenesis is a major route of disposal of H₂ in the human colon since molar ratios of acetate, propionate, and butyrate are similar in methanogenic and non-methanogenic people.⁵¹ Comparison of SCFA concentrations in the gut of known acetogenic animals, such as the termite, with man, shows that in the insect acetate accounts for 94% of the total compared with only 57% in the human caecum.

What are the clinical implications of all this? Effective H₂ disposal during fermentation is essential for normal large bowel function. In man, the principal routes are via methanogenesis or dissimilatory sulphate reduction. Methanogenesis is a safe and effective disposal route for H₂. Methane is non-toxic and allows a large volume of H₂ to be removed, thereby potentially reducing any gas related symptoms. In the absence of either methanogenic bacteria or sulphate reducing bacteria in the colon, H₂ accumulates in excessive quantities and this may be an important factor in pneumatosis cystoides intestinalis.⁵² Other ways of disposing of H₂ include acetogenesis and dissimilatory nitrate reduction, for which H₂ is the preferred electron donor. Acetogenesis is probably the most favourable H₂ disposal method, since this makes additional energy available to the host from fermentation through acetate absorption. However, the evidence for significant homoacetogenesis in man is still inconclusive.

Sulphate reduction is probably the least desirable route. The main product, sulphide, is toxic to cells and impairs cellular metabolism through inhibition of cytochrome oxidase.^{53,54} Sulphide may also destroy the disulphide bridges in mucus, in turn leading to a breakdown of the protective layer of the epithelium. Moreover, other sulphhydryl compounds have been implicated in colonic disease since Roediger *et al*^{55,56} have shown that mercaptoacetate and mercaptobutyrate affect fatty acid metabolism in colonocytes in a manner that is characteristic of the defect found in ulcerative colitis. We have shown that ulcerative colitis patients have a very high carriage rate of sulphate reducing bacteria and raised sulphide values in faeces.^{57,58}

Thus the removal of H₂ in the large gut may follow a number of pathways. It is clear from studies in man that either methanogenesis or dissimilatory sulphate reduction predominate in some populations. Because of the known toxicity of sulphide in the gut we suggest that the role of sulphate reducing bacteria in the pathogenesis of colonic disease warrants further investigation.

G R GIBSON
G T MACFARLANE
J H CUMMINGS

MRC Dunn Clinical Nutrition Centre,
100 Tennis Court Road,
Cambridge CB2 1QL

- Cummings JH, Macfarlane GT. A review: The control and consequences of bacterial fermentation in the human colon. *J Appl Bacteriol* 1991; 70: 443-59.
- Bond JH, Engel RR, Levitt MD. Factors influencing pulmonary excretion in man. *J Exp Med* 1971; 133: 572-8.
- Macfarlane GT. Fermentation reactions in the large intestine. In: Cummings JH, Rombeau JL, Sakata T. *Short chain fatty acids: Metabolism and clinical importance*. Columbus: Ross Laboratories Press, 1991; 5-10.
- Wolin MJ. Interactions between H₂-producing and methane producing species. In: Schlegel HG, Gottschalk G, Pfennig N, eds. *Microbial formation and utilization of gases*. Göttingen: Goltze Press, 1976: 141-50.
- Bond JH, Levitt MD. Use of pulmonary hydrogen measurements to quantitate carbohydrate absorption. *J Clin Invest* 1972; 51: 1219-25.
- Anderson IH, Levine AS, Levitt MD. Incomplete absorption of the carbohydrates in all-purpose wheat flour. *N Engl J Med* 1981; 304: 891-2.
- Levitt MD, Hirsch P, Fetzter CA, Sheahan M, Levine A. H₂ excretion after ingestion of complex carbohydrates. *Gastroenterology* 1987; 92: 383-9.
- Zehnder AJB. Ecology of methane formation. In: Mitchell R, ed. *Water pollution microbiology*. Vol 2. New York: John Wiley, 1978: 349-76.
- Ljungdahl LG. The autotrophic pathway of acetate synthesis in acetogenic bacteria. *Ann Rev Microbiol* 1986; 40: 415-50.
- Postgate JR. *The sulphate-reducing bacteria*. 2nd ed. Cambridge: Cambridge University Press, 1984.

- 11 Thauer RK, Jungermann K, Decker K. Energy conservation in chemotrophic anaerobic bacteria. *Bacteriol Rev* 1977; **41**: 100-80.
- 12 Gibson GR, Cummings JH, Macfarlane GT, Allison C, Segal I, Vorster HH, et al. Alternative pathways for hydrogen disposal during fermentation in the human colon. *Gut* 1990; **31**: 679-83.
- 13 Karrer P. *Organic chemistry*. Amsterdam: Elsevier, 1960.
- 14 Gibson GR, Macfarlane GT, Cummings JH. Occurrence of sulphate-reducing bacteria in human faeces and the relationship of dissimilatory sulphate reduction to methanogenesis in the large gut. *J Appl Bacteriol* 1988; **65**: 103-11.
- 15 Gibson GR, Macfarlane GT. Chemostat enrichment of sulphate-reducing bacteria from the large gut. *Lett Appl Microbiol* 1988; **7**: 127-33.
- 16 Gibson GR. A review: physiology and ecology of the sulphate-reducing bacteria. *J Appl Bacteriol* 1990; **69**: 769-97.
- 17 Gibson GR, Cummings JH, Macfarlane GT. Factors affecting hydrogen uptake by bacteria growing in the human large intestine. In: Belaich JP, Bruschi M, Garcia JL, eds. *Microbiology and biochemistry of strict anaerobes involved in interspecies hydrogen transfer*. New York: Plenum Press, 1990: 191-202.
- 18 Gibson GR, Cummings JH, Macfarlane GT. Competition for hydrogen between sulphate-reducing bacteria and methanogenic bacteria from the human large intestine. *J Appl Bacteriol* 1988; **65**: 241-7.
- 19 Stocchi A, Furne JK, Ellis CJ, Levitt MD. Competition for hydrogen by human faecal bacteria: evidence for the predominance of methane producing bacteria. *Gut* 1991; **32**: 1498-501.
- 20 Kristjansson JK, Schönheit P, Thauer RK. Different Ks values for hydrogen of methanogenic bacteria and sulfate reducing bacteria: an explanation for the apparent inhibition of methanogenesis by sulfate. *Arch Microbiol* 1982; **131**: 278-82.
- 21 Lovley DR, Klug MJ. Sulfate-reducers can outcompete methanogens at freshwater sulfate concentrations. *Appl Environ Microbiol* 1983; **45**: 187-92.
- 22 Lovley DR, Dwyer DF, Klug MJ. Kinetic analysis of competition between sulfate reducers and methanogens for hydrogen in sediments. *Appl Environ Microbiol* 1982; **43**: 1373-9.
- 23 Abram JW, Nedwell DB. Inhibition of methanogenesis by sulphate reducing bacteria competing for transferred hydrogen. *Arch Microbiol* 1978; **117**: 89-92.
- 24 Abram JW, Nedwell DB. Hydrogen as a substrate for methanogenesis and sulphate reduction in anaerobic saltmarsh sediment. *Arch Microbiol* 1978; **117**: 93-7.
- 25 King GM, Wiebe WJ. Tracer analysis of methanogenesis in salt marsh soil. *Appl Environ Microbiol* 1980; **39**: 877-81.
- 26 Martens CS, Berner RA. Methane production in the interstitial waters of sulfate-depleted sediments. *Science* 1974; **185**: 1167-9.
- 27 Mountford DD, Asher RA. Role of sulfate reduction versus methanogenesis in terminal carbon flow in polluted intertidal sediments of Waimea Inlet, Nelson, New Zealand. *Appl Environ Microbiol* 1981; **42**: 252-8.
- 28 Mountford DD, Asher RA, Mays EL, Tiedje JM. Carbon and electron flow in mud and sand intertidal sediments at Delaware Inlet, Nelson, New Zealand. *Appl Environ Microbiol* 1980; **39**: 686-94.
- 29 Winfrey MR, Zeikus JG. Effect of sulfate on carbon and electron flow during microbial methanogenesis in freshwater sediments. *Appl Environ Microbiol* 1977; **33**: 275-81.
- 30 Oremland RS, Taylor BF. Sulfate reduction and methanogenesis in marine sediments. *Geochimica Cosmochimica Acta* 1978; **42**: 209-14.
- 31 Oremland RS, Polcin S. Methanogenesis and sulfate reduction: competitive and non-competitive substrates in estuarine sediments. *Appl Environ Microbiol* 1982; **44**: 1270-6.
- 32 Christl SU, Gibson GR, Florin THJ, Cummings JH. The role of dietary sulfate in the regulation of methanogenesis in the human large intestine. *Gastroenterology* 1990; **98**: A164.
- 33 Christl SU, Murgatroyd PR, Gibson GR, Cummings JH. Production, excretion and metabolism of hydrogen in the large intestine. *Gastroenterology* 1990; **102**: 1269-77.
- 34 Florin THJ, Neale G, Gibson GR, Christl SU, Cummings JH. Metabolism of dietary sulphate: absorption and excretion in humans. *Gut* 1991; **32**: 766-73.
- 35 Salyers AA, O'Brien M. Cellular location of enzymes involved in chondroitin sulfate breakdown by *Bacteroides thetaiotaomicron*. *J Bacteriol* 1980; **143**: 772-80.
- 36 Ishioka T, Kuwabara N, Oohashi Y, Wakabayashi K. Induction of colorectal tumours in rats by sulfated polysaccharides. *CRC Crit Rev Toxicol* 1987; **17**: 215-44.
- 37 Jennings MA, Florey H. Autoradiographic observations on the mucous cells of the stomach and intestine. *Q J Exp Physiol* 1956; **41**: 131-51.
- 38 Filipe MI. Mucins in the human gastroepithelium: a review. *Invest Cell Pathol* 1979; **2**: 195-216.
- 39 Dogson KS, White GF, Fitzgerald JW. *Sulfatases of microbial origin*. Vol 1. Boca Raton: CRC Press, 1982.
- 40 Macfarlane GT, Gibson GR. Formation of glycoprotein degrading enzymes by *Bacteroides fragilis*. *FEMS Microbiol Ecol* 1991; **77**: 289-94.
- 41 Gibson GR, Cummings JH, Macfarlane GT. Use of a three-stage continuous culture system to study the effect of mucin on dissimilatory sulfate reduction and methanogenesis by mixed populations of human gut bacteria. *Appl Environ Microbiol* 1988; **54**: 2750-5.
- 42 Macfarlane GT, Hay S, Gibson GR. Influence of mucin on glycosidase, protease and arylamidase activities of human gut bacteria grown in a 3-stage continuous culture system. *J Appl Bacteriol* 1989; **66**: 407-17.
- 43 Tsai HH, Hart CA, Rhodes JM. Production of mucin degrading sulphatase and glycosidases by *Bacteroides thetaiotaomicron*. *Lett Appl Microbiol* 1991; **13**: 97-101.
- 44 Robertson AM, Stanley RA. *In vitro* utilization of mucin by *Bacteroides fragilis*. *Appl Environ Microbiol* 1982; **43**: 325-30.
- 45 Vercellotti JR, Salyers AA, Bullard WS, Wilkins TD. Breakdown of mucin and plant polysaccharides in the human colon. *Can J Biochem* 1977; **55**: 1190-6.
- 46 Prins RA, Lankhorst A. Synthesis of acetate from CO₂ in the cecum of some rodents. *FEMS Microbiol Letts* 1977; **1**: 255-8.
- 47 Breznak JA, Switzer JM. Acetate synthesis from H₂ plus CO₂ by termite microbes. *Appl Environ Microbiol* 1986; **52**: 623-30.
- 48 Lajoie SF, Bank S, Miller TL, Wolin MJ. Acetate production from hydrogen and [¹³C] carbon dioxide by the microflora of human feces. *Appl Environ Microbiol* 1988; **54**: 2723-7.
- 49 Coutts DAP, Balba MTM, Senior E. Acetogenesis and acetotrophy in a hexanoate-catabolizing microbial association isolated from landfill. *J Appl Bacteriol* 1987; **63**: 343-53.
- 50 Zeikus JG. Microbial populations in digestors. In: Stafford DA, Wheatley BI, Hughes DE, eds. *Proceedings of the 1st International Symposium on Anaerobic Digestion*. London: Applied Science, 1980: 73-103.
- 51 Macfarlane GT, Gibson GR, Cummings JH. Comparison of fermentation reactions in different regions of the human colon. *J Appl Bacteriol* 1972; **57**: 57-64.
- 52 Christl SU, Gibson GR, Murgatroyd PR, Scheppach W, Cummings JH. Impaired H₂ metabolism in pneumatosis cystoides intestinalis. *Gastroenterology* 1991; **100**: A203.
- 53 Smith L, Kruszyna H, Smith RP. The effect of methaemoglobin on the inhibition of cytochrome c oxidase by cyanide, sulfide or azide. *Biochem Pharmacol* 1977; **22**: 47-50.
- 54 Florin THJ, Neale G, Cummings JH. The effect of hydrogen sulphide on isolated rat colonic epithelium. *Falk Symposium on Advances in the Treatment of Ulcerative Colitis* 1990 [Abstract].
- 55 Roediger WEW, Duncan A, Nance SH. Bacterial mercaptoacetate inhibits lipid synthesis in colonocytes - a factor in aetiology of ulcerative colitis? *Aust NZ J Med* 1990; **20**: 377.
- 56 Roediger WEW, Nance S. Selective reduction of fatty acid oxidation in colonocytes: correlation with ulcerative colitis. *Lipids* 1990; **25**: 646-52.
- 57 Florin THJ, Gibson GR, Neale G, Cummings JH. A role for sulfate reducing bacteria in ulcerative colitis? *Gastroenterology* 1990; **98**: A170.
- 58 Gibson GR, Cummings JH, Macfarlane GT. Growth and activities of sulphate-reducing bacteria in gut contents of health subjects and patients with ulcerative colitis. *FEMS Microbiol Ecol* 1991; **86**: 103-12.