

Effect of acute hyperglycaemia on gall bladder contraction induced by cholecystokinin in humans

S Y de Boer, A A M Masclee, M C W Jebbink, J Schipper, H H P J Lemkes, J B M J Jansen, C B H W Lamers

Abstract

This study examined the effect of acute hyperglycaemia, induced by intravenous glucose, on gall bladder motility. Six healthy volunteers were studied in random order on three occasions during normoglycaemia and hyperglycaemia with blood glucose concentrations stabilised at 8 and 15 mmol/l. Gall bladder volumes, measured with ultrasonography, were studied before and during infusion of stepwise increasing doses of cholecystokinin (CCK-33; 0.25, 0.5, and 1.0 IDU.kg⁻¹.h⁻¹). Each dose was given for 30 minutes. Pancreatic polypeptide (PP) secretion was determined as an indirect measure of cholinergic tone. Infusion of CCK-33 resulted in significant dose dependent reductions in gall bladder volume in all three experiments. Compared with normoglycaemia the gall bladder contraction was significantly ($p < 0.05$) reduced during infusion of 0.25 and 0.5 IDU.kg⁻¹.h⁻¹ CCK-33 in the 8 mmol/l hyperglycaemic experiment, and during infusion of 0.25, 0.5, and 1.0 IDU.kg⁻¹.h⁻¹ CCK-33 in the 15 mmol hyperglycaemic experiment. During hyperglycaemia basal plasma PP concentrations and PP secretion in response to CCK-33 were significantly ($p < 0.05$) reduced. It is concluded that blood glucose concentrations affect gall bladder motility, that an acute hyperglycaemia at 8 and 15 mmol/l reduces the gall bladder responsiveness to CCK-33 in a dose dependent manner, and that hyperglycaemia reduces basal and CCK-33 stimulated plasma PP concentrations, suggesting impaired cholinergic activity during hyperglycaemia.

(Gut 1993; 34: 1128-1132)

Recently several studies have emphasised the importance of serum glucose concentrations in the regulation of gastrointestinal function.¹⁻⁶ In healthy subjects hyperglycaemia induced by intravenous infusion of glucose reduces gastric emptying of solid and liquid meals,¹⁻³ gastric acid secretion,^{4,5} and pancreatic secretion.^{4,6} Little is known, however, about the effect of hyperglycaemia on gall bladder motility. It has been suggested that hyperglycaemia reduces gastrointestinal function through inhibition of the vagal cholinergic system.^{2,3}

This study was undertaken to investigate the influence of hyperglycaemia on gall bladder motility. Gall bladder contraction in response to infusion of graded doses of cholecystokinin (CCK-33) was measured during normoglycaemia and during acute hyperglycaemia with blood glucose concentrations stabilised at 8 and 15 mmol/l. A second aim of the study was to

examine the tone of the cholinergic system during hyperglycaemia. This was accomplished indirectly by measuring pancreatic polypeptide (PP) secretion, which is known to be dependent on cholinergic tone and reduced during hyperglycaemia in healthy subjects.^{7,8}

Materials and methods

SUBJECTS

Six healthy young men (aged 24-32) participated in the study. None was taking chronic medication or had a history of gastrointestinal disease, surgery, or diabetes mellitus. Informed consent was obtained from each person and the protocol was approved by the ethics committee of Leiden University Hospital.

EQUIPMENT

Gall bladder volumes, measured by real time ultrasonography (Technicare, 3.5 MHz transducer), were calculated by the sum of cylinders method with a computerised system.^{9,10} In this method the longitudinal scan of the gall bladder is divided into series of cylinders of equal height, with diameters perpendicular to the longitudinal axis of the gall bladder image. The uncorrected volume is the sum of volumes of these separate cylinders. To correct for the displacement of the longitudinal image of the gall bladder from the central axis, a correction factor is calculated from the longitudinal and transversal scans of the gall bladder. Gall bladder volume is calculated by multiplication of the uncorrected volume by the square of the correction factor. The mean of two measurements was used for further analysis. The assumptions and the mathematical formula used to calculate gall bladder volume have been described and validated previously by Everson *et al* and Hopman *et al*.^{9,10}

TEST PROCEDURE

Each subject participated in three experiments performed in random order, with an interval of at least seven days between each. In the three experiments the subjects were either normoglycaemic or hyperglycaemic with blood glucose concentrations stabilised at 8 or 15 mmol/l. All experiments were started at 0900 after an overnight fast. An intravenous cannula was inserted into an antecubital vein of one arm for blood sampling. A second intravenous line was positioned in the antecubital vein of the other arm for infusions of 20% glucose and 0.9% saline. An acute hyperglycaemia at 8 or 15 mmol/l was obtained with a modified glucose clamp tech-

Department of
Gastroenterology-
Hepatology,
S Y de Boer
A A M Masclee
M C W Jebbink
J B M J Jansen
C B H W Lamers

Department of
Radiology,
J Schipper

Department of
Endocrinology,
University Hospital
Leiden, The Netherlands
H H P J Lemkes

Correspondence to:
A A M Masclee, Department
of Gastroenterology-
Hepatology, University
Hospital, Building 1, C4-P,
PO Box 9600, 2300 RC
Leiden, The Netherlands.

Accepted for publication
8 December 1992




Figure 1: Serum glucose concentrations (mmol/l, mean (SEM)) in six healthy subjects during normoglycaemia and hyperglycaemic clamping.

nique.^{11,12} The acute rise in blood glucose concentrations was induced by an intravenous bolus injection of a glucose 20% solution 45 minutes before the start of CCK-33 infusion. The amount of glucose given was calculated from body weight. After a bolus injection the rate of glucose infusion was adjusted to maintain a blood glucose concentration of about 8 or 15 mmol/l according to the protocol. Serum glucose concentrations were measured every 2.5 to five minutes by the glucose oxidase method (Beckman glucose analyser, Beckman Instruments).

At zero minutes a continuous intravenous infusion of CCK-33 (Ferring, Malmo, Sweden) was started. Cholecystokinin was given in stepwise increasing doses of 0.25 IDU kg⁻¹.h⁻¹, 0.5 IDU kg⁻¹.h⁻¹, and 1.0 IDU kg⁻¹.h⁻¹. Each dose was for 30 minutes. Previous studies have shown that during infusions of 0.25, 0.5, and 1.0 IDU kg⁻¹.h⁻¹ plasma cholecystokinin (CCK) concentrations correspond with plasma CCK




Figure 2: Plasma insulin concentrations (mU/l, mean (SEM)) during normoglycaemia and hyperglycaemic clamping.

concentrations measured after ingestion of a light, normal, and fatty meal.¹³ Gall bladder images were taken in duplicate every 15 minutes at -60, -45, -30, -15, 0, 15, 30, 45, 60, 75, and 90 minutes. Blood samples for measurement of PP, CCK, and insulin were drawn each time a gall bladder image was taken. Plasma PP concentrations were measured by a sensitive and specific radioimmunoassay as described previously.¹⁴ Plasma CCK was measured by a sensitive and specific radioimmunoassay with antibody T204.^{15,16} This antibody binds to all carboxy terminal CCK peptides containing the sulphated tryosyl region. The detection limit of the assay was 0.5 pmol/l plasma. The within assay variation ranged from 4.6% to 11.5% and the between assay variation from 11.3% to 26.1%.¹⁵ Plasma insulin was measured by radioimmunoassay as described previously.¹⁷

STATISTICAL ANALYSIS

Results are expressed as mean (SEM). The integrated incremental PP secretion in response to CCK-33 infusion was determined by calculating the area under the plasma concentration-time curve after subtraction of the basal value. Gall bladder volumes are expressed in cm³ or as percentages, considering the gall bladder volume at the start of the infusion of CCK-33 as 100% (zero minutes). Data were analysed for the significance of differences by analysis of variance (ANOVA). When this indicated a probability of less than 0.05 for the null hypothesis Student-Newman-Keuls analyses were performed to determine which values differed significantly. The significance level was set at $p < 0.05$.

Results

SERUM GLUCOSE CONCENTRATIONS

Fasting serum glucose concentrations in the normo and hyperglycaemic experiments were 5.1 (0.3) mmol/l, 5.0 (0.7) mmol/l, and 5.0 (0.3) mmol/l respectively. Figure 1 shows mean serum glucose concentrations reached during clamping.

PLASMA INSULIN CONCENTRATIONS

Fasting plasma insulin concentrations were not significantly different between the three experiments (Fig 2). In the normoglycaemic experiment no significant changes in plasma insulin were found during infusion of CCK-33. After initiating the hyperglycaemic clamp at -45 minutes, plasma insulin concentrations increased significantly ($p < 0.05$) starting from -30 minutes in the 8 and 15 mmol/l glucose clamps. At the end of the 15 mmol/l glucose clamp plasma insulin concentrations of 212 (45) mU/l were reached.

GALL BLADDER VOLUMES

Basal gall bladder volumes were not significantly different between the three experiments (28 (8) cm³ (normoglycaemia), 29 (9) cm³ (8 mmol/l hyperglycaemia), and 29 (9) cm³ (15 mmol/l




Figure 3: Gall bladder volumes (% mean (SEM)) during normoglycaemia and hyperglycaemia clamping. From 0 to 90 minutes gall bladder volumes were in response to cholecystokinin (CCK-33) infusion.

hyperglycaemia)). Infusion of 0.25, 0.5, and 1.0 IDU kg⁻¹.h⁻¹ CCK-33 resulted in significant reductions in gall bladder volumes from 100% to 70% (3%) (p<0.05), 34% (7%) (p<0.05), and 8% (3%) (p<0.05) during normoglycaemia, from 100% to 84% (3%) (p<0.05), 45% (4%) (p<0.05), and 10% (2%) (p<0.05) during the 8 mmol/l glucose clamp, and from 100% to 91% (4%) (p<0.05), 63% (5%) (p<0.05), and 24% (5%) (p<0.05) during the 15 mmol/l glucose clamp (Fig 3). During the 15 mmol/l hyperglycaemia gall bladder contraction was significantly reduced (p<0.05) from 30 to 90 minutes compared with normoglycaemia and from 45 to 90 minutes compared with the 8 mmol/l glucose clamp experiment. During the 8 mmol/l hyperglycaemia gall bladder contraction was significantly reduced at 30, 45, and 60 minutes compared with normoglycaemia.

PLASMA CCK CONCENTRATIONS

Fasting plasma CCK concentrations were not significantly different between the three experiments (Table). Infusion of 0.25, 0.5, and 1.0 IDU kg⁻¹.h⁻¹ CCK-33 resulted in significant (p<0.05) increases in plasma CCK concentrations over baseline in all experiments. The plasma CCK concentrations in response to infusion of graded doses of CCK-33 were not significantly different between the three experiments.

Plasma cholecystokinin (CCK) concentrations (pmol/l, mean (SEM)) before and during infusion of stepwise increasing doses of CCK-33

CCK-33 infusion rate (IDU kg ⁻¹ .h ⁻¹)	Normoglycaemia	Hyperglycaemia (8 mmol/l)	Hyperglycaemia (15 mmol/l)
Baseline	3.4 (0.3)	3.5 (0.4)	3.5 (0.3)
0.25	4.5 (0.2)*	4.4 (0.3)*	4.5 (0.2)*
0.5	6.5 (0.4)*	6.6 (0.3)*	5.6 (0.3)*
1.0	10.9 (0.7)*	11.0 (0.8)*	10.2 (0.6)*

*p<0.05 compared with baseline.




Figure 4: Integrated incremental pancreatic polypeptide (PP) secretion (pmol/l, mean (SEM)) response to cholecystokinin (CCK-33) infusion during normoglycaemia and hyperglycaemic clamping.

PLASMA PANCREATIC POLYPEPTIDE CONCENTRATIONS

Fasting plasma PP concentrations were not significantly different between the three experiments (17 (2) pmol/l (normoglycaemia), 17 (2) pmol/l (8 mmol/l hyperglycaemia), and 19 (2) pmol/l (15 mmol/l hyperglycaemia)). From -45 to zero minutes no significant changes in PP concentration were found during normoglycaemia whereas after initiating hyperglycaemia, plasma PP concentrations decreased significantly (p<0.05) both during the 8 and 15 mmol/l glucose clamp. At 0 minutes mean plasma PP concentrations were 17 (2) pmol/l (normoglycaemia), 13 (1) pmol/l (8 mmol/l hyperglycaemia), and 12 (1) pmol/l (15 mmol/l hyperglycaemia). Infusion of CCK-33 from 0.25 IDU kg⁻¹.h⁻¹ to 1.0 IDU kg⁻¹.h⁻¹ resulted in significant (p<0.05) increases in plasma PP in the normoglycaemic and 8 mmol/l hyperglycaemic experiments whereas during the 15 mmol/l hyperglycaemic experiment plasma PP concentrations increased significantly only during infusion of 1.0 IDU kg⁻¹.h⁻¹ CCK-33. Compared with normoglycaemia, during the 15 mmol/l hyperglycaemia the integrated incremental PP secretion in response to CCK-33 was significantly reduced during infusion of 0.25, 0.5, and 1.0 IDU kg⁻¹.h⁻¹ CCK-33 (Fig 4). During the 8 mmol/l hyperglycaemia the integrated PP secretion in response to CCK-33 was significantly (p<0.05) reduced compared with normoglycaemia during infusion of 0.25 and 0.5 IDU kg⁻¹.h⁻¹.

Discussion

The results of this study indicate that in healthy humans, gall bladder motility induced by CCK-33 is related to blood glucose concentrations. During an acute hyperglycaemia of 8 and 15 mmol/l the gall bladder contraction in response to CCK-33 was inhibited in a dose dependent manner.

In several studies it has been suggested that gastrointestinal function is related to blood glucose concentrations.¹⁻⁶ During an acute

hyperglycaemia induced by intravenous infusion of glucose, gastric emptying,¹³ gastric acid secretion⁴⁵ and pancreatic enzyme secretion⁴⁶ are reduced in healthy subjects. McGregor *et al* have shown that dextrose given intravenously inhibits meal stimulated intraduodenal bilirubin output.⁴ It is not known from that study, however, whether the reduction in intraduodenal bilirubin output resulted from hyperglycaemia induced alterations in gastric emptying of the meal, alterations in endogenous CCK secretion, or alterations in the responsiveness of the gall bladder to CCK. Unfortunately, in several of the studies that analysed the effect of intravenous glucose on gastrointestinal function, serum glucose concentrations have not been reported.^{13 46} Also, serum glucose concentrations may vary considerably during a constant infusion of glucose due to insulin secretion. To accurately determine the effect of hyperglycaemia on gastrointestinal function we think it is essential to stabilise serum glucose concentrations as in the present study by glucose clamp techniques.

In recent years it has been possible to study the mechanisms involved in the regulation postprandial gall bladder motility in detail because of the development of sensitive and specific radioimmunoassays for measurement of CCK and the availability of ultrasonography and cholescintigraphy as non-invasive methods to measure human gall bladder motility.^{9 10 15 18 19} Two findings point to an important role of CCK in the regulation of postprandial gall bladder motility. Firstly, after eating a meal gall bladder contraction is closely related to plasma CCK concentrations.^{18 20} Secondly, intravenous infusion of CCK to plasma concentrations comparable with those found after meals induces similar gall bladder contraction.^{13 18 19} By stimulating gall bladder contraction in our study through infusion of CCK-33 instead of meal ingestion we precluded the influences of alterations in gastric emptying and endogenous secretion of CCK on gall bladder motility induced by hyperglycaemia. Infusion of stepwise increasing doses of CCK-33 in our experiments resulted in dose dependent increments in plasma CCK, corresponding to plasma CCK concentrations found after ingestion of a light, normal, and fatty meal.^{18 20} Plasma CCK concentrations reached during infusion of CCK-33 were in the same range in the three experiments, thus excluding the possibility that differences in gall bladder contraction between the hyperglycaemic and normoglycaemic experiments are related to differences in plasma CCK concentrations.

The mechanism responsible for the inhibitory action of hyperglycaemia on gall bladder motility is not known. It is possible that metabolic, hormonal, or neural changes secondary to hyperglycaemia, rather than glucose itself, modulate gastrointestinal function. For instance, high glucose concentrations stimulate insulin secretion in healthy volunteers and hyperinsulinemia affects gastrointestinal function. Previous studies have shown, however, that insulin exerts a stimulatory, rather than an inhibitory effect on intestinal motility.^{21 22} Furthermore, in type I diabetic patients, hyperglycaemia reduces

gastric emptying indicating that insulin is not responsible for the inhibitory effect of hyperglycaemia on gastrointestinal function.²³

Several arguments are in favour of an inhibitory effect of hyperglycaemia on gall bladder motility by suppressing activity of the vagal cholinergic system. Firstly, the effect of CCK on the gall bladder in vivo is dependent on vagal cholinergic activity. Cholinergic blockade with atropine reduces contractile responses of the gall bladder to CCK,²⁴ and action similar to the effect of hyperglycaemia on gall bladder contraction induced by CCK. Secondly, it has been shown in animals that infusion of glucose suppresses vagal nerve activity.²⁵ Thirdly, in our study hyperglycaemia significantly reduced basal and CCK-33 stimulated PP concentrations. The release of PP from the pancreas is known to be under cholinergic control.⁷ Although CCK is a physiological stimulus of PP secretion in humans the effect of CCK on PP secretion is abolished by atropine, indicating that the cholinergic system regulates CCK induced PP secretion.^{7 26} Fourthly, hyperglycaemia is known to suppress PP secretion through inhibition of cholinergic activity as the inhibitory effect of hyperglycaemia on PP secretion is blunted after truncal vagotomy.²⁷ Based on these findings it is postulated that hyperglycaemia affects gall bladder motility, at least in part, through inhibition of cholinergic activity.

It is not known whether in patients with diabetes mellitus, as in healthy subjects, the gall bladder sensitivity to CCK is influenced by serum glucose concentrations. A reduced gall bladder emptying in response to a meal and to infusion of CCK-33 has been reported in such patients.²⁸ This has been associated with the presence of autonomic neuropathy as in diabetic patients without neuropathy the gall bladder contraction in response to CCK is not different from that in healthy subjects. No significant correlation has been found between impaired gall bladder emptying and overall diabetic control as determined by the percentage of total haemoglobin present as glycated haemoglobin.²⁸ It remains to be established whether in diabetic patients gall bladder motility is dependent on actual serum glucose concentrations.

In conclusion blood glucose concentrations affect gall bladder motility in healthy subjects, an acute hyperglycaemia at 8 and 15 mmol/l dose dependently reduces the gall bladder sensitivity to CCK-33, and hyperglycaemia reduces basal and CCK-33 stimulated plasma PP concentrations, suggesting impaired cholinergic activity during hyperglycaemia.

- 1 McGregor IL, Gueller R, Watts HD, Meyer JH. The effect of acute hyperglycemia on gastric emptying in man. *Gastroenterology* 1976; 70: 190-6.
- 2 Barnett JL, Owyang C. Serum glucose concentration as a modulator of interdigestive gastric motility. *Gastroenterology* 1988; 94: 739-44.
- 3 Oster-Jorgensen E, Pedersen SA, Larsen ML. The influence of induced hyperglycemia on gastric emptying in healthy humans. *Scand J Clin Lab Invest* 1990; 50: 831-6.
- 4 McGregor IL, Deveney C, Way LW, Meyer JH. The effect of acute hyperglycemia on meal-stimulated gastric, biliary, and pancreatic secretion, and serum gastrin. *Gastroenterology* 1976; 70: 197-202.
- 5 Loud FB, Holst JJ, Rehfeld JF, Christiansen J. Inhibition of

- gastric acid secretion in humans by glucagon during euglycemia, hyperglycemia, and hypoglycemia. *Dig Dis Sci* 1988; **33**: 530-4.
- 6 Niederau C, Sonnenberg A, Erckenbrecht J. Effects of intravenous infusion of amino acids, fat or glucose on unstimulated pancreatic secretion in healthy humans. *Dig Dis Sci* 1985; **30**: 445-55.
 - 7 Schwartz TW. Pancreatic polypeptide: a hormone under vagal control. *Gastroenterology* 1983; **85**: 1411-25.
 - 8 Marco J, Hedo JA, Castillo-Olivares J, Villanueva ML. Human pancreatic polypeptide secretion in conditions of exogenous and endogenous hyperglycemia. *Diabetologia* 1980; **19**: 418-22.
 - 9 Everson GT, Braverman DJ, Johnson ML, Kern F. A critical evaluation of real-time ultrasonography for the study of gallbladder volume and contraction. *Gastroenterology* 1980; **79**: 40-6.
 - 10 Hopman WPM, Brouwer WFM, Rosenbusch G, et al. A computerized method for rapid quantification of gallbladder volume from real-time sonograms. *Radiology* 1985; **154**: 236-7.
 - 11 Ward WK, Halter JB, Beard JC, Poter D. Adaptation of B and A cell function during prolonged glucose infusion in human subjects. *Am J Physiol* 1984; **246**: E405-11.
 - 12 DeFronzo RA, Tobin JD, Andres R. Glucose clamp technique for quantifying insulin secretion and resistance. *Am J Physiol* 1979; **237**: E214-23.
 - 13 Hopman WPM, Kerstens PJSM, Jansen JBMJ, Rosenbusch J, Lamers CBHW. Effect of graded physiologic doses of cholecystokinin on gallbladder contraction measured by ultrasonography. *Gastroenterology* 1985; **89**: 1242-7.
 - 14 Lamers CBHW, Diemel JM, van Leer E, van Leusen R, Peetoom J. Mechanism of elevated serum pancreatic polypeptide concentration in chronic renal failure. *J Clin Endocrinol Metab* 1982; **55**: 922-6.
 - 15 Jansen JBMJ, Lamers CBHW. Radioimmunoassay of cholecystokinin in human tissue and plasma. *Clin Chim Acta* 1983; **131**: 305-16.
 - 16 Jansen JBMJ, Lamers CBHW. Molecular forms of cholecystokinin in plasma from normal and gastrectomized human subjects following a fat meal. *Peptides* 1987; **8**: 801-5.
 - 17 Nijs HGT, Radder JK, Fröhlich M, Krans HMJ. In vivo relation between insulin clearance and insulin action in healthy subjects and insulin-dependent diabetic patients. *Diabetes* 1990; **39**: 333-9.
 - 18 Wiener I, Inoue K, Fagan CJ, Lilja P, Watson LC, Thompson JC. Release of cholecystokinin in man: correlation of blood levels with gallbladder contraction. *Ann Surg* 1981; **194**: 321-7.
 - 19 Masclee AAM, Hopman WPM, Corstens FHM, Rosenbusch G, Jansen JBMJ, Lamers CBHW. Simultaneous measurement of gallbladder emptying with cholecintigraphy and ultrasonography during infusion of physiologic doses of cholecystokinin: a comparison. *Radiology* 1989; **173**: 407-10.
 - 20 Hopman WPM, Rosenbusch G, Jansen JBMJ, de Jong AJL, Lamers CBHW. Gallbladder contraction: effects of fatty meals and of cholecystokinin. *Radiology* 1985; **157**: 37-9.
 - 21 Bueno L, Ruckebusch M. Insulin and jejunal electrical activity in dog and sheep. *Am J Physiol* 1976; **230**: 1538-44.
 - 22 Prasad KR, Sarna SK. The central and peripheral effects of insulin on migrating myoelectric complexes [Abstract]. *Gastroenterology* 1986; **90**: 1589.
 - 23 Fraser RJ, Horowitz M, Maddox AF, Harving PE, Chatterton BE, Dent J. Hyperglycemia slows gastric emptying in type I (insulin-dependent) diabetes mellitus. *Diabetologia* 1990; **33**: 675-80.
 - 24 Hopman WPM, Jansen JBMJ, Rosenbusch G, Lamers CBHW. Role of cholecystokinin and the cholinergic system in intestinal stimulation of gallbladder contraction in man. *Hepatology* 1990; **11**: 261-5.
 - 25 Hirano T, Niiijima A. Effects of 2-deoxy-D-glucose, glucose and insulin on efferent activity in gastric vagus nerve. *Experientia* 1980; **36**: 1197-8.
 - 26 Meier R, Hildebrand P, Thumshirn M, et al. Effect of loxiglumide, a cholecystokinin antagonist, on pancreatic polypeptide release in humans. *Gastroenterology* 1990; **99**: 1757-62.
 - 27 Tsuda K, Seino Y, Mori K, et al. Effect of truncal vagotomy on pancreatic polypeptide response after intravenous glucose administration. *Regul Pept* 1981; **1**: 347-52.
 - 28 Stone BG, Gavalier JS, Belle SH, et al. Impairment of gallbladder emptying in diabetes mellitus. *Gastroenterology* 1988; **95**: 170-6.